

1969 AB

1. B
2. C
3. B
4. D
5. E
6. B
7. D
8. B
9. C
10. E
11. B
12. A
13. C
14. E
15. B
16. B
17. B
18. E
19. C
20. A
21. B
22. E
23. C

24. C
25. A
26. C
27. C
28. C
29. A
30. E
31. C
32. B
33. A
34. D
35. A
36. B
37. D
38. C
39. D
40. E
41. D
42. D
43. D
44. C
45. D

1969 BC

1. C
2. E
3. B
4. D
5. E
6. B
7. D
8. C
9. D
10. A
11. B
12. E
13. C
14. D
15. B
16. B
17. B
18. E
19. C
20. A
21. B
22. E
23. D

24. C
25. A
26. C
27. C
28. D
29. C
30. D
31. C
32. B
33. A
34. D
35. A
36. B
37. D
38. A
39. D
40. E
41. D
42. B
43. E
44. E
45. E

1973 AB

1. E
2. E
3. B
4. A
5. A
6. D
7. B
8. B
9. A
10. C
11. B
12. C
13. D
14. D
15. C
16. C
17. C
18. D
19. D
20. D
21. B
22. B
23. C

24. B
25. B
26. E
27. E
28. C
29. C
30. B
31. D
32. D
33. A
34. C
35. C
36. A
37. A
38. B
39. B
40. E
41. D
42. D
43. E
44. B
45. C

1973 BC

1. A
2. D
3. A
4. C
5. B
6. D
7. D
8. B
9. A
10. A
11. E
12. D
13. D
14. A
15. C
16. A
17. C
18. D
19. D
20. E
21. B
22. C
23. C

24. A
25. B
26. D
27. E
28. C
29. A
30. B
31. E
32. C
33. A
34. C
35. C
36. E
37. E
38. B
39. D
40. C
41. D
42. D
43. E
44. A
45. E

1985 AB

1. D
2. E
3. A
4. C
5. D
6. C
7. E
8. B
9. D
10. D
11. B
12. C
13. A
14. D
15. C
16. B
17. C
18. C
19. B
20. A
21. B
22. A
23. B

24. D
25. E
26. E
27. D
28. C
29. D
30. B
31. C
32. D
33. B
34. A
35. D
36. B
37. D
38. C
39. E
40. D
41. E
42. C
43. B
44. A
45. A

1985 BC

1. D
2. A
3. B
4. D
5. D
6. E
7. A
8. C
9. B
10. A
11. A
12. A
13. B
14. C
15. C
16. C
17. B
18. C
19. D
20. C
21. B
22. A
23. C

24. D
25. C
26. E
27. E
28. E
29. D
30. B
31. D
32. E
33. C
34. A
35. B
36. E
37. A
38. C
39. A
40. A
41. C
42. E
43. E
44. A
45. D

1988 AB

1. C
2. D
3. A
4. E
5. A
6. D
7. D
8. B
9. E
10. C
11. A
12. B
13. A
14. D
15. B
16. C
17. D
18. E
19. B
20. C
21. C
22. C
23. B

24. C
25. B
26. E
27. E
28. C
29. B
30. A
31. C
32. A
33. A
34. D
35. B
36. C
37. D
38. E
39. E
40. B
41. A
42. C
43. B
44. C
45. D

1988 BC

1. A
2. D
3. B
4. E
5. C
6. C
7. A
8. A
9. D
10. D
11. A
12. B
13. B
14. A
15. E
16. A
17. D
18. E
19. B
20. E
21. D
22. E
23. E

24. D
25. D
26. C
27. B
28. E
29. B
30. C
31. C
32. E
33. E
34. C
35. A
36. E or D
37. D
38. C
39. C
40. E
41. B
42. A
43. A
44. A
45. B

1993 AB

1. C
2. B
3. D
4. A
5. A
6. D
7. B
8. E
9. E
10. D
11. C
12. B
13. A
14. A
15. D
16. B
17. E
18. D
19. E
20. B
21. C
22. E
23. C

24. A
25. C
26. D
27. C
28. B
29. C
30. C
31. E
32. A
33. B
34. D
35. E
36. D
37. C
38. A
39. D
40. C
41. D
42. B
43. B
44. C
45. B

1993 BC

1. A
2. C
3. E
4. B
5. D
6. A
7. A
8. B
9. D
10. E
11. E
12. E
13. C
14. B
15. D
16. A
17. A
18. B
19. B
20. E
21. A
22. B
23. D

24. C
25. D
26. B
27. C
28. A
29. E
30. C
31. A
32. B
33. A
34. E
35. A
36. E
37. B
38. C
39. C
40. C
41. C
42. E
43. A
44. E
45. D

1997 AB

1. C
2. A
3. C
4. D
5. E
6. C
7. D
8. C
9. B
10. E
11. E
12. B
13. A
14. C
15. B
16. D
17. A
18. C
19. D
20. E

21. E
22. D
23. A
24. B
25. A
76. E
77. D
78. D
79. C
80. A
81. A
82. B
83. C
84. C
85. C
86. A
87. B
88. E
89. B
90. D

1997 BC

1. C
2. E
3. A
4. C
5. C
6. A
7. C
8. E
9. A
10. B
11. C
12. A
13. B
14. C
15. D
16. B
17. B
18. C
19. D
20. E

21. A
22. C
23. E
24. D
25. A
76. D
77. E
78. A
79. D
80. B
81. D
82. B
83. E
84. C
85. D
86. A
87. B
88. C
89. D
90. B

1998 AB

1. D
2. B
3. C
4. B
5. E
6. A
7. E
8. E
9. D
10. D
11. A
12. E
13. B
14. C
15. D
16. E
17. D
18. B
19. C
20. A
21. B
22. C
23. A
24. D
25. D
26. A
27. A
28. E
76. A
77. C
78. B
79. A
80. B
81. D
82. E
83. B
84. A
85. C
86. C
87. D
88. C
89. B
90. D
91. E
92. D

1998 BC

1. C
2. A
3. D
4. A
5. A
6. E
7. E
8. B
9. D
10. E
11. A
12. E
13. B
14. E
15. B
16. C
17. D
18. B
19. D
20. E
21. C
22. A
23. E
24. C
25. C
26. E
27. D
28. C
76. D
77. E
78. B
79. A
80. B
81. B
82. B
83. C
84. B
85. C
86. C
87. D
88. C
89. A
90. A
91. E
92. D

1. B Sine is the only odd function listed. $\sin(-x) = -\sin(x)$.

2. C $\ln t < 0$ for $0 < t < 1 \Rightarrow \ln(x-2) < 0$ for $2 < x < 3$.

3. B Need to have $\lim_{x \rightarrow 2} f(x) = f(2) = k$.

$$\begin{aligned} k &= \lim_{x \rightarrow 2} \frac{\sqrt{2x+5} - \sqrt{x+7}}{x-2} = \lim_{x \rightarrow 2} \frac{\sqrt{2x+5} - \sqrt{x+7}}{x-2} \cdot \frac{\sqrt{2x+5} + \sqrt{x+7}}{\sqrt{2x+5} + \sqrt{x+7}} \\ &= \lim_{x \rightarrow 2} \frac{2x+5 - (x+7)}{x-2} \cdot \frac{1}{\sqrt{2x+5} + \sqrt{x+7}} = \lim_{x \rightarrow 2} \frac{1}{\sqrt{2x+5} + \sqrt{x+7}} = \frac{1}{6} \end{aligned}$$

4. D $\int_0^8 \frac{dx}{\sqrt{1+x}} = 2\sqrt{1+x} \Big|_0^8 = 2(3-1) = 4$

5. E Using implicit differentiation, $6x + 2xy' + 2y + 2y \cdot y' = 0$. Therefore $y' = \frac{-2y-6x}{2x+2y}$.

When $x=1$, $3+2y+y^2 = 2 \Rightarrow 0 = y^2 + 2y + 1 = (y+1)^2 \Rightarrow y = -1$

Therefore $2x+2y = 0$ and so $\frac{dy}{dx}$ is not defined at $x=1$.

6. B This is the derivative of $f(x) = 8x^8$ at $x = \frac{1}{2}$

$$f'\left(\frac{1}{2}\right) = 64\left(\frac{1}{2}\right)^7 = \frac{1}{2}$$

7. D With $f(x) = x + \frac{k}{x}$, we need $0 = f'(-2) = 1 - \frac{k}{4}$ and so $k = 4$. Since $f''(-2) < 0$ for $k = 4$, f does have a relative maximum at $x = -2$.

8. B $p(x) = q(x)(x-1) + 12$ for some polynomial $q(x)$ and so $12 = p(1) = (1+2)(1+k) \Rightarrow k = 3$

9. C $A = \pi r^2$, $\frac{dA}{dt} = 2\pi r \cdot \frac{dr}{dt}$ and from the given information in the problem $\frac{dA}{dt} = 2 \frac{dr}{dt}$.

$$\text{So, } 2 \frac{dr}{dt} = 2\pi r \cdot \frac{dr}{dt} \Rightarrow r = \frac{1}{\pi}$$

10. E $x = e^y \Rightarrow y = \ln x$

11. B Let L be the distance from $\left(x, -\frac{x^2}{2}\right)$ and $\left(0, -\frac{1}{2}\right)$.

$$L^2 = (x-0)^2 + \left(\frac{x^2}{2} - \frac{1}{2}\right)^2$$

$$2L \cdot \frac{dL}{dx} = 2x + 2\left(\frac{x^2}{2} - \frac{1}{2}\right)(x)$$

$$\frac{dL}{dx} = \frac{2x + 2\left(\frac{x^2}{2} - \frac{1}{2}\right)(x)}{2L} = \frac{2x + x^3 - x}{2L} = \frac{x^3 + x}{2L} = \frac{x(x^2 + 1)}{2L}$$

$\frac{dL}{dx} < 0$ for all $x < 0$ and $\frac{dL}{dx} > 0$ for all $x > 0$, so the minimum distance occurs at $x = 0$.

The nearest point is the origin.

12. A $\frac{4}{2x-1} = 2\left(\frac{4}{x-1}\right) \Rightarrow x-1 = 4x-2; x = \frac{1}{3}$

13. C $\int_{-\pi/2}^k \cos x \, dx = 3 \int_k^{\pi/2} \cos x \, dx; \sin k - \sin\left(-\frac{\pi}{2}\right) = 3\left(\sin \frac{\pi}{2} - \sin k\right)$
 $\sin k + 1 = 3 - 3 \sin k; 4 \sin k = 2 \Rightarrow k = \frac{\pi}{6}$

14. E $y = x^5 - 1$ has an inverse $x = y^5 - 1 \Rightarrow y = \sqrt[5]{x+1}$

15. B The graphs do not need to intersect (eg. $f(x) = -e^{-x}$ and $g(x) = e^{-x}$). The graphs could intersect (e.g. $f(x) = 2x$ and $g(x) = x$). However, if they do intersect, they will intersect no more than once because $f(x)$ grows faster than $g(x)$.

16. B $y' > 0 \Rightarrow y$ is increasing; $y'' < 0 \Rightarrow$ the graph is concave down. Only B meets these conditions.

17. B $y' = 20x^3 - 5x^4, y'' = 60x^2 - 20x^3 = 20x^2(3-x)$. The only sign change in y'' is at $x = 3$. The only point of inflection is $(3, 162)$.

18. E There is no derivative at the vertex which is located at $x = 3$.

19. C $\frac{dv}{dt} = \frac{1 - \ln t}{t^2} > 0$ for $0 < t < e$ and $\frac{dv}{dt} < 0$ for $t > e$, thus v has its maximum at $t = e$.

20. A $y(0) = 0$ and $y'(0) = \frac{1/2}{\sqrt{1 - \frac{x^2}{4}}}\bigg|_{x=0} = \frac{1}{\sqrt{4 - x^2}}\bigg|_{x=0} = \frac{1}{2}$. The tangent line is
 $y = \frac{1}{2}x \Rightarrow x - 2y = 0$.

21. B $f'(x) = 2x - 2e^{-2x}$, $f'(0) = -2$, so f is decreasing

22. E $\ln e^{2x} = 2x \Rightarrow \frac{d}{dx}(\ln e^{2x}) = \frac{d}{dx}(2x) = 2$

23. C $\int_0^2 e^{2x} dx = \frac{1}{2} e^{2x} \bigg|_0^2 = \frac{1}{2}(e^4 - 1)$

24. C $y = \ln \sin x$, $y' = \frac{\cos x}{\sin x} = \cot x$

25. A $\int_m^{2m} \frac{1}{x} dx = \ln x \bigg|_m^{2m} = \ln(2m) - \ln(m) = \ln 2$ so the area is independent of m .

26. C $\int_0^1 \sqrt{x^2 - 2x + 1} dx = \int_0^1 |x - 1| dx = \int_0^1 -(x - 1) dx = -\frac{1}{2}(x - 1)^2 \bigg|_0^1 = \frac{1}{2}$

Alternatively, the graph of the region is a right triangle with vertices at $(0,0)$, $(0,1)$, and $(1,0)$.

The area is $\frac{1}{2}$.

27. C $\int \tan x dx = \int \frac{\sin x}{\cos x} dx = -\ln |\cos x| + C = \ln |\sec x| + C$

28. C $\sqrt{3} \cos x + 3 \sin x$ can be thought of as the expansion of $\sin(x + y)$. Since $\sqrt{3}$ and 3 are too large for values of $\sin y$ and $\cos y$, multiply and divide by the result of the Pythagorean Theorem used on those values, i.e. $2\sqrt{3}$. Then

$$\begin{aligned}\sqrt{3} \cos x + 3 \sin x &= 2\sqrt{3} \left(\frac{\sqrt{3}}{2\sqrt{3}} \cos x + \frac{3}{2\sqrt{3}} \sin x \right) = 2\sqrt{3} \left(\frac{1}{2} \cos x + \frac{\sqrt{3}}{2} \sin x \right) \\ &= 2\sqrt{3} (\sin y \cos x + \cos y \sin x) = 2\sqrt{3} \sin(y + x)\end{aligned}$$

where $y = \sin^{-1}\left(\frac{1}{2}\right)$. The amplitude is $2\sqrt{3}$.

Alternatively, the function $f(x)$ is periodic with period 2π . $f'(x) = -\sqrt{3} \sin x + 3 \cos x = 0$ when $\tan x = \sqrt{3}$. The solutions over one period are $x = \frac{\pi}{3}, \frac{4\pi}{3}$. Then $f\left(\frac{\pi}{3}\right) = 2\sqrt{3}$ and $f\left(\frac{4\pi}{3}\right) = -2\sqrt{3}$. So the amplitude is $2\sqrt{3}$.

29. A $\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin x} dx = \ln(\sin x) \Big|_{\pi/4}^{\pi/2} = \ln 1 - \ln \frac{1}{\sqrt{2}} = \ln \sqrt{2}$
30. E Because f is continuous for all x , the Intermediate Value Theorem implies that the graph of f must intersect the x -axis. The graph must also intersect the y -axis since f is defined for all x , in particular, at $x = 0$.
31. C $\frac{dy}{dx} = -y \Rightarrow y = ce^{-x}$ and $1 = ce^{-1} \Rightarrow c = e$; $y = e \cdot e^{-x} = e^{1-x}$
32. B If $a < 0$ then $\lim_{x \rightarrow -\infty} y = \infty$ and $\lim_{x \rightarrow \infty} y = -\infty$ which would mean that there is at least one root. If $a > 0$ then $\lim_{x \rightarrow -\infty} y = -\infty$ and $\lim_{x \rightarrow \infty} y = \infty$ which would mean that there is at least one root. In both cases the equation has at least one root.
33. A $\frac{1}{3} \int_{-1}^2 3t^3 - t^2 dt = \frac{1}{3} \left(\frac{3}{4} t^4 - \frac{1}{3} t^3 \right) \Big|_{-1}^2 = \frac{1}{3} \left(\left(12 - \frac{8}{3} \right) - \left(\frac{3}{4} + \frac{1}{3} \right) \right) = \frac{11}{4}$
34. D $y' = -\frac{1}{x^2}$, so the desired curve satisfies $y' = x^2 \Rightarrow y = \frac{1}{3} x^3 + C$

35. A $a(t) = 24t^2$, $v(t) = 8t^3 + C$ and $v(0) = 0 \Rightarrow C = 0$. The particle is always moving to the right, so distance $= \int_0^2 8t^3 dt = 2t^4 \Big|_0^2 = 32$.
36. B $y = \sqrt{4 + \sin x}$, $y(0) = 2$, $y'(0) = \frac{\cos 0}{2\sqrt{4 + \sin 0}} = \frac{1}{4}$. The linear approximation to y is $L(x) = 2 + \frac{1}{4}x$. $L(1.2) = 2 + \frac{1}{4}(1.2) = 2.03$
37. D All options have the same value at $x = 0$. We want the one that has the same first and second derivatives at $x = 0$ as $y = \cos 2x$: $y'(0) = -2 \sin 2x \Big|_{x=0} = 0$ and $y''(0) = -4 \cos 2x \Big|_{x=0} = -4$.
For $y = 1 - 2x^2$, $y'(0) = -4x \Big|_{x=0} = 0$ and $y''(0) = -4$ and no other option works.
38. C $\int \frac{x^2}{e^{-x^3}} dx = -\frac{1}{3} \int e^{-x^3} (-3x^2 dx) = -\frac{1}{3} e^{-x^3} + C = -\frac{1}{3e^{-x^3}} + C$
39. D $x = e \Rightarrow v = 1$, $u = 0$, $y = 0$; $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dx} = (\sec^2 u) \left(1 + \frac{1}{v^2}\right) \left(\frac{1}{x}\right) = (1)(2)(e^{-1}) = \frac{2}{e}$
40. E One solution technique is to evaluate each integral and note that the value is $\frac{1}{n+1}$ for each.
Another technique is to use the substitution $u = 1 - x$; $\int_0^1 (1-x)^n dx = \int_1^0 u^n (-du) = \int_0^1 u^n du$.
Integrals do not depend on the variable that is used and so $\int_0^1 u^n du$ is the same as $\int_0^1 x^n dx$.
41. D $\int_{-1}^3 f(x) dx = \int_{-1}^2 (8-x^2) dx + \int_2^3 x^2 dx = \left(8x - \frac{1}{3}x^3\right) \Big|_{-1}^2 + \frac{1}{3}x^3 \Big|_2^3 = 27 \frac{1}{3}$
42. D $y = x^3 - 3x^2 + k$, $y' = 3x^2 - 6x = 3x(x-2)$. So f has a relative maximum at $(0, k)$ and a relative minimum at $(2, k-4)$. There will be 3 distinct x -intercepts if the maximum and minimum are on the opposite sides of the x -axis. We want $k-4 < 0 < k \Rightarrow 0 < k < 4$.
43. D $\int \sin(2x+3) dx = -\frac{1}{2} \cos(2x+3) + C$

44. C Since $\cos 2A = 2 \cos^2 A - 1$, we have $3 - 2 \cos^2 \frac{\pi x}{3} = 3 - (1 + \cos \frac{2\pi x}{3})$ and the latter expression has period $\frac{2\pi}{\left(\frac{2\pi}{3}\right)} = 3$

45. D Let $y = f(x^3)$. We want y'' where $f'(x) = g(x)$ and $f''(x) = g'(x) = f'(x^2)$

$$y = f(x^3)$$

$$y' = f'(x^3) \cdot 3x^2$$

$$y'' = 3x^2 \left(f''(x^3) \cdot 3x^2 \right) + f'(x^3) \cdot 6x$$

$$= 9x^4 f''(x^3) + 6x f'(x^3) = 9x^4 f'((x^3)^2) + 6x g(x^3) = 9x^4 f'(x^6) + 6x g(x^3)$$

1. C For horizontal asymptotes consider the limit as $x \rightarrow \pm\infty$: $t \rightarrow 0 \Rightarrow y = 0$ is an asymptote
For vertical asymptotes consider the limit as $y \rightarrow \pm\infty$: $t \rightarrow -1 \Rightarrow x = -1$ is an asymptote

2. E $y = (x+1)\tan^{-1}x$, $y' = \frac{x+1}{1+x^2} + \tan^{-1}x$

$$y'' = \frac{(1+x^2)(1) - (x+1)(2x)}{(1+x^2)^2} + \frac{1}{1+x^2} = \frac{2-2x}{(1+x^2)^2}$$

y'' changes sign at $x = 1$ only. The point of inflection is $(1, \pi/2)$

3. B $y = \sqrt{x}$, $y' = \frac{1}{2\sqrt{x}}$. By the Mean Value Theorem we have $\frac{1}{2\sqrt{c}} = \frac{2}{4} \Rightarrow c = 1$.

The point is (1,1).

4. D $\int_0^8 \frac{dx}{\sqrt{1+x}} = 2\sqrt{1+x} \Big|_0^8 = 2(3-1) = 4$

5. E Using implicit differentiation, $6x + 2xy' + 2y + 2y \cdot y' = 0$. Therefore $y' = \frac{-2y-6x}{2x+2y}$.

When $x = 1$, $3 + 2y + y^2 = 2 \Rightarrow 0 = y^2 + 2y + 1 = (y+1)^2 \Rightarrow y = -1$

Therefore $2x + 2y = 0$ and so $\frac{dy}{dx}$ is not defined at $x = 1$.

6. B This is the derivative of $f(x) = 8x^8$ at $x = \frac{1}{2}$.

$$f'\left(\frac{1}{2}\right) = 64\left(\frac{1}{2}\right)^7 = \frac{1}{2}$$

7. D With $f(x) = x + \frac{k}{x}$, we need $0 = f'(-2) = 1 - \frac{k}{4}$ and so $k = 4$. Since $f''(-2) < 0$ for $k = 4$, f does have a relative maximum at $x = -2$.

8. C $h'(x) = 2f(x) \cdot f'(x) - 2g(x) \cdot g'(x) = 2f(x) \cdot (-g(x)) - 2g(x) \cdot f(x) = -4f(x) \cdot g(x)$

9. D $A = \frac{1}{2} \int_0^{2\pi} (\sqrt{3 + \cos \theta})^2 d\theta = 2 \cdot \frac{1}{2} \int_0^{\pi} (\sqrt{3 + \cos \theta})^2 d\theta = \int_0^{\pi} (3 + \cos \theta) d\theta$

10. A $\int_0^1 \frac{x^2}{x^2 + 1} dx = \int_0^1 \frac{x^2 + 1 - 1}{x^2 + 1} dx = \int_0^1 \left(\frac{x^2 + 1}{x^2 + 1} - \frac{1}{x^2 + 1} \right) dx = \left(x - \tan^{-1} x \right) \Big|_0^1 = 1 - \frac{\pi}{4} = \frac{4 - \pi}{4}$

11. B Let L be the distance from $\left(x, -\frac{x^2}{2} \right)$ and $\left(0, -\frac{1}{2} \right)$.

$$L^2 = (x - 0)^2 + \left(\frac{x^2}{2} - \frac{1}{2} \right)^2$$

$$2L \cdot \frac{dL}{dx} = 2x + 2 \left(\frac{x^2}{2} - \frac{1}{2} \right) (x)$$

$$\frac{dL}{dx} = \frac{2x + 2 \left(\frac{x^2}{2} - \frac{1}{2} \right) (x)}{2L} = \frac{2x + x^3 - x}{2L} = \frac{x^3 + x}{2L} = \frac{x(x^2 + 1)}{2L}$$

$\frac{dL}{dx} < 0$ for all $x < 0$ and $\frac{dL}{dx} > 0$ for all $x > 0$, so the minimum distance occurs at $x = 0$.

The nearest point is the origin.

12. E By the Fundamental Theorem of Calculus, if $F(x) = \int_0^x e^{-t^2} dt$ then $F'(x) = e^{-x^2}$.

13. C $\int_{-\pi/2}^k \cos x dx = 3 \int_k^{\pi/2} \cos x dx$; $\sin k - \sin \left(-\frac{\pi}{2} \right) = 3 \left(\sin \frac{\pi}{2} - \sin k \right)$

$$\sin k + 1 = 3 - 3 \sin k; 4 \sin k = 2 \Rightarrow k = \frac{\pi}{6}$$

14. D $y = x^2 + 2$ and $u = 2x - 1$, $\frac{dy}{du} = \frac{dy}{dx} \cdot \frac{dx}{du} = (2x) \left(\frac{1}{2} \right) = x$

15. B The graphs do not need to intersect (eg. $f(x) = -e^{-x}$ and $g(x) = e^{-x}$). The graphs could intersect (e.g. $f(x) = 2x$ and $g(x) = x$). However, if they do intersect, they will intersect no more than once because $f(x)$ grows faster than $g(x)$.

16. B $y' > 0 \Rightarrow y$ is increasing; $y'' < 0 \Rightarrow$ the graph is concave down. Only B meets these conditions.
17. B $y' = 20x^3 - 5x^4$, $y'' = 60x^2 - 20x^3 = 20x^2(3 - x)$. The only sign change in y'' is at $x = 3$. The only point of inflection is $(3, 162)$.
18. E There is no derivative at the vertex which is located at $x = 3$.
19. C $\frac{dv}{dt} = \frac{1 - \ln t}{t^2} > 0$ for $0 < t < e$ and $\frac{dv}{dt} < 0$ for $t > e$, thus v has its maximum at $t = e$.
20. A $y(0) = 0$ and $y'(0) = \left. \frac{\frac{1/2}{\sqrt{1 - \frac{x^2}{4}}}}{\sqrt{1 - \frac{x^2}{4}}} \right|_{x=0} = \left. \frac{1}{\sqrt{4 - x^2}} \right|_{x=0} = \frac{1}{2}$. The tangent line is $y = \frac{1}{2}x \Rightarrow x - 2y = 0$.
21. B $f'(x) = 2x - 2e^{-2x}$, $f'(0) = -2$, so f is decreasing
22. E $f(x) = \int_0^x \frac{1}{\sqrt{t^3 + 2}} dt$, $f(-1) = \int_0^{-1} \frac{1}{\sqrt{t^3 + 2}} dt = -\int_{-1}^0 \frac{1}{\sqrt{t^3 + 2}} dt < 0$
 $f(-1) < 0$ so E is false.
23. D $\frac{dy}{dx} = \frac{-xe^{-x^2}}{y} \Rightarrow 2y dy = -2xe^{-x^2} dx \Rightarrow y^2 = e^{-x^2} + C$
 $4 = 1 + C \Rightarrow C = 3$; $y^2 = e^{-x^2} + 3 \Rightarrow y = \sqrt{e^{-x^2} + 3}$
24. C $y = \ln \sin x$, $y' = \frac{\cos x}{\sin x} = \cot x$
25. A $\int_m^{2m} \frac{1}{x} dx = \ln x \Big|_m^{2m} = \ln(2m) - \ln(m) = \ln 2$ so the area is independent of m .

$$26. \quad C \quad \int_0^1 \sqrt{x^2 - 2x + 1} \, dx = \int_0^1 |x - 1| \, dx = \int_0^1 -(x - 1) \, dx = -\frac{1}{2}(x - 1)^2 \Big|_0^1 = \frac{1}{2}$$

Alternatively, the graph of the region is a right triangle with vertices at (0,0), (0,1), and (1,0).

The area is $\frac{1}{2}$.

$$27. \quad C \quad \int \tan x \, dx = \int \frac{\sin x}{\cos x} \, dx = -\ln |\cos x| + C = \ln |\sec x| + C$$

$$28. \quad D \quad \text{Use L'Hôpital's Rule: } \lim_{x \rightarrow 0} \frac{e^{2x} - 1}{\tan x} = \lim_{x \rightarrow 0} \frac{2e^{2x}}{\sec^2 x} = 2$$

29. C Make the substitution $x = 2 \sin \theta \Rightarrow dx = 2 \cos \theta \, d\theta$.

$$\int_0^1 (4 - x^2)^{-3/2} \, dx = \int_0^{\pi/6} \frac{2 \cos \theta}{8 \cos^3 \theta} \, d\theta = \frac{1}{4} \int_0^{\pi/6} \sec^2 \theta \, d\theta = \frac{1}{4} \tan \theta \Big|_0^{\pi/6} = \frac{1}{4} \cdot \frac{\sqrt{3}}{3} = \frac{\sqrt{3}}{12}$$

$$30. \quad D \quad \text{Substitute } -x \text{ for } x \text{ in } \sum_{n=0}^{\infty} \frac{x^n}{n!} = e^x \text{ to get } \sum_{n=0}^{\infty} \frac{(-1)^n x^n}{n!} = e^{-x}$$

$$31. \quad C \quad \frac{dy}{dx} = -y \Rightarrow y = ce^{-x} \text{ and } 1 = ce^{-1} \Rightarrow c = e; y = e \cdot e^{-x} = e^{1-x}$$

32. B $1 + 2^x + 3^x + 4^x + \cdots + n^x + \cdots = \sum_{n=1}^{\infty} \frac{1}{n^p}$ where $p = -x$. This is a p -series and is convergent if $p > 1 \Rightarrow -x > 1 \Rightarrow x < -1$.

$$33. \quad A \quad \frac{1}{3} \int_{-1}^2 3t^3 - t^2 \, dt = \frac{1}{3} \left(\frac{3}{4} t^4 - \frac{1}{3} t^3 \right) \Big|_{-1}^2 = \frac{1}{3} \left(\left(12 - \frac{8}{3} \right) - \left(\frac{3}{4} + \frac{1}{3} \right) \right) = \frac{11}{4}$$

$$34. \quad D \quad y' = -\frac{1}{x^2}, \text{ so the desired curve satisfies } y' = x^2 \Rightarrow y = \frac{1}{3} x^3 + C$$

35. A $a(t) = 24t^2$, $v(t) = 8t^3 + C$ and $v(0) = 0 \Rightarrow C = 0$. The particle is always moving to the right, so distance $= \int_0^2 8t^3 \, dt = 2t^4 \Big|_0^2 = 32$.

36. B $y = \sqrt{4 + \sin x}$, $y(0) = 2$, $y'(0) = \frac{\cos 0}{2\sqrt{4 + \sin 0}} = \frac{1}{4}$. The linear approximation to y is

$$L(x) = 2 + \frac{1}{4}x. L(1.2) = 2 + \frac{1}{4}(1.2) = 2.03$$

37. D This item uses the formal definition of a limit and is no longer part of the AP Course Description. Need to have $|(1-3x) - (-5)| < \varepsilon$ whenever $0 < |x-2| < \delta$.
 $|(1-3x) - (-5)| = |6-3x| = 3|x-2| < \varepsilon$ if $|x-2| < \varepsilon/3$.
 Thus we can use any $\delta < \varepsilon/3$. Of the five choices, the largest satisfying this condition is $\delta = \varepsilon/4$.

38. A Note $f(1) = \frac{1}{2}$. Take the natural logarithm of each side of the equation and then differentiate.

$$\ln f(x) = (2-3x)\ln(x^2+1); \frac{f'(x)}{f(x)} = (2-3x) \cdot \frac{2x}{x^2+1} - 3\ln(x^2+1)$$

$$f'(1) = f(1) \left((-1) \cdot \frac{2}{2} - 3\ln(2) \right) \Rightarrow f'(1) = \frac{1}{2}(-1 - 3\ln 2) = -\frac{1}{2}(\ln e + \ln 2^3) = -\frac{1}{2}\ln 8e$$

39. D $x = e \Rightarrow v = 1$, $u = 0$, $y = 0$; $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dv} \cdot \frac{dv}{dx} = (\sec^2 u) \left(1 + \frac{1}{v^2} \right) \left(\frac{1}{x} \right) = (1)(2)(e^{-1}) = \frac{2}{e}$

40. E One solution technique is to evaluate each integral and note that the value is $\frac{1}{n+1}$ for each.

$$\text{Another technique is to use the substitution } u = 1-x; \int_0^1 (1-x)^n dx = \int_1^0 u^n (-du) = \int_0^1 u^n du.$$

$$\text{Integrals do not depend on the variable that is used and so } \int_0^1 u^n du \text{ is the same as } \int_0^1 x^n dx.$$

41. D $\int_{-1}^3 f(x) dx = \int_{-1}^2 (8-x^2) dx + \int_2^3 x^2 dx = \left(8x - \frac{1}{3}x^3 \right) \Big|_{-1}^2 + \frac{1}{3}x^3 \Big|_2^3 = 27 \frac{1}{3}$

42. B Use the technique of antiderivatives by parts to evaluate $\int x^2 \cos x \, dx$

$$u = x^2 \quad dv = \cos x \, dx$$

$$du = 2x \, dx \quad v = \sin x$$

$$f(x) - \int 2x \sin x \, dx = \int x^2 \cos x \, dx = x^2 \sin x - \int 2x \sin x \, dx + C$$

$$f(x) = x^2 \sin x + C$$

43. E
$$L = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} \, dx = \int_a^b \sqrt{1 + (\sec^2 x)^2} \, dx = \int_a^b \sqrt{1 + \sec^4 x} \, dx$$

44. E $y'' - y' - 2y = 0$, $y'(0) = -2$, $y(0) = 2$; the characteristic equation is $r^2 - r - 2 = 0$.

The solutions are $r = -1$, $r = 2$ so the general solution to the differential equation is

$$y = c_1 e^{-x} + c_2 e^{2x} \quad \text{with} \quad y' = -c_1 e^{-x} + 2c_2 e^{2x}. \quad \text{Using the initial conditions we have the system:}$$

$$2 = c_1 + c_2 \quad \text{and} \quad -2 = -c_1 + 2c_2 \Rightarrow c_2 = 0, \quad c_1 = 2. \quad \text{The solution is} \quad f(x) = 2e^{-x} \Rightarrow f(1) = 2e^{-1}.$$

45. E The ratio test shows that the series is convergent for any value of x that makes $|x+1| < 1$. The solutions to $|x+1| = 1$ are the endpoints of the interval of convergence. Test $x = -2$ and $x = 0$ in the series. The resulting series are $\sum_{k=1}^{\infty} \frac{(-1)^k}{k^2}$ and $\sum_{k=1}^{\infty} \frac{1}{k^2}$ which are both convergent. The interval is $-2 \leq x \leq 0$.


1. E $\int (x^3 - 3x) dx = \frac{1}{4}x^4 - \frac{3}{2}x^2 + C$
2. E $g(x) = 5 \Rightarrow g(f(x)) = 5$
3. B $y = \ln x^2$; $y' = \frac{2x}{x^2} = \frac{2}{x}$. At $x = e^2$, $y' = \frac{2}{e^2}$.
4. A $f(x) = x + \sin x$; $f'(x) = 1 - \cos x$
5. A $\lim_{x \rightarrow -\infty} e^x = 0 \Rightarrow y = 0$ is a horizontal asymptote
6. D $f'(x) = \frac{(1)(x+1) - (x-1)(1)}{(x+1)^2}$, $f'(1) = \frac{2}{4} = \frac{1}{2}$
7. B Replace x with $(-x)$ and see if the result is the opposite of the original. This is true for B.
 $-(-x)^5 + 3(-x) = x^5 - 3x = -(-x^5 + 3x)$.
8. B Distance $= \int_1^2 |t^2| dx = \int_1^2 t^2 dt = \frac{1}{3}t^3 \Big|_1^2 = \frac{1}{3}(2^3 - 1^3) = \frac{7}{3}$
9. A $y' = 2 \cos 3x \cdot \frac{d}{dx}(\cos 3x) = 2 \cos 3x \cdot (-\sin 3x) \cdot \frac{d}{dx}(3x) = 2 \cos 3x \cdot (-\sin 3x) \cdot (3)$
 $y' = -6 \sin 3x \cos 3x$
10. C $f(x) = \frac{x^4}{3} - \frac{x^5}{5}$; $f'(x) = \frac{4x^3}{3} - x^4$; $f''(x) = 4x^2 - 4x^3 = 4x^2(1-x)$
 $f'' > 0$ for $x < 1$ and $f'' < 0$ for $x > 1 \Rightarrow f'$ has its maximum at $x = 1$.
11. B Curve and line have the same slope when $3x^2 = \frac{3}{4} \Rightarrow x = \frac{1}{2}$. Using the line, the point of tangency is $\left(\frac{1}{2}, \frac{3}{8}\right)$. Since the point is also on the curve, $\frac{3}{8} = \left(\frac{1}{2}\right)^3 + k \Rightarrow k = \frac{1}{4}$.
12. C Substitute the points into the equation and solve the resulting linear system.
 $3 = 16 + 4A + 2B - 5$ and $-37 = -16 + 4A - 2B - 5$; $A = -3$, $B = 2 \Rightarrow A + B = -1$.

13. D $v(t) = 8t - 3t^2 + C$ and $v(1) = 25 \Rightarrow C = 20$ so $v(t) = 8t - 3t^2 + 20$.
 $s(4) - s(2) = \int_2^4 v(t) dt = (4t^2 - t^3 + 20t) \Big|_2^4 = 32$
14. D $f(x) = x^{1/3}(x-2)^{2/3}$
 $f'(x) = x^{1/3} \cdot \frac{2}{3}(x-2)^{-1/3} + (x-2)^{2/3} \cdot \frac{1}{3}x^{-2/3} = \frac{1}{3}x^{-2/3}(x-2)^{-1/3}(3x-2)$
 f' is not defined at $x = 0$ and at $x = 2$.
15. C $\text{Area} = \int_0^2 e^{\frac{x}{2}} dx = 2e^{\frac{x}{2}} \Big|_0^2 = 2(e-1)$
16. C $\frac{dN}{dt} = 3000e^{\frac{2}{5}t}$, $N = 7500e^{\frac{2}{5}t} + C$ and $N(0) = 7500 \Rightarrow C = 0$
 $N = 7500e^{\frac{2}{5}t}$, $N(5) = 7500e^2$
17. C Determine where the curves intersect. $-x^2 + x + 6 = 4 \Rightarrow x^2 - x - 2 = 0$
 $(x-2)(x+1) = 0 \Rightarrow x = -1, x = 2$. Between these two x values the parabola lies above the line $y = 4$.
 $\text{Area} = \int_{-1}^2 ((-x^2 + x + 6) - 4) dx = \left(-\frac{1}{3}x^3 + \frac{1}{2}x^2 + 2x \right) \Big|_{-1}^2 = \frac{9}{2}$
18. D $\frac{d}{dx}(\arcsin 2x) = \frac{1}{\sqrt{1-(2x)^2}} \cdot \frac{d}{dx}(2x) = \frac{2}{\sqrt{1-(2x)^2}} = \frac{2}{\sqrt{1-4x^2}}$
19. D If f is strictly increasing then it must be one to one and therefore have an inverse.
20. D By the Fundamental Theorem of Calculus, $\int_a^b f(x) dx = F(b) - F(a)$ where $F'(x) = f(x)$.
21. B $\int_0^1 (x+1)e^{x^2+2x} dx = \frac{1}{2} \int_0^1 e^{x^2+2x} ((2x+2) dx) = \frac{1}{2} (e^{x^2+2x}) \Big|_0^1 = \frac{1}{2} (e^3 - e^0) = \frac{e^3 - 1}{2}$
22. B $f(x) = 3x^5 - 20x^3$; $f'(x) = 15x^4 - 60x^2$; $f''(x) = 60x^3 - 120x = 60x(x^2 - 2)$
The graph of f is concave up where $f'' > 0$: $f'' > 0$ for $x > \sqrt{2}$ and for $-\sqrt{2} < x < 0$.

23. C $\lim_{h \rightarrow 0} \frac{\ln(2+h) - \ln 2}{h} = f'(2)$ where $f(x) = \ln x$; $f'(x) = \frac{1}{x} \Rightarrow f'(2) = \frac{1}{2}$
24. B $f(x) = \cos(\arctan x)$; $-\frac{\pi}{2} < \arctan x < \frac{\pi}{2}$ and the cosine in this domain takes on all values in the interval $(0, 1]$.
25. B $\int_0^{\frac{\pi}{4}} \tan^2 x \, dx = \int_0^{\frac{\pi}{4}} (\sec^2 x - 1) \, dx = (\tan x - x) \Big|_0^{\frac{\pi}{4}} = 1 - \frac{\pi}{4}$
26. E $\frac{dV}{dt} = 4\pi r^2 \cdot \frac{dr}{dt} = S \cdot \frac{dr}{dt} = 100\pi(0.3) = 30\pi$
27. E $\int_0^{\frac{1}{2}} \frac{2x}{\sqrt{1-x^2}} \, dx = -\int_0^{\frac{1}{2}} (1-x^2)^{-\frac{1}{2}} (-2x \, dx) = -2(1-x^2)^{\frac{1}{2}} \Big|_0^{\frac{1}{2}} = 2 - \sqrt{3}$
28. C $v(t) = 8 - 6t$ changes sign at $t = \frac{4}{3}$. Distance = $\left| x(1) - x\left(\frac{4}{3}\right) \right| + \left| x(2) - x\left(\frac{4}{3}\right) \right| = \frac{5}{3}$.
- Alternative Solution: Distance = $\int_1^2 |v(t)| \, dt = \int_1^2 |8 - 6t| \, dt = \frac{5}{3}$
29. C $-1 \leq \sin x \leq 1 \Rightarrow -\frac{3}{2} \leq \sin x - \frac{1}{2} \leq \frac{1}{2}$; The maximum for $\left| \sin x - \frac{1}{2} \right|$ is $\frac{3}{2}$.
30. B $\int_1^2 \frac{x-4}{x^2} \, dx = \int_1^2 \left(\frac{1}{x} - 4x^{-2} \right) \, dx = \left(\ln x + \frac{4}{x} \right) \Big|_1^2 = (\ln 2 + 2) - (\ln 1 + 4) = \ln 2 - 2$
31. D $\log_a(2^a) = \frac{a}{4} \Rightarrow \log_a 2 = \frac{1}{4} \Rightarrow 2 = a^{\frac{1}{4}}; a = 16$
32. D $\int \frac{5}{1+x^2} \, dx = 5 \int \frac{1}{1+x^2} \, dx = 5 \tan^{-1}(x) + C$
33. A $f(-x) = -f(x) \Rightarrow f'(-x) \cdot (-1) = -f'(x) \Rightarrow f'(-x) = -f'(x)$ thus $f'(-x_0) = -f'(x_0)$.

34. C $\frac{1}{2} \int_0^2 \sqrt{x} dx = \frac{1}{2} \cdot \frac{2}{3} x^{\frac{3}{2}} \Big|_0^2 = \frac{1}{3} \cdot 2^{\frac{3}{2}} = \frac{2}{3} \sqrt{2}$

35. C Washers: $\sum \pi r^2 \Delta x$ where $r = y = \sec x$.
 Volume = $\pi \int_0^{\frac{\pi}{4}} \sec^2 x dx = \pi \tan x \Big|_0^{\frac{\pi}{4}} = \pi(\tan \frac{\pi}{4} - \tan 0) = \pi$



36. A $y = e^{nx}$, $y' = ne^{nx}$, $y'' = n^2 e^{nx}$, ..., $y^{(n)} = n^n e^{nx}$

37. A $\frac{dy}{dx} = 4y$, $y(0) = 4$. This is exponential growth. The general solution is $y = Ce^{4x}$. Since $y(0) = 4$, $C = 4$ and so the solution is $y = 4e^{4x}$.

38. B Let $z = x - c$. Then $5 = \int_1^2 f(x - c) dx = \int_{1-c}^{2-c} f(z) dz$

39. B Use the distance formula to determine the distance, L , from any point $(x, y) = (x, \frac{1}{2}x^2)$ on the curve to the point $(4, 1)$. The distance L satisfies the equation $L^2 = (x - 4)^2 + (\frac{1}{2}x^2 - 1)^2$. Determine where L is a maximum by examining critical points. Differentiating with respect to x , $2L \cdot \frac{dL}{dx} = 2(x - 4) + 2(\frac{1}{2}x^2 - 1)x = x^3 - 8$. $\frac{dL}{dx}$ changes sign from positive to negative at $x = 2$ only. The point on the curve has coordinates $(2, 2)$.

40. E $\sec^2(xy) \cdot (xy' + y) = 1$, $xy' \sec^2(xy) + y \sec^2(xy) = 1$, $y' = \frac{1 - y \sec^2(xy)}{x \sec^2(xy)} = \frac{\cos^2(xy) - y}{x}$

41. D $\int_{-1}^1 f(x) dx = \int_{-1}^0 (x+1) dx + \int_0^1 \cos(\pi x) dx = \frac{1}{2}(x+1)^2 \Big|_{-1}^0 + \frac{1}{\pi} \sin(\pi x) \Big|_0^1$
 $= \frac{1}{2} + \frac{1}{\pi}(\sin \pi - \sin 0) = \frac{1}{2}$

42. D $\Delta x = \frac{1}{3}; T = \frac{1}{2} \cdot \frac{1}{3} \left(1^2 + 2 \left(\frac{4}{3} \right)^2 + 2 \left(\frac{5}{3} \right)^2 + 2^2 \right) = \frac{127}{54}$

43. E Solve $\frac{x}{2} = -1$ and $\frac{x}{2} = 2; x = -2, 4$

44. B Use the linearization of $f(x) = \sqrt[4]{x}$ at $x = 16$. $f'(x) = \frac{1}{4}x^{-\frac{3}{4}}, f'(16) = \frac{1}{32}$
 $L(x) = 2 + \frac{1}{32}(x-16); f(16+h) \approx L(16+h) = 2 + \frac{h}{32}$

45. C This uses the definition of continuity of f at $x = x_0$.

$$1. \quad A \quad f'(x) = e^{\frac{1}{x}} \cdot \frac{d\left(\frac{1}{x}\right)}{dx} = e^{\frac{1}{x}} \left(-\frac{1}{x^2}\right) = -\frac{e^{\frac{1}{x}}}{x^2}$$

$$2. \quad D \quad \int_0^3 (x+1)^{\frac{1}{2}} dx = \frac{2}{3} (x+1)^{\frac{3}{2}} \Big|_0^3 = \frac{2}{3} \left(4^{\frac{3}{2}} - 1^{\frac{3}{2}}\right) = \frac{2}{3} (8-1) = \frac{14}{3}$$

$$3. \quad A \quad f'(x) = 1 - \frac{1}{x^2} = \frac{(x+1)(x-1)}{x^2}. \quad f'(x) > 0 \text{ for } x < -1 \text{ and for } x > 1.$$

f is increasing for $x \leq -1$ and for $x \geq 1$.

4. C The slopes will be negative reciprocals at the point of intersection.

$$3x^2 = 3 \Rightarrow x = \pm 1 \text{ and } x \geq 0, \text{ thus } x = 1 \text{ and the } y \text{ values must be the same at } x = 1.$$

$$-\frac{1}{3} + b = 1 \Rightarrow b = \frac{4}{3}$$

$$5. \quad B \quad \int_{-1}^2 \frac{|x|}{x} dx = \int_{-1}^0 -1 dx + \int_0^2 1 dx = -1 + 2 = 1$$

$$6. \quad D \quad f'(x) = \frac{(1)(x+1) - (x-1)(1)}{(x+1)^2}, \quad f'(1) = \frac{2}{4} = \frac{1}{2}$$

$$7. \quad D \quad \frac{dy}{dx} = \frac{2x + 2y \cdot \frac{dy}{dx}}{x^2 + y^2} \text{ at } (1, 0) \Rightarrow y' = \frac{2}{1} = 2$$

$$8. \quad B \quad y = \sin x, \quad y' = \cos x, \quad y'' = -\sin x, \quad y''' = -\cos x, \quad y^{(4)} = \sin x$$

$$9. \quad A \quad y' = 2 \cos 3x \cdot \frac{d}{dx}(\cos 3x) = 2 \cos 3x \cdot (-\sin 3x) \cdot \frac{d}{dx}(3x) = 2 \cos 3x \cdot (-\sin 3x) \cdot 3$$

$$y' = -6 \sin 3x \cos 3x$$

$$10. \quad A \quad L = \int_0^b \sqrt{1+(y')^2} \, dx = \int_0^b \sqrt{1+\left(\frac{\sec x \tan x}{\sec x}\right)^2} \, dx$$

$$= \int_0^b \sqrt{1+(\tan x)^2} \, dx = \int_0^b \sqrt{\sec^2 x} \, dx = \int_0^b \sec x \, dx$$

$$11. \quad E \quad dy = \left(x \cdot \frac{1}{2} (1+x^2)^{-\frac{1}{2}} (2x) + (1+x^2)^{\frac{1}{2}} \right) dx; \quad dy = (0+1)(2) = 2$$

$$12. \quad D \quad \frac{1}{n} = \int_1^k x^{n-1} \, dx = \frac{x^n}{n} \Big|_1^k \Rightarrow \frac{1}{n} = \frac{k^n}{n} - \frac{1}{n}; \quad \frac{k^n}{n} = \frac{2}{n} \Rightarrow k = 2^{\frac{1}{n}}$$

$$13. \quad D \quad v(t) = 8t - 3t^2 + C \quad \text{and} \quad v(1) = 25 \Rightarrow C = 20 \quad \text{so} \quad v(t) = 8t - 3t^2 + 20.$$

$$s(4) - s(2) = \int_2^4 v(t) \, dt = \left(4t^2 - t^3 + 20t \right) \Big|_2^4 = 32$$

$$14. \quad A \quad \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{2e^t}{2t} = \frac{e^t}{t}$$

$$15. \quad C \quad \text{Area} = \int_0^2 e^{\frac{1}{2}x} \, dx = 2e^{\frac{1}{2}x} \Big|_0^2 = 2(e-1)$$

$$16. \quad A \quad \sin t = t - \frac{t^3}{3!} + \frac{t^5}{5!} - \frac{t^7}{7!} + \dots \Rightarrow \frac{\sin t}{t} = 1 - \frac{t^2}{3!} + \frac{t^4}{5!} - \frac{t^6}{7!} + \dots$$

$$17. \quad C \quad \frac{dN}{dt} = 3000e^{\frac{2}{5}t}, \quad N = 7500e^{\frac{2}{5}t} + C \quad \text{and} \quad N(0) = 7500 \Rightarrow C = 0$$

$$N = 7500e^{\frac{2}{5}t}, \quad N(5) = 7500e^2$$

18. D D could be false, consider $g(x) = 1 - x$ on $[0, 1]$. A is true by the Extreme Value Theorem, B is true because g is a function, C is true by the Intermediate Value Theorem, and E is true because g is continuous.

19. D I is a convergent p -series, $p = 2 > 1$

II is the Harmonic series and is known to be divergent,

III is convergent by the Alternating Series Test.

$$20. E \quad \int x\sqrt{4-x^2} dx = -\frac{1}{2} \int (4-x^2)^{\frac{1}{2}} (-2x dx) = -\frac{1}{2} \cdot \frac{2}{3} (4-x^2)^{\frac{3}{2}} + C = -\frac{1}{3} (4-x^2)^{\frac{3}{2}} + C$$

$$21. B \quad \int_0^1 (x+1)e^{x^2+2x} dx = \frac{1}{2} \int_0^1 e^{x^2+2x} ((2x+2) dx) = \frac{1}{2} \left(e^{x^2+2x} \right) \Big|_0^1 = \frac{1}{2} (e^3 - e^0) = \frac{e^3 - 1}{2}$$

$$22. C \quad x'(t) = t+1 \Rightarrow x(t) = \frac{1}{2}(t+1)^2 + C \quad \text{and} \quad x(0) = 1 \Rightarrow C = \frac{1}{2} \Rightarrow x(t) = \frac{1}{2}(t+1)^2 + \frac{1}{2}$$

$$x(1) = \frac{5}{2}, \quad y(1) = \ln \frac{5}{2}; \quad \left(\frac{5}{2}, \ln \frac{5}{2} \right)$$

$$23. C \quad \lim_{h \rightarrow 0} \frac{\ln(2+h) - \ln 2}{h} = f'(2) \quad \text{where} \quad f(x) = \ln x; \quad f'(x) = \frac{1}{x} \Rightarrow f'(2) = \frac{1}{2}$$

24. A This item uses the formal definition of a limit and is no longer part of the AP Course Description. $|f(x) - 7| = |(3x+1) - 7| = |3x-6| = 3|x-2| < \varepsilon$ whenever $|x-2| < \frac{\varepsilon}{3}$.

Any $\delta < \frac{\varepsilon}{3}$ will be sufficient and $\frac{\varepsilon}{4} < \frac{\varepsilon}{3}$, thus the answer is $\frac{\varepsilon}{4}$.


$$25. B \quad \int_0^{\frac{\pi}{4}} \tan^2 x dx = \int_0^{\frac{\pi}{4}} (\sec^2 x - 1) dx = (\tan x - x) \Big|_0^{\pi/4} = 1 - \frac{\pi}{4}$$

26. D For x in the interval $(-1, 1)$, $g(x) = |x^2 - 1| = -(x^2 - 1)$ and so $y = \ln g(x) = \ln(-(x^2 - 1))$.


Therefore

$$y' = \frac{2x}{x^2 - 1}, \quad y'' = \frac{(x^2 - 1)(2) - (2x)(2x)}{(x^2 - 1)^2} = \frac{-2x^2 - 2}{(x^2 - 1)^2} < 0$$

Alternative graphical solution: Consider the graphs of $g(x) = |x^2 - 1|$ and $\ln g(x)$.



$$g(x) = |x^2 - 1|$$



$$\ln|x^2 - 1|$$

concave
down

27. E $f'(x) = x^2 - 8x + 12 = (x - 2)(x - 6)$; the candidates are: $x = 0, 2, 6, 9$

x	0	2	6	9
$f(x)$	-5	$17/3$	-5	22

the maximum is at $x = 9$

28. C $x = \sin^2 y \Rightarrow dx = 2 \sin y \cos y dy$; when $x = 0$, $y = 0$ and when $x = \frac{1}{2}$, $y = \frac{\pi}{4}$

$$\int_0^{\frac{1}{2}} \frac{\sqrt{x}}{\sqrt{1-x}} dx = \int_0^{\frac{\pi}{4}} \frac{\sin y}{\sqrt{1-\sin^2 y}} \cdot 2 \sin y \cos y dy = \int_0^{\frac{\pi}{4}} 2 \sin^2 y dy$$

29. A Let $z = y'$. Then $z = e$ when $x = 0$. Thus $y'' = 2y' \Rightarrow z' = 2z$. Solve this differential equation.

$$z = Ce^{2x}; e = Ce^0 \Rightarrow C = e \Rightarrow y' = z = e^{2x+1}. \text{ Solve this differential equation.}$$

$$y = \frac{1}{2}e^{2x+1} + K; e = \frac{1}{2}e^1 + K \Rightarrow K = \frac{1}{2}e; y = \frac{1}{2}e^{2x+1} + \frac{1}{2}e, y(1) = \frac{1}{2}e^3 + \frac{1}{2}e = \frac{1}{2}e(e^2 + 1)$$

Alternative Solution: $y'' = 2y' \Rightarrow y' = Ce^{2x} = e \cdot e^{2x}$. Therefore $y'(1) = e^3$.

$$y'(1) - y'(0) = \int_0^1 y''(x) dx = \int_0^1 2y'(x) dx = 2y(1) - 2y(0) \text{ and so}$$

$$y(1) = \frac{y'(1) - y'(0) + 2y(0)}{2} = \frac{e^3 + e}{2}.$$

30. B $\int_1^2 \frac{x-4}{x^2} dx = \int_1^2 \left(\frac{1}{x} - 4x^{-2} \right) dx = \left(\ln x + \frac{4}{x} \right) \Big|_1^2 = (\ln 2 + 2) - (\ln 1 + 4) = \ln 2 - 2$

31. E $f'(x) = \frac{\frac{d}{dx}(\ln x)}{\ln x} = \frac{\frac{1}{x}}{\ln x} = \frac{1}{x \ln x}$

32. C Take the log of each side of the equation and differentiate. $\ln y = \ln x^{\ln x} = \ln x \cdot \ln x = (\ln x)^2$


$$\frac{y'}{y} = 2 \ln x \cdot \frac{d}{dx}(\ln x) = \frac{2}{x} \ln x \Rightarrow y' = x^{\ln x} \left(\frac{2}{x} \ln x \right)$$

33. A $f(-x) = -f(x) \Rightarrow f'(-x) \cdot (-1) = -f'(x) \Rightarrow f'(-x) = -f'(x)$ thus $f'(-x_0) = -f'(x_0)$.

34. C $\frac{1}{2} \int_0^2 \sqrt{x} dx = \frac{1}{2} \cdot \frac{2}{3} x^{\frac{3}{2}} \Big|_0^2 = \frac{1}{3} \cdot 2^{\frac{3}{2}} = \frac{2}{3} \sqrt{2}$

35. C Washers: $\sum \pi r^2 \Delta x$ where $r = y = \sec x$.

$$\text{Volume} = \pi \int_0^{\frac{\pi}{4}} \sec^2 x dx = \pi \tan x \Big|_0^{\frac{\pi}{4}} = \pi \left(\tan \frac{\pi}{4} - \tan 0 \right) = \pi$$



36. E $\int_0^1 \frac{x+1}{x^2+2x-3} dx = \frac{1}{2} \lim_{L \rightarrow 1^-} \int_0^L \frac{2x+2}{x^2+2x-3} dx = \frac{1}{2} \lim_{L \rightarrow 1^-} \ln |x^2+2x-3| \Big|_0^L$

$$= \frac{1}{2} \lim_{L \rightarrow 1^-} \left(\ln |L^2+2L-3| - \ln |-3| \right) = -\infty. \text{ Divergent}$$

37. E $\lim_{x \rightarrow 0} \frac{1 - \cos^2 2x}{x^2} = \lim_{x \rightarrow 0} \frac{\sin^2 2x}{x^2} = \lim_{x \rightarrow 0} \frac{\sin 2x}{2x} \cdot \frac{\sin 2x}{2x} \cdot 4 = 1 \cdot 1 \cdot 4 = 4$

38. B Let $z = x - c$. $5 = \int_1^2 f(x-c) dx = \int_{1-c}^{2-c} f(z) dz$

39. D $h'(x) = f'(g(x)) \cdot g'(x)$; $h'(1) = f'(g(1)) \cdot g'(1) = f'(2) \cdot g'(1) = (-4)(-3) = 12$

$$40. \quad C \quad \text{Area} = \frac{1}{2} \int_0^{2\pi} (1 - \cos \theta)^2 d\theta = \int_0^{\pi} (1 - 2 \cos \theta + \cos^2 \theta) d\theta; \quad \cos^2 \theta = \frac{1}{2}(1 + \cos 2\theta)$$

$$\text{Area} = \int_0^{\pi} \left(1 - 2 \cos \theta + \frac{1}{2}(1 + \cos 2\theta) \right) d\theta = \left(\frac{3}{2}\theta - 2 \sin \theta + \frac{1}{4} \sin 2\theta \right) \Big|_0^{\pi} = \frac{3}{2}\pi$$

$$41. \quad D \quad \int_{-1}^1 f(x) dx = \int_{-1}^0 (x+1) dx + \int_0^1 \cos(\pi x) dx$$

$$= \frac{1}{2}(x+1)^2 \Big|_{-1}^0 + \frac{1}{\pi} \sin(\pi x) \Big|_0^1 = \frac{1}{2} + \frac{1}{\pi}(\sin \pi - \sin 0) = \frac{1}{2}$$

$$42. \quad D \quad \Delta x = \frac{1}{3}; \quad T = \frac{1}{2} \cdot \frac{1}{3} \left(1^2 + 2 \left(\frac{4}{3} \right)^2 + 2 \left(\frac{5}{3} \right)^2 + 2^2 \right) = \frac{127}{54}$$

43. E Use the technique of antiderivatives by part:

$$u = \sin^{-1} x \quad dv = dx$$

$$du = \frac{dx}{\sqrt{1-x^2}} \quad v = x$$

$$\int \sin^{-1} x dx = x \sin^{-1} x - \int \frac{x}{\sqrt{1-x^2}} dx$$

44. A Multiply both sides of $x = x f'(x) - f(x)$ by $\frac{1}{x^2}$. Then $\frac{1}{x} = \frac{x f'(x) - f(x)}{x^2} = \frac{d}{dx} \left(\frac{f(x)}{x} \right)$.

Thus we have $\frac{f(x)}{x} = \ln|x| + C \Rightarrow f(x) = x(\ln|x| + C) = x(\ln|x| - 1)$ since $f(-1) = 1$.

$$\text{Therefore } f(e^{-1}) = e^{-1} (\ln|e^{-1}| - 1) = e^{-1}(-1 - 1) = -2e^{-1}$$

This was most likely the solution students were expected to produce while solving this problem on the 1973 multiple-choice exam. However, the problem itself is not well-defined. A solution to an initial value problem should be a function that is differentiable on an interval containing the initial point. In this problem that would be the domain $x < 0$ since the solution requires the choice of the branch of the logarithm function with $x < 0$. Thus one cannot ask about the value of the function at $x = e^{-1}$.

45. E $F'(x) = xg'(x)$ with $x \geq 0$ and $g'(x) < 0 \Rightarrow F'(x) < 0 \Rightarrow F$ is not increasing.


1. D $\int_1^2 x^{-3} dx = -\frac{1}{2}x^{-2} \Big|_1^2 = -\frac{1}{2}\left(\frac{1}{4}-1\right) = \frac{3}{8}$.
2. E $f'(x) = 4(2x+1)^3 \cdot 2$, $f''(1) = 4 \cdot 3(2x+1)^2 \cdot 2^2$, $f'''(1) = 4 \cdot 3 \cdot 2(2x+1)^1 \cdot 2^3$,
 $f^{(4)}(1) = 4! \cdot 2^4 = 384$
3. A $y = 3(4+x^2)^{-1}$ so $y' = -3(4+x^2)^{-2}(2x) = \frac{-6x}{(4+x^2)^2}$
 Or using the quotient rule directly gives $y' = \frac{(4+x^2)(0) - 3(2x)}{(4+x^2)^2} = \frac{-6x}{(4+x^2)^2}$
4. C $\int \cos(2x) dx = \frac{1}{2} \int \cos(2x)(2 dx) = \frac{1}{2} \sin(2x) + C$
5. D $\lim_{n \rightarrow \infty} \frac{4n^2}{n^2 + 10000n} = \lim_{n \rightarrow \infty} \frac{4}{1 + \frac{10000}{n}} = 4$
6. C $f'(x) = 1 \Rightarrow f'(5) = 1$
7. E $\int_1^4 \frac{1}{t} dt = \ln t \Big|_1^4 = \ln 4 - \ln 1 = \ln 4$
8. B $y = \ln\left(\frac{x}{2}\right) = \ln x - \ln 2$, $y' = \frac{1}{x}$, $y'(4) = \frac{1}{4}$
9. D Since e^{-x^2} is even, $\int_{-1}^0 e^{-x^2} dx = \frac{1}{2} \int_{-1}^1 e^{-x^2} dx = \frac{1}{2}k$
10. D $y' = 10^{(x^2-1)} \cdot \ln(10) \cdot \frac{d}{dx}\left((x^2-1)\right) = 2x \cdot 10^{(x^2-1)} \cdot \ln(10)$
11. B $v(t) = 2t + 4 \Rightarrow a(t) = 2 \therefore a(4) = 2$
12. C $f(g(x)) = \ln(g(x)^2) = \ln(x^2 + 4) \Rightarrow g(x) = \sqrt{x^2 + 4}$

13. A $2x + x \cdot y' + y + 3y^2 \cdot y' = 0 \Rightarrow y' = -\frac{2x + y}{x + 3y^2}$
14. D Since $v(t) \geq 0$, distance $= \int_0^4 |v(t)| dt = \int_0^4 \left(3t^{\frac{1}{2}} + 5t^{\frac{3}{2}} \right) dt = \left(2t^{\frac{3}{2}} + 2t^{\frac{5}{2}} \right) \Big|_0^4 = 80$
15. C $x^2 - 4 > 0 \Rightarrow |x| > 2$
16. B $f'(x) = 3x^2 - 6x = 3x(x - 2)$ changes sign from positive to negative only at $x = 0$.
17. C Use the technique of antiderivatives by parts:
 $u = x \quad dv = e^{-x} dx$
 $du = dx \quad v = -e^{-x}$
 $-xe^{-x} + \int e^{-x} dx = \left(-xe^{-x} - e^{-x} \right) \Big|_0^1 = 1 - 2e^{-1}$
18. C $y = \cos^2 x - \sin^2 x = \cos 2x, \quad y' = -2 \sin 2x$
19. B Quick solution: lines through the origin have this property.
 Or, $f(x_1) + f(x_2) = 2x_1 + 2x_2 = 2(x_1 + x_2) = f(x_1 + x_2)$
20. A $\frac{dy}{dx} = \frac{1}{1 + \cos^2 x} \cdot \frac{d}{dx}(\cos x) = \frac{-\sin x}{1 + \cos^2 x}$
21. B $|x| > 1 \Rightarrow x^2 > 1 \Rightarrow f(x) < 0$ for all x in the domain. $\lim_{|x| \rightarrow \infty} f(x) = 0$. $\lim_{|x| \rightarrow 1} f(x) = -\infty$. The only option that is consistent with these statements is (B).
22. A $\int_1^2 \frac{x^2 - 1}{x + 1} dx = \int_1^2 \frac{(x+1)(x-1)}{x+1} dx = \int_1^2 (x-1) dx = \frac{1}{2}(x-1)^2 \Big|_1^2 = \frac{1}{2}$
23. B $\frac{d}{dx} \left(x^{-3} - x^{-1} + x^2 \right) \Big|_{x=-1} = \left(-3x^{-4} + x^{-2} + 2x \right) \Big|_{x=-1} = -3 + 1 - 2 = -4$
24. D $16 = \int_{-2}^2 (x^7 + k) dx = \int_{-2}^2 x^7 dx + \int_{-2}^2 k dx = 0 + (2 - (-2))k = 4k \Rightarrow k = 4$

25. E $f'(e) = \lim_{h \rightarrow 0} \frac{f(e+h) - f(e)}{h} = \lim_{h \rightarrow 0} \frac{e^{e+h} - e^e}{h}$

26. E I: Replace y with $(-y)$: $(-y)^2 = x^2 + 9 \Rightarrow y^2 = x^2 + 9$, no change, so yes.
 II: Replace x with $(-x)$: $y^2 = (-x)^2 + 9 \Rightarrow y^2 = x^2 + 9$, no change, so yes.
 III: Since there is symmetry with respect to both axes there is origin symmetry.

27. D The graph is a V with vertex at $x = 1$. The integral gives the sum of the areas of the two triangles that the V forms with the horizontal axis for x from 0 to 3. These triangles have areas of $1/2$ and 2 respectively.



28. C Let $x(t) = -5t^2$ be the position at time t . Average velocity $= \frac{x(3) - x(0)}{3 - 0} = \frac{-45 - 0}{3} = -15$

29. D The tangent function is not defined at $x = \pi/2$ so it cannot be continuous for all real numbers. Option E is the only one that includes item III. In fact, the functions in I and II are a power and an exponential function that are known to be continuous for all real numbers x .

30. B $\int \tan(2x) dx = -\frac{1}{2} \int \frac{-2 \sin(2x)}{\cos(2x)} dx = -\frac{1}{2} \ln |\cos(2x)| + C$

31. C $V = \frac{1}{3} \pi r^2 h$, $\frac{dV}{dt} = \frac{1}{3} \pi \left(2rh \frac{dr}{dt} + r^2 \frac{dh}{dt} \right) = \frac{1}{3} \pi \left(2(6)(9) \left(\frac{1}{2} \right) + 6^2 \left(\frac{1}{2} \right) \right) = 24\pi$

32. D $\int_0^{\pi/3} \sin(3x) dx = -\frac{1}{3} \cos(3x) \Big|_0^{\pi/3} = -\frac{1}{3} (\cos \pi - \cos 0) = \frac{2}{3}$

33. B f' changes sign from positive to negative at $x = -1$ and therefore f changes from increasing to decreasing at $x = -1$.

Or f' changes sign from positive to negative at $x = -1$ and from negative to positive at $x = 1$. Therefore f has a local maximum at $x = -1$ and a local minimum at $x = 1$.

34. A $\int_0^1 ((x+8) - (x^3+8)) dx = \int_0^1 (x - x^3) dx = \left(\frac{1}{2} x^2 - \frac{1}{4} x^4 \right) \Big|_0^1 = \frac{1}{4}$

35. D The amplitude is 2 and the period is 2.

$$y = A \sin Bx \text{ where } |A| = \text{amplitude} = 2 \text{ and } B = \frac{2\pi}{\text{period}} = \frac{2\pi}{2} = \pi$$

36. B II is true since $|-7| = 7$ will be the maximum value of $|f(x)|$. To see why I and III do not

$$\text{have to be true, consider the following: } f(x) = \begin{cases} 5 & \text{if } x \leq -5 \\ -x & \text{if } -5 < x < 7 \\ -7 & \text{if } x \geq 7 \end{cases}$$

For $f(|x|)$, the maximum is 0 and the minimum is -7 .

37. D $\lim_{x \rightarrow 0} x \csc x = \lim_{x \rightarrow 0} \frac{x}{\sin x} = 1$

38. C To see why I and II do not have to be true consider $f(x) = \sin x$ and $g(x) = 1 + e^x$. Then $f(x) \leq g(x)$ but neither $f'(x) \leq g'(x)$ nor $f''(x) < g''(x)$ is true for all real values of x .

III is true, since

$$f(x) \leq g(x) \Rightarrow g(x) - f(x) \geq 0 \Rightarrow \int_0^1 (g(x) - f(x)) dx \geq 0 \Rightarrow \int_0^1 f(x) dx \leq \int_0^1 g(x) dx$$

39. E $f'(x) = \frac{1}{x} \cdot \frac{1}{x} - \frac{1}{x^2} \ln x = \frac{1}{x^2} (1 - \ln x) < 0$ for $x > e$. Hence f is decreasing for $x > e$.

40. D $\int_0^2 f(x) dx \leq \int_0^2 4 dx = 8$

41. E Consider the function whose graph is the horizontal line $y = 2$ with a hole at $x = a$. For this function $\lim_{x \rightarrow a} f(x) = 2$ and none of the given statements are true.


42. C This is a direct application of the Fundamental Theorem of Calculus: $f'(x) = \sqrt{1+x^2}$

43. B $y' = 3x^2 + 6x$, $y'' = 6x + 6 = 0$ for $x = -1$. $y'(-1) = -3$. Only option B has a slope of -3 .

44. A $\frac{1}{2} \int_0^2 x^2 (x^3 + 1)^{1/2} dx = \frac{1}{2} \cdot \frac{1}{3} \int_0^2 (x^3 + 1)^{1/2} (3x^2 dx) = \frac{1}{6} (x^3 + 1)^{3/2} \cdot \frac{2}{3} \Big|_0^2 = \frac{26}{9}$

45. A Washers: $\sum \pi(R^2 - r^2)\Delta y$ where $R = 2$, $r = x$

$$\text{Volume} = \pi \int_0^4 (2^2 - x^2) dy = \pi \int_0^4 (4 - y) dy = \pi \left(4y - \frac{1}{2}y^2 \right) \Big|_0^4 = 8\pi$$



1. D $\int_0^2 (4x^3 + 2) dx = (x^4 + 2x) \Big|_0^2 = (16 + 4) - (1 + 2) = 17$
2. A $f'(x) = 15x^4 - 15x^2 = 15x^2(x^2 - 1) = 15x^2(x-1)(x+1)$, changes sign from positive to negative only at $x = -1$. So f has a relative maximum at $x = -1$ only.
3. B $\int_1^2 \frac{x+1}{x^2+2x} dx = \frac{1}{2} \int_1^2 \frac{(2x+2) dx}{x^2+2x} = \frac{1}{2} \ln |x^2+2x| \Big|_1^2 = \frac{1}{2} (\ln 8 - \ln 3)$
4. D $x(t) = t^2 - 1 \Rightarrow \frac{dx}{dt} = 2t$ and $\frac{d^2x}{dt^2} = 2$; $y(t) = t^4 - 2t^3 \Rightarrow \frac{dy}{dt} = 4t^3 - 6t^2$ and $\frac{d^2y}{dt^2} = 12t^2 - 12t$
 $a(t) = \left(\frac{d^2x}{dt^2}, \frac{d^2y}{dt^2} \right) = (2, 12t^2 - 12t) \Rightarrow a(1) = (2, 0)$
5. D Area = $\int_{x_1}^{x_2} (\text{top curve} - \text{bottom curve}) dx$, $x_1 < x_2$; Area = $\int_{-1}^a (f(x) - g(x)) dx$
6. E $f(x) = \frac{x}{\tan x}$, $f'(x) = \frac{\tan x - x \sec^2 x}{\tan^2 x}$, $f'\left(\frac{\pi}{4}\right) = \frac{1 - \frac{\pi}{4} \cdot (\sqrt{2})^2}{1} = 1 - \frac{\pi}{2}$
7. A $\int \frac{du}{\sqrt{a^2 - u^2}} du = \sin^{-1}\left(\frac{u}{a}\right) \Rightarrow \int \frac{dx}{\sqrt{25 - x^2}} dx = \sin^{-1}\left(\frac{x}{5}\right) + C$
8. C $\lim_{x \rightarrow 2} \frac{f(x) - f(2)}{x - 2} = f'(2)$ so the derivative of f at $x = 2$ is 0.
9. B Take the derivative of each side of the equation with respect to x .
 $2xyy' + y^2 + 2xy' + 2y = 0$, substitute the point $(1, 2)$
 $(1)(4)y' + 2^2 + (2)(1)y' + (2)(2) = 0 \Rightarrow y = -\frac{4}{3}$
10. A Take the derivative of the general term with respect to x : $\sum_{n=1}^{\infty} (-1)^{n+1} x^{2n-2}$
11. A $\frac{d}{dx} \left(\ln \left(\frac{1}{1-x} \right) \right) = \frac{d}{dx} (-\ln(1-x)) = -\left(\frac{-1}{1-x} \right) = \frac{1}{1-x}$

12. A Use partial fractions to rewrite $\frac{1}{(x-1)(x+2)}$ as $\frac{1}{3}\left(\frac{1}{x-1} - \frac{1}{x+2}\right)$

$$\int \frac{1}{(x-1)(x+2)} dx = \frac{1}{3} \int \left(\frac{1}{x-1} - \frac{1}{x+2} \right) dx = \frac{1}{3} (\ln|x-1| - \ln|x+2|) + C = \frac{1}{3} \ln \left| \frac{x-1}{x+2} \right| + C$$

13. B $f(0) = 0$, $f(3) = 0$, $f'(x) = 3x^2 - 6x$; by the Mean Value Theorem,

$$f'(c) = \frac{f(3) - f(0)}{3} = 0 \text{ for } c \in (0, 3).$$

So, $0 = 3c^2 - 6c = 3c(c - 2)$. The only value in the open interval is 2.

14. C I. convergent: p -series with $p = 2 > 1$
 II. divergent: Harmonic series which is known to diverge
 III. convergent: Geometric with $|r| = \frac{1}{3} < 1$

15. C $x(t) = 4 + \int_0^t (2w - 4) dw = 4 + (w^2 - 4w) \Big|_0^t = 4 + t^2 - 4t = t^2 - 4t + 4$

or, $x(t) = t^2 - 4t + C$, $x(0) = 4 \Rightarrow C = 4$ so, $x(t) = t^2 - 4t + 4$

16. C For $f(x) = x^{\frac{1}{3}}$ we have continuity at $x = 0$, however, $f'(x) = \frac{1}{3}x^{-\frac{2}{3}}$ is not defined at $x = 0$.

17. B $f'(x) = (1) \cdot \ln(x^2) + x \cdot \frac{\frac{d}{dx}(x^2)}{x^2} = \ln(x^2) + \frac{2x^2}{x^2} = \ln(x^2) + 2$

18. C $\int \sin(2x+3) dx = \frac{1}{2} \int \sin(2x+3) (2dx) = -\frac{1}{2} \cos(2x+3) + C$

19. D $g(x) = e^{f(x)}$, $g'(x) = e^{f(x)} \cdot f'(x)$, $g''(x) = e^{f(x)} \cdot f''(x) + f'(x) \cdot e^{f(x)} \cdot f'(x)$
 $g''(x) = e^{f(x)} (f''(x) + (f'(x))^2) = h(x)e^{f(x)} \Rightarrow h(x) = f''(x) + (f'(x))^2$

20. C Look for concavity changes, there are 3.

21. B Use the technique of antiderivatives by parts:

$$u = f(x) \quad dv = \sin x \, dx$$

$$du = f'(x) \, dx \quad v = -\cos x$$

$$\int f(x) \sin x \, dx = -f(x) \cos x + \int f'(x) \cos x \, dx \quad \text{and we are given that}$$

$$\int f(x) \sin x \, dx = -f(x) \cos x + \int 3x^2 \cos x \, dx \Rightarrow f'(x) = 3x^2 \Rightarrow f(x) = x^3$$

22. A $A = \pi r^2$, $A = 64\pi$ when $r = 8$. Take the derivative with respect to t .

$$\frac{dA}{dt} = 2\pi r \cdot \frac{dr}{dt}; \quad 96\pi = 2\pi(8) \cdot \frac{dr}{dt} \Rightarrow \frac{dr}{dt} = 6$$

23. C $\lim_{h \rightarrow 0} \frac{\int_1^{1+h} \sqrt{x^5 + 8} \, dx}{h} = \lim_{h \rightarrow 0} \frac{F(1+h) - F(1)}{h} = F'(1)$ where $F'(x) = \sqrt{x^5 + 8}$. $F'(1) = 3$

Alternate solution by L'Hôpital's Rule: $\lim_{h \rightarrow 0} \frac{\int_1^{1+h} \sqrt{x^5 + 8} \, dx}{h} = \lim_{h \rightarrow 0} \frac{\sqrt{(1+h)^5 + 8}}{1} = \sqrt{9} = 3$

24. D $\text{Area} = \frac{1}{2} \int_0^{\pi/2} \sin^2(2\theta) \, d\theta = \frac{1}{2} \int_0^{\pi/2} \frac{1}{2} (1 - \cos 4\theta) \, d\theta = \frac{1}{4} \left(\theta - \frac{1}{4} \sin 4\theta \right) \Big|_0^{\pi/2} = \frac{\pi}{8}$

25. C At rest when $v(t) = 0$. $v(t) = e^{-2t} - 2te^{-2t} = e^{-2t}(1 - 2t)$, $v(t) = 0$ at $t = \frac{1}{2}$ only.

26. E Apply the log function, simplify, and differentiate. $\ln y = \ln(\sin x)^x = x \ln(\sin x)$

$$\frac{y'}{y} = \ln(\sin x) + x \cdot \frac{\cos x}{\sin x} \Rightarrow y' = y(\ln(\sin x) + x \cdot \cot x) = (\sin x)^x (\ln(\sin x) + x \cdot \cot x)$$

27. E Each of the right-hand sides represent the area of a rectangle with base length $(b - a)$.

I. Area under the curve is less than the area of the rectangle with height $f(b)$.

II. Area under the curve is more than the area of the rectangle with height $f(a)$.

III. Area under the curve is the same as the area of the rectangle with height $f(c)$, $a < c < b$.

Note that this is the Mean Value Theorem for Integrals.

28. E $\int e^{x+e^x} \, dx = \int e^{e^x} (e^x \, dx)$. This is of the form $\int e^u \, du$, $u = e^x$, so $\int e^{x+e^x} \, dx = e^{e^x} + C$

29. D Let $x - \frac{\pi}{4} = t$. $\lim_{x \rightarrow \frac{\pi}{4}} \frac{\sin\left(x - \frac{\pi}{4}\right)}{x - \frac{\pi}{4}} = \lim_{t \rightarrow 0} \frac{\sin t}{t} = 1$

30. B At $t = 1$, $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{\frac{3}{2\sqrt{3t+1}}}{3t^2-1} \Big|_{t=1} = \frac{\frac{3}{4}}{3-1} = \frac{3}{8}$

31. D The center is $x = 1$, so only C, D, or E are possible. Check the endpoints.

At $x = 0$: $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ converges by alternating series test.

At $x = 2$: $\sum_{n=1}^{\infty} \frac{1}{n}$ which is the harmonic series and known to diverge.

32. E $y(-1) = -6$, $y'(-1) = 3x^2 + 6x + 7 \Big|_{x=-1} = 4$, the slope of the normal is $-\frac{1}{4}$ and an equation for the normal is $y + 6 = -\frac{1}{4}(x + 1) \Rightarrow x + 4y = -25$.

33. C This is the differential equation for exponential growth.

$$y = y(0)e^{-2t} = e^{-2t}; \frac{1}{2} = e^{-2t}; -2t = \ln\left(\frac{1}{2}\right) \Rightarrow t = -\frac{1}{2}\ln\left(\frac{1}{2}\right) = \frac{1}{2}\ln 2$$


34. A This topic is no longer part of the AP Course Description. $\sum 2\pi\rho \Delta s$ where $\rho = x = y^3$

$$\text{Surface Area} = \int_0^1 2\pi y^3 \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy = \int_0^1 2\pi y^3 \sqrt{1 + (3y^2)^2} dy = 2\pi \int_0^1 y^3 \sqrt{1 + 9y^4} dy$$

35. B Use shells (which is no longer part of the AP Course Description)

$$\sum 2\pi rh \Delta x \text{ where } r = x \text{ and } h = y = 6x - x^2$$

$$\text{Volume} = 2\pi \int_0^6 x(6x - x^2) dx$$



36. E $\int_{-1}^1 \frac{3}{x^2} dx = 2 \int_0^1 \frac{3}{x^2} dx = 2 \lim_{L \rightarrow 0^+} \int_L^1 \frac{3}{x^2} dx = 2 \lim_{L \rightarrow 0^+} -\frac{3}{x} \Big|_L^1$ which does not exist.
37. A This topic is no longer part of the AP Course Description. $y = y_h + y_p$ where $y_h = ce^{-x}$ is the solution to the homogeneous equation $\frac{dy}{dx} + y = 0$ and $y_p = (Ax^2 + Bx)e^{-x}$ is a particular solution to the given differential equation. Substitute y_p into the differential equation to determine the values of A and B . The answer is $A = \frac{1}{2}$, $B = 0$.
38. C $\lim_{x \rightarrow \infty} (1 + 5e^x)^{1/x} = \lim_{x \rightarrow \infty} e^{\ln(1+5e^x)^{1/x}} = \lim_{x \rightarrow \infty} \ln(1+5e^x)^{1/x} = \lim_{x \rightarrow \infty} \frac{\ln(1+5e^x)}{x} = \lim_{x \rightarrow \infty} \frac{5e^x}{1+5e^x} = e$
39. A Square cross sections: $\sum y^2 \Delta x$ where $y = e^{-x}$. $V = \int_0^3 e^{-2x} dx = -\frac{1}{2} e^{-2x} \Big|_0^3 = \frac{1}{2} (1 - e^{-6})$
40. A $u = \frac{x}{2}$, $du = \frac{1}{2} dx$; when $x = 2$, $u = 1$ and when $x = 4$, $u = 2$
- $$\int_2^4 \frac{1 - \left(\frac{x}{2}\right)^2}{x} dx = \int_1^2 \frac{1 - u^2}{2u} \cdot 2 du = \int_1^2 \frac{1 - u^2}{u} du$$
41. C $y' = x^{\frac{1}{2}}$, $L = \int_0^3 \sqrt{1 + (y')^2} dx = \int_0^3 \sqrt{1 + x} dx = \frac{2}{3} (1 + x)^{3/2} \Big|_0^3 = \frac{2}{3} (4^{3/2} - 1^{3/2}) = \frac{2}{3} (8 - 1) = \frac{14}{3}$
42. E Since $e^u = 1 + u + \frac{u^2}{2!} + \frac{u^3}{3!} + \dots$, then $e^{3x} = 1 + 3x + \frac{(3x)^2}{2!} + \frac{(3x)^3}{3!} + \dots$
- The coefficient we want is $\frac{3^3}{3!} = \frac{9}{2}$
43. E Graphs A and B contradict $f'' < 0$. Graph C contradicts $f'(0)$ does not exist. Graph D contradicts continuity on the interval $[-2, 3]$. Graph E meets all given conditions.
44. A $\frac{dy}{dx} = 3x^2 y \Rightarrow \frac{dy}{y} = 3x^2 dx \Rightarrow \ln|y| = x^3 + K$; $y = Ce^{x^3}$ and $y(0) = 8$ so, $y = 8e^{x^3}$

45. D The expression is a Riemann sum with $\Delta x = \frac{1}{n}$ and $f(x) = x^2$.

The evaluation points are: $\frac{1}{n}, \frac{2}{n}, \frac{3}{n}, \dots, \frac{3n}{n}$

Thus the right Riemann sum is for $x = 0$ to $x = 3$. The limit is equal to $\int_0^3 x^2 dx$.

1. C $\frac{dy}{dx} = x^2 \cdot \frac{d}{dx}(e^x) + e^x \cdot \frac{d}{dx}(x^2) = x^2 e^x + 2xe^x = xe^x(x+2)$
2. D $x^2 - 4 \geq 0$ and $x \neq 3 \Rightarrow |x| \geq 2$ and $x \neq 3$
3. A Distance $= \int_0^2 |v(t)| dt = \int_0^2 e^t dt = e^t \Big|_0^2 = e^2 - e^0 = e^2 - 1$
4. E Students should know what the graph looks like without a calculator and choose option E.
Or $y = -5(x-2)^{-1}$; $y' = 5(x-2)^{-2}$; $y'' = -10(x-2)^{-3}$. $y'' < 0$ for $x > 2$.
5. A $\int \sec^2 x dx = \int d(\tan x) = \tan x + C$
6. D $\frac{dy}{dx} = \frac{x \cdot \frac{d}{dx}(\ln x) - \ln x \cdot \frac{d}{dx}(x)}{x^2} = \frac{x \cdot \left(\frac{1}{x}\right) - \ln x \cdot (1)}{x^2} = \frac{1 - \ln x}{x^2}$
7. D $\int x(3x^2 + 5)^{-\frac{1}{2}} dx = \frac{1}{6} \int (3x^2 + 5)^{-\frac{1}{2}} (6x dx) = \frac{1}{6} \cdot 2(3x^2 + 5)^{\frac{1}{2}} + C = \frac{1}{3}(3x^2 + 5)^{\frac{1}{2}} + C$
8. B $\frac{dy}{dx} > 0 \Rightarrow y$ is increasing; $\frac{d^2y}{dx^2} < 0 \Rightarrow$ graph is concave down. This is only on $b < x < c$.
9. E $1 + (2x \cdot y' + 2y) - 2y \cdot y' = 0$; $y' = \frac{1+2y}{2y-2x}$. This cannot be evaluated at (1,1) and so y' does not exist at (1,1).
10. C $18 = \left(kx^2 - \frac{1}{3}x^3\right) \Big|_0^k = \frac{2}{3}k^3 \Rightarrow k^3 = 27$, so $k = 3$
11. A $f'(x) = x \cdot 3(1-2x)^2(-2) + (1-2x)^3$; $f'(1) = -7$. Only option A has a slope of -7 .
12. B $f'\left(\frac{\pi}{3}\right) = \cos\left(\frac{\pi}{3}\right) = \frac{1}{2}$

13. A By the Fundamental Theorem of Calculus $\int_0^c f'(x) dx = f(x) \Big|_0^c = f(c) - f(0)$
14. D $\int_0^{\frac{\pi}{2}} (1 + \sin \theta)^{-1/2} (\cos \theta d\theta) = 2(1 + \sin \theta)^{1/2} \Big|_0^{\frac{\pi}{2}} = 2(\sqrt{2} - 1)$
15. B $f(x) = \sqrt{2x} = \sqrt{2} \cdot \sqrt{x}$; $f'(x) = \sqrt{2} \cdot \frac{1}{2\sqrt{x}}$; $f'(2) = \sqrt{2} \cdot \frac{1}{2\sqrt{2}} = \frac{1}{2}$
16. C At rest when $0 = v(t) = x'(t) = 3t^2 - 6t - 9 = 3(t^2 - 2t - 3) = 3(t-3)(t+1)$
 $t = -1, 3$ and $t \geq 0 \Rightarrow t = 3$
17. D $\int_0^1 (3x-2)^2 dx = \frac{1}{3} \int_0^1 (3x-2)^2 (3 dx) = \frac{1}{3} \cdot \frac{1}{3} (3x-2)^3 \Big|_0^1 = \frac{1}{9} (1 - (-8)) = 1$
18. E $y' = 2 \cdot \left(-\sin\left(\frac{x}{2}\right) \cdot \frac{1}{2} \right) = -\sin\left(\frac{x}{2}\right)$; $y'' = -\left(\cos\left(\frac{x}{2}\right) \cdot \left(\frac{1}{2}\right) \right) = -\frac{1}{2} \cos\left(\frac{x}{2}\right)$
19. B $\int_2^3 \frac{x}{x^2+1} dx = \frac{1}{2} \int_2^3 \frac{2x dx}{x^2+1} = \frac{1}{2} \ln(x^2+1) \Big|_2^3 = \frac{1}{2} (\ln 10 - \ln 5) = \frac{1}{2} \ln 2$
20. C Consider the cases:
 I. false if $f(x) = 1$
 II. This is true by the Mean Value Theorem
 III. false if the graph of f is a parabola with vertex at $x = \frac{a+b}{2}$.
 Only II must be true.
21. C $x = x^2 - 3x + 3$ at $x = 1$ and at $x = 3$.
 Area = $\int_1^3 (x - (x^2 - 3x + 3)) dx = \int_1^3 (-x^2 + 4x - 3) dx = \left(-\frac{1}{3}x^3 + 2x^2 - 3x \right) \Big|_1^3 = \frac{4}{3}$
22. C $2 = \ln x - \ln \frac{1}{x} = \ln x + \ln x \Rightarrow \ln x = 1 \Rightarrow x = e$

23. B By L'Hôpital's rule (which is no longer part of the AB Course Description),

$$\lim_{x \rightarrow 0} \frac{f(x)}{g(x)} = \lim_{x \rightarrow 0} \frac{f'(x)}{g'(x)} = \frac{f'(0)}{g'(0)} = \frac{\cos 0}{1} = \frac{1}{1} = 1$$

Alternatively, $f'(x) = \cos x$ and $f(0) = 0 \Rightarrow f(x) = \sin x$. Also $g'(x) = 1$ and

$$g(0) = 0 \Rightarrow g(x) = x. \text{ Hence } \lim_{x \rightarrow 0} \frac{f(x)}{g(x)} = \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1.$$

24. C Let $y = x^{\ln x}$ and take the ln of each side. $\ln y = \ln x^{\ln x} = \ln x \cdot \ln x$. Take the derivative of each side with respect to x . $\frac{y'}{y} = 2 \ln x \cdot \frac{1}{x} \Rightarrow y' = 2 \ln x \cdot \frac{1}{x} \cdot x^{\ln x}$


25. B Use the Fundamental Theorem of Calculus. $f'(x) = \frac{1}{x}$

26. E Use the technique of antiderivatives by parts: Let $u = x$ and $dv = \cos x \, dx$.

$$\int_0^{\frac{\pi}{2}} x \cos x \, dx = \left(x \sin x - \int \sin x \, dx \right) \Big|_0^{\frac{\pi}{2}} = (x \sin x + \cos x) \Big|_0^{\frac{\pi}{2}} = \frac{\pi}{2} - 1$$

27. E The function is continuous at $x = 3$ since $\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^+} f(x) = 9 = f(3)$. Also, the derivative as you approach $x = 3$ from the left is 6 and the derivative as you approach $x = 3$ from the right is also 6. These two facts imply that f is differentiable at $x = 3$. The function is clearly continuous and differentiable at all other values of x .

28. C The graph is a V with vertex at $x = 3$. The integral gives the sum of the areas of the two triangles that the V forms with the horizontal axis for x from 1 to 4. These triangles have areas of 2 and 0.5 respectively.




29. B This limit gives the derivative of the function $f(x) = \tan(3x)$. $f'(x) = 3 \sec^2(3x)$

30. A Shells (which is no longer part of the AB Course Description)

$$\sum 2\pi rh\Delta x, \text{ where } r = x, h = e^{2x}$$

$$\text{Volume} = 2\pi \int_0^1 xe^{2x} dx$$



31. C Let $y = f(x)$ and solve for x .

$$y = \frac{x}{x+1}; xy + y = x; x(y-1) = -y; x = \frac{y}{1-y} \Rightarrow f^{-1}(x) = \frac{x}{1-x}$$

32. A The period for $\sin\left(\frac{x}{2}\right)$ is $\frac{2\pi}{\frac{1}{2}} = 4\pi$.

33. A Check the critical points and the endpoints.

$$f'(x) = 3x^2 - 6x = 3x(x-2) \text{ so the critical points are } 0 \text{ and } 2.$$

x	-2	0	2	4
$f(x)$	-8	12	8	28

Absolute maximum is at $x = 4$.

34. D The interval is $x = a$ to $x = c$. The height of a rectangular slice is the top curve, $f(x)$, minus the bottom curve, $g(x)$. The area of the rectangular slice is therefore $(f(x) - g(x))\Delta x$. Set up a Riemann sum and take the limit as Δx goes to 0 to get a definite integral.

$$\begin{aligned} 35. \text{ B } \quad 4 \cos\left(x + \frac{\pi}{3}\right) &= 4 \left(\cos x \cdot \cos\left(\frac{\pi}{3}\right) - \sin x \cdot \sin\left(\frac{\pi}{3}\right) \right) \\ &= 4 \left(\cos x \cdot \frac{1}{2} - \sin x \cdot \frac{\sqrt{3}}{2} \right) = 2 \cos x - 2\sqrt{3} \sin x \end{aligned}$$

36. C $3x - x^2 = x(3-x) > 0$ for $0 < x < 3$

$$\text{Average value} = \frac{1}{3} \int_0^3 (3x - x^2) dx = \frac{1}{3} \left(\frac{3}{2}x^2 - \frac{1}{3}x^3 \right) \Big|_0^3 = \frac{3}{2}$$

37. D Since $e^x > 0$ for all x , the zeros of $f(x)$ are the zeros of $\sin x$, so $x = 0, \pi, 2\pi$.

38. E $\int \left(\frac{1}{x} \int_1^x \frac{du}{u} \right) dx = \int \frac{1}{x} \ln x \, dx = \int \ln x \left(\frac{dx}{x} \right)$. This is $\int u \, du$ with $u = \ln x$, so the value is $\frac{(\ln x)^2}{2} + C$

39. E $\int_3^{10} f(x) \, dx = -\int_{10}^3 f(x) \, dx$; $\int_1^3 f(x) \, dx = \int_1^{10} f(x) \, dx - \int_3^{10} f(x) \, dx = 4 - (-7) = 11$


40. B $x^2 + y^2 = z^2$, take the derivative of both sides with respect to t . $2x \cdot \frac{dx}{dt} + 2y \cdot \frac{dy}{dt} = 2z \cdot \frac{dz}{dt}$
Divide by 2 and substitute: $4 \cdot \frac{dx}{dt} + 3 \cdot \frac{1}{3} \frac{dx}{dt} = 5 \cdot 1 \Rightarrow \frac{dx}{dt} = 1$

41. A The statement makes no claim as to the behavior of f at $x = 3$, only the value of f for input arbitrarily close to $x = 3$. None of the statements are true.

42. C $\lim_{x \rightarrow \infty} \frac{x}{x+1} = \lim_{x \rightarrow \infty} \frac{\frac{x}{x}}{\frac{x}{x} + \frac{1}{x}} = \lim_{x \rightarrow \infty} \frac{1}{1 + \frac{1}{x}} = 1$.

None of the other functions have a limit of 1 as $x \rightarrow \infty$

43. B The cross-sections are disks with radius $r = y$ where $y = \frac{1}{3} \sqrt{9 - x^2}$.



$$\text{Volume} = \pi \int_{-3}^3 y^2 \, dx = 2\pi \int_0^3 \frac{1}{9} (9 - x^2) \, dx = \frac{2\pi}{9} \left(9x - \frac{1}{3} x^3 \right) \Big|_0^3 = 4\pi$$

44. C For I: $p(-x) = f(g(-x)) = f(-g(x)) = -f(g(x)) = -p(x) \Rightarrow p$ is odd.

For II: $r(-x) = f(-x) + g(-x) = -f(x) - g(x) = -(f(x) + g(x)) = -r(x) \Rightarrow r$ is odd.

For III: $s(-x) = f(-x) \cdot g(-x) = (-f(x)) \cdot (-g(x)) = f(x) \cdot g(x) = s(x) \Rightarrow s$ is not odd.

45. D Volume = $\pi r^2 h = 16\pi \Rightarrow h = 16r^{-2}$. $A = 2\pi r h + 2\pi r^2 = 2\pi(16r^{-1} + r^2)$

$$\frac{dA}{dr} = 2\pi(-16r^{-2} + 2r) = 4\pi r^{-2}(-8 + r^3); \quad \frac{dA}{dr} < 0 \text{ for } 0 < r < 2 \text{ and } \frac{dA}{dr} > 0 \text{ for } r > 2$$

The minimum surface area of the can is when $r = 2 \Rightarrow h = 4$.

1. A $\int_0^1 (x - x^2) dx = \frac{1}{2}x^2 - \frac{1}{3}x^3 \Big|_0^1 = \frac{1}{2} - \frac{1}{3} = \frac{1}{6}$ only.
2. D $\int_0^1 x(x^2 + 2)^2 dx = \frac{1}{2} \int_0^1 (x^2 + 2)^2 (2x dx) = \frac{1}{2} \cdot \frac{1}{3} (x^2 + 2)^3 \Big|_0^1 = \frac{1}{6} (3^3 - 2^3) = \frac{19}{6}$
3. B $f(x) = \ln \sqrt{x} = \frac{1}{2} \ln x$; $f'(x) = \frac{1}{2} \cdot \frac{1}{x} \Rightarrow f''(x) = -\frac{1}{2x^2}$
4. E $\left(\frac{uv}{w}\right)' = \frac{(uv' + u'v)w - uvw'}{w^2} = \frac{uv'w + u'vw - uvw'}{w^2}$
5. C $\lim_{x \rightarrow a} f(x) = f(a)$ for all values of a except 2.
 $\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} (x - 2) = 0 \neq -1 = f(2)$
6. C $2y \cdot y' - 2x \cdot y' - 2y = 0 \Rightarrow y' = \frac{y}{y - x}$
7. A $\int_2^\infty \frac{dx}{x^2} = \lim_{L \rightarrow \infty} \int_2^L \frac{dx}{x^2} = \lim_{L \rightarrow \infty} \left(-\frac{1}{x}\right) \Big|_2^L = \lim_{L \rightarrow \infty} \left(\frac{1}{2} - \frac{1}{L}\right) = \frac{1}{2}$
8. A $f'(x) = e^x$, $f'(2) = e^2$, $\ln e^2 = 2$
9. D II does not work since the slope of f at $x = 0$ is not equal to $f'(0)$. Both I and III could work. For example, $f(x) = e^x$ in I and $f(x) = \sin x$ in III.
10. D This limit is the derivative of $\sin x$.
11. A The slope of the line is $-\frac{1}{7}$, so the slope of the tangent line at $x = 1$ is $7 \Rightarrow f'(1) = 7$.
12. B $v(t) = 3t + C$ and $v(2) = 10 \Rightarrow C = 4$ and $v(t) = 3t + 4$.
 Distance $= \int_0^2 (3t + 4) dt = \frac{3}{2}x^2 + 4t \Big|_0^2 = 14$

13. B The Maclaurin series for $\sin t$ is $t - \frac{t^3}{3!} + \frac{t^5}{5!} - \dots$. Let $t = 2x$.

$$\sin(2x) = 2x - \frac{(2x)^3}{3!} + \frac{(2x)^5}{5!} - \dots + \frac{(-1)^{n-1}(2x)^{2n-1}}{(2n-1)!} + \dots$$

14. A Use the Fundamental Theorem of Calculus: $\sqrt{1+(x^2)^3} \cdot \frac{d(x^2)}{dx} = 2x\sqrt{1+x^6}$

15. E $x = t^2 + 1$, $\frac{dx}{dt} = 2t$, $\frac{d^2x}{dt^2} = 2$; $y = \ln(2t+3)$, $\frac{dy}{dt} = \frac{2}{2t+3}$; $\frac{d^2y}{dt^2} = -\frac{4}{(2t+3)^2}$

16. A Use the technique of antiderivatives by parts

$$u = x \quad dv = e^{2x} dx$$

$$du = dx \quad v = \frac{1}{2}e^{2x}$$

$$\frac{1}{2}xe^{2x} - \frac{1}{2}\int e^{2x} dx = \frac{1}{2}xe^{2x} - \frac{1}{4}e^{2x} + C$$

17. D Use partial fractions:

$$\int_2^3 \frac{3}{(x-1)(x+1)} dx = \int_2^3 \left(\frac{1}{x-1} - \frac{1}{x+2} \right) dx = (\ln|x-1| - \ln|x+2|) \Big|_2^3 = \ln 2 - \ln 5 - \ln 1 + \ln 4 = \ln \frac{8}{5}$$

18. E $\Delta x = \frac{4 - (-2)}{3} = 2$, $T = \frac{1}{2}(2) \left(\frac{e^4}{2} + 2 \cdot \frac{e^2}{2} + 2 \cdot \frac{e^0}{2} + \frac{e^{-2}}{2} \right) = \frac{1}{2}(e^4 + 2e^2 + 2e^0 + e^{-2})$

19. B Make a sketch. $x < -2$ one zero, $-2 < x < 5$ no zeros, $x > 5$ one zero for a total of 2 zeros

20. E This is the definition of a limit.

21. D $\frac{1}{2} \int_1^3 \frac{1}{x} dx = \frac{1}{2} \ln x \Big|_1^3 = \frac{1}{2}(\ln 3 - \ln 1) = \frac{1}{2} \ln 3$

22. E Quick Solution: f' must have a factor of f which makes E the only option. Or,

$$\ln f(x) = x \ln(x^2 + 1) \Rightarrow \frac{f'(x)}{f(x)} = x \cdot \frac{2x}{x^2 + 1} + \ln(x^2 + 1) \Rightarrow f'(x) = f(x) \cdot \left(\frac{2x^2}{x^2 + 1} + \ln(x^2 + 1) \right)$$

23. E $r = 0$ when $\cos 3\theta = 0 \Rightarrow \theta = \pm \frac{\pi}{6}$. The region is for the interval from $\theta = -\frac{\pi}{6}$ to $\theta = \frac{\pi}{6}$.

$$\text{Area} = \frac{1}{2} \int_{-\frac{\pi}{6}}^{\frac{\pi}{6}} (4 \cos 3\theta)^2 d\theta$$

24. D $f'(x) = 3x^2 - 4x$, $f(0) = 0$ and $f(2) = 0$. By the Mean Value Theorem,

$$0 = \frac{f(2) - f(0)}{2 - 0} = f'(c) = 3c^2 - 4c \text{ for } c \in (0, 2). \text{ So, } c = \frac{4}{3}.$$


25. D Square cross-sections: $\sum y^2 \Delta x$ where $y = 4x^2$. Volume = $\int_0^1 16x^4 dx = \frac{16}{5} x^5 \Big|_0^1 = \frac{16}{5}$.

26. C This is not true if f is not an even function.

27. B $y'(x) = 3x^2 + 2ax + b$, $y''(x) = 6x + 2a$, $y''(1) = 0 \Rightarrow a = -3$
 $y(1) = -6$ so, $-6 = 1 + a + b - 4 \Rightarrow -6 = 1 - 3 + b - 4 \Rightarrow b = 0$

28. E
$$\frac{\frac{d}{dx} \left(\cos \left(\frac{\pi}{x} \right) \right)}{\cos \left(\frac{\pi}{x} \right)} = \frac{-\sin \left(\frac{\pi}{x} \right) \cdot \frac{d}{dx} \left(\frac{\pi}{x} \right)}{\cos \left(\frac{\pi}{x} \right)} = \frac{-\sin \left(\frac{\pi}{x} \right) \cdot \left(-\frac{\pi}{x^2} \right)}{\cos \left(\frac{\pi}{x} \right)} = \frac{\pi}{x^2} \tan \left(\frac{\pi}{x} \right)$$

29. B Disks: $\sum \pi x^2 \Delta y$ where $x^2 = y - 1$.
 Volume = $\pi \int_1^5 (y - 1) dy = \frac{\pi}{2} (y - 1)^2 \Big|_1^5 = 8\pi$



30. C This is an infinite geometric series with ratio $\frac{1}{3}$ and first term $\frac{1}{3^n}$.

$$\text{Sum} = \frac{\text{first}}{1 - \text{ratio}} = \frac{\left(\frac{1}{3^n}\right)}{1 - \frac{1}{3}} = \frac{3}{2} \cdot \left(\frac{1}{3^n}\right)$$

31. C This integral gives $\frac{1}{4}$ of the area of the circle with center at the origin and radius = 2.

$$\frac{1}{4}(\pi \cdot 2^2) = \pi$$

32. E No longer covered in the AP Course Description. The solution is of the form $y = y_h + y_p$ where y_h is the solution to $y' - y = 0$ and the form of y_p is $Ax^2 + Bx + K$. Hence $y_h = Ce^x$. Substitute y_p into the original differential equation to determine the values of A , B , and K .

Another technique is to substitute each of the options into the differential equation and pick the one that works. Only (A), (B), and (E) are viable options because of the form for y_h . Both (A) and (B) fail, so the solution is (E).

33. E
$$L = \int_0^2 \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx = \int_0^2 \sqrt{1 + (3x^2)^2} dx = \int_0^2 \sqrt{1 + 9x^4} dx$$

34. C At $t = 1$, $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{4t^3 + 4t}{3t^2 + 1} \Big|_{t=1} = \frac{8}{4} = 2$; the point at $t = 1$ is $(2, 3)$. $y = 3 + 2(x - 2) = 2x - 1$

35. A Quick solution: For large x the exponential function dominates any polynomial, so

$$\lim_{x \rightarrow +\infty} \frac{x^k}{e^x} = 0.$$

Or, repeated use of L'Hôpital's rule gives $\Rightarrow \lim_{x \rightarrow +\infty} \frac{x^k}{e^x} = \lim_{x \rightarrow +\infty} \frac{k!}{e^x} = 0$


36. E Disks: $\sum \pi(R^2 - r^2)\Delta x$ where $R = 1$, $r = \sin x$

$$\text{Volume} = \pi \int_0^{\pi/2} (1 - \sin^2 x) dx$$

Note that the expression in (E) can also be written as


$$\begin{aligned} \pi \int_0^{\pi/2} \cos^2 x dx &= -\pi \int_{\pi/2}^0 \cos^2 \left(\frac{\pi}{2} - x\right) dx \\ &= \pi \int_0^{\pi/2} \sin^2 x dx \end{aligned}$$

and therefore option (D) is also a correct answer.



37. D $\frac{x+S}{8} = \frac{S}{2} \Rightarrow x = 3S$

$$\frac{dx}{dt} = 3 \frac{dS}{dt} = 3 \cdot \frac{4}{9} = \frac{4}{3}$$



38. C Check $x = -1$, $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ which is convergent by alternating series test

Check $x = 1$, $\sum_{n=1}^{\infty} \frac{1}{n}$ which is the harmonic series and known to diverge.

39. C $\frac{dy}{y} = \sec^2 x dx \Rightarrow \ln|y| = \tan x + k \Rightarrow y = Ce^{\tan x}$. $y(0) = 5 \Rightarrow y = 5e^{\tan x}$

40. E Since f and g are inverses their derivatives at the inverse points are reciprocals. Thus,

$$g'(-2) \cdot f'(5) = 1 \Rightarrow g'(-2) = \frac{1}{f'(5)} = -\frac{1}{2}$$

41. B Take the interval $[0,1]$ and divide it into n pieces of equal length and form the right Riemann Sum for the function $f(x) = \sqrt{x}$. The limit of this sum is what is given and its value is given by $\int_0^1 \sqrt{x} dx$

42. A Let $5-x=u$, $dx=-du$, substitute

$$\int_1^4 f(5-x) dx = \int_4^1 f(u)(-du) = \int_1^4 f(u) du = \int_1^4 f(x) dx = 6$$

43. A This is an example of exponential growth, $B = B_0 \cdot 2^{t/3}$. Find the value of t so $B = 3B_0$.

$$3B_0 = B_0 \cdot 2^{t/3} \Rightarrow 3 = 2^{t/3} \Rightarrow \ln 3 = \frac{t}{3} \ln 2 \Rightarrow t = \frac{3 \ln 3}{\ln 2}$$

44. A I. Converges by Alternate Series Test

II Diverges by the nth term test: $\lim_{n \rightarrow \infty} \frac{1}{n} \left(\frac{3}{2}\right)^n \neq 0$

III Diverges by Integral test: $\int_2^{\infty} \frac{1}{x \ln x} dx = \lim_{L \rightarrow \infty} \ln(\ln x) \Big|_2^L = \infty$


45. B $A = (2x)(2y) = 4xy$ and $y = \sqrt{4 - \frac{4}{9}x^2}$.

$$\text{So } A = 8x \sqrt{1 - \frac{1}{9}x^2}.$$

$$\begin{aligned} A' &= 8 \left(\left(1 - \frac{1}{9}x^2\right)^{1/2} + \frac{1}{2}x \left(1 - \frac{1}{9}x^2\right)^{-1/2} \left(-\frac{2}{9}x\right) \right) \\ &= \frac{8}{9} \left(1 - \frac{1}{9}x^2\right)^{-1/2} (9 - 2x^2) \end{aligned}$$

$A' = 0$ at $x = \pm 3, \frac{3}{\sqrt{2}}$. The maximum area occurs when $x = \frac{3}{\sqrt{2}}$ and $y = \sqrt{2}$. The value of


the largest area is $A = 4xy = 4 \cdot \frac{3}{\sqrt{2}} \cdot \sqrt{2} = 12$



1. C $f'(x) = \frac{3}{2}x^{\frac{1}{2}}; f'(4) = \frac{3}{2} \cdot 4^{\frac{1}{2}} = \frac{3}{2} \cdot 2 = 3$
2. B Summing pieces of the form: (vertical) · (small width), vertical = $(d - f(x))$, width = Δx
 Area = $\int_a^b (d - f(x)) dx$
3. D Divide each term by n^3 . $\lim_{n \rightarrow \infty} \frac{3n^3 - 5n}{n^3 - 2n^2 + 1} = \lim_{n \rightarrow \infty} \frac{3 - \frac{5}{n^2}}{1 - \frac{2}{n} + \frac{1}{n^3}} = 3$
4. A $3x^2 + 3(y + x \cdot y') + 6y^2 \cdot y' = 0; y'(3x + 6y^2) = -(3x^2 + 3y)$
 $y' = -\frac{3x^2 + 3y}{3x + 6y^2} = -\frac{x^2 + y}{x + 2y^2}$
5. A $\lim_{x \rightarrow -2} \frac{x^2 - 4}{x + 2} = \lim_{x \rightarrow -2} \frac{(x+2)(x-2)}{x+2} = \lim_{x \rightarrow -2} (x-2) = -4$. For continuity $f(-2)$ must be -4 .
6. D Area = $\int_3^4 \frac{1}{x-1} dx = (\ln|x-1|) \Big|_3^4 = \ln 3 - \ln 2 = \ln \frac{3}{2}$
7. B $y' = \frac{2 \cdot (3x-2) - (2x+3) \cdot 3}{(3x-2)^2}; y'(1) = -13$. Tangent line: $y - 5 = -13(x - 1) \Rightarrow 13x + y = 18$
8. E $y' = \sec^2 x + \csc^2 x$
9. E $h(x) = f(|x|) = 3|x|^2 - 1 = 3x^2 - 1$
10. D $f'(x) = 2(x-1) \cdot \sin x + (x-1)^2 \cos x; f'(0) = (-2) \cdot 0 + 1 \cdot 1 = 1$
11. C $a(t) = 6t - 2; v(t) = 3t^2 - 2t + C$ and $v(3) = 25 \Rightarrow 25 = 27 - 6 + C; v(t) = 3t^2 - 2t + 4$
 $x(t) = t^3 - t^2 + 4t + K$; Since $x(1) = 10, K = 6; x(t) = t^3 - t^2 + 4t + 6$.

12. B The only one that is true is II. The others can easily be seen as false by examples. For example, let $f(x) = 1$ and $g(x) = 1$ with $a = 0$ and $b = 2$. Then I gives $2 = 4$ and III gives $2 = \sqrt{2}$, both false statements.
13. A $\text{period} = \frac{2\pi}{B} = \frac{2\pi}{3}$
14. A Let $u = x^3 + 1$. Then $\int \frac{3x^2}{\sqrt{x^3 + 1}} dx = \int u^{-1/2} du = 2u^{1/2} + C = 2\sqrt{x^3 + 1} + C$
15. D $f'(x) = (x-3)^2 + 2(x-2)(x-3) = (x-3)(3x-7)$; $f'(x)$ changes from positive to negative at $x = \frac{7}{3}$.
16. B $y' = 2 \frac{\sec x \tan x}{\sec x} = 2 \tan x$; $y'(\pi/4) = 2 \tan(\pi/4) = 2$. The slope of the normal line $-\frac{1}{y'(\pi/4)} = -\frac{1}{2}$
17. E Expand the integrand. $\int (x^2 + 1)^2 dx = \int (x^4 + 2x^2 + 1) dx = \frac{1}{5}x^5 + \frac{2}{3}x^3 + x + C$
18. D Want c so that $f'(c) = \frac{f\left(\frac{3\pi}{2}\right) - f\left(\frac{\pi}{2}\right)}{\frac{3\pi}{2} - \frac{\pi}{2}} = \frac{\sin\left(\frac{3\pi}{4}\right) - \sin\left(\frac{\pi}{4}\right)}{\pi} = \frac{0}{\pi}$.
 $f'(c) = \frac{1}{2} \cos\left(\frac{c}{2}\right) = 0 \Rightarrow c = \pi$
19. E The only one that is true is E. A consideration of the graph of $y = f(x)$, which is a standard cubic to the left of 0 and a line with slope 1 to the right of 0, shows the other options to be false.

20. B Use Cylindrical Shells which is no part of the AP Course Description. The volume of each shell is of the form $(2\pi rh)\Delta x$ with $r = x$ and $h = y$. Volume = $2\pi \int_0^7 x(x+1)^{\frac{1}{3}} dx$.



21. C $y = x^{-2} - x^{-3}$; $y' = -2x^{-3} + 3x^{-4}$; $y'' = 6x^{-4} - 12x^{-5} = 6x^{-5}(x-2)$. The only domain value at which there is a sign change in y'' is $x = 2$. Inflection point at $x = 2$.
22. E $\int \frac{1}{x^2 - 2x + 2} dx = \int \frac{1}{(x^2 - 2x + 1) + 1} dx = \int \frac{1}{(x-1)^2 + 1} dx = \tan^{-1}(x-1) + C$
23. C A quick way to do this problem is to use the effect of the multiplicity of the zeros of f on the graph of $y = f(x)$. There is point of inflection and a horizontal tangent at $x = -2$. There is a horizontal tangent and turning point at $x = 3$. There is a horizontal tangent on the interval $(-2, 3)$. Thus, there must be 3 critical points. Also, $f'(x) = (x-3)^3(x+2)^4(9x-7)$.
24. A $f'(x) = \frac{2}{3}(x^2 - 2x - 1)^{-\frac{1}{3}}(2x - 2)$, $f'(0) = \frac{2}{3} \cdot (-1) \cdot (-2) = \frac{4}{3}$
25. C $\frac{d}{dx}(2^x) = 2^x \cdot \ln 2$
26. D $v(t) = 4 \sin t - t$; $a(t) = 4 \cos t - 1 = 0$ at $t = \cos^{-1}(1/4) = 1.31812$; $v(1.31812) = 2.55487$
27. C $f'(x) = 3x^2 + 12 > 0$. Thus f is increasing for all x .
28. B $\int_1^{500} (13^x - 11^x) dx + \int_2^{500} (11^x - 13^x) dx = \int_1^{500} (13^x - 11^x) dx - \int_2^{500} (13^x - 11^x) dx$
 $= \int_1^2 (13^x - 11^x) dx = \left(\frac{13^x}{\ln 13} - \frac{11^x}{\ln 11} \right) \Big|_1^2 = \frac{13^2 - 13}{\ln 13} - \frac{11^2 - 11}{\ln 11} = 14.946$

29. C Use L'Hôpital's Rule (which is no longer part of the AB Course Description).

$$\lim_{\theta \rightarrow 0} \frac{1 - \cos \theta}{2 \sin^2 \theta} = \lim_{\theta \rightarrow 0} \frac{\sin \theta}{4 \sin \theta \cos \theta} = \lim_{\theta \rightarrow 0} \frac{1}{4 \cos \theta} = \frac{1}{4}$$

A way to do this without L'Hôpital's rule is the following

$$\lim_{\theta \rightarrow 0} \frac{1 - \cos \theta}{2 \sin^2 \theta} = \lim_{\theta \rightarrow 0} \frac{1 - \cos \theta}{2(1 - \cos^2 \theta)} = \lim_{\theta \rightarrow 0} \frac{1 - \cos \theta}{2(1 - \cos \theta)(1 + \cos \theta)} = \lim_{\theta \rightarrow 0} \frac{1}{2(1 + \cos \theta)} = \frac{1}{4}$$

30. C Each slice is a disk whose volume is given by $\pi r^2 \Delta x$, where $r = \sqrt{x}$.

$$\text{Volume} = \pi \int_0^3 (\sqrt{x})^2 dx = \pi \int_0^3 x dx = \frac{\pi}{2} x^2 \Big|_0^3 = \frac{9}{2} \pi.$$

31. E $f(x) = e^{3 \ln(x^2)} = e^{\ln(x^6)} = x^6$; $f'(x) = 6x^5$

32. A $\int \frac{du}{\sqrt{a^2 - u^2}} = \sin^{-1} \left(\frac{u}{a} \right) + C, a > 0$

$$\int_0^{\sqrt{3}} \frac{dx}{\sqrt{4 - x^2}} = \sin^{-1} \left(\frac{x}{2} \right) \Big|_0^{\sqrt{3}} = \sin^{-1} \left(\frac{\sqrt{3}}{2} \right) - \sin^{-1}(0) = \frac{\pi}{3}$$

33. B Separate the variables. $y^{-2} dy = 2 dx$; $-\frac{1}{y} = 2x + C$; $y = \frac{-1}{2x + C}$. Substitute the point $(1, -1)$

to find the value of C . Then $-1 = \frac{-1}{2 + C} \Rightarrow C = -1$, so $y = \frac{1}{1 - 2x}$. When $x = 2$, $y = -\frac{1}{3}$.

34. D Let x and y represent the horizontal and vertical sides of the triangle formed by the ladder, the wall, and the ground.

$$x^2 + y^2 = 25; 2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0; 2(24) \frac{dx}{dt} + 2(7)(-3) = 0; \frac{dx}{dt} = \frac{7}{8}.$$

35. E For there to be a vertical asymptote at $x = -3$, the value of c must be 3. For $y = 2$ to be a horizontal asymptote, the value of a must be 2. Thus $a + c = 5$.

36. D Rectangle approximation = $e^0 + e^1 = 1 + e$

$$\text{Trapezoid approximation.} = \left(1 + 2e + e^4 \right) / 2.$$

$$\text{Difference} = (e^4 - 1) / 2 = 26.799.$$

37. C I and II both give the derivative at a . In III the denominator is fixed. This is not the derivative of f at $x = a$. This gives the slope of the secant line from $(a, f(a))$ to $(a+h, f(a+h))$.
38. A $f'(x) = x^2 - \sin x + C$, $f(x) = \frac{1}{3}x^3 + \cos x + Cx + K$. Option A is the only one with this form.
39. D $A = \pi r^2$ and $C = 2\pi r$; $\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$ and $\frac{dC}{dt} = 2\pi \frac{dr}{dt}$. For $\frac{dA}{dt} = \frac{dC}{dt}$, $r = 1$.
40. C The graph of $y = f(|x|)$ is symmetric to the y -axis. This leaves only options C and E. For $x > 0$, x and $|x|$ are the same, so the graphs of $f(x)$ and $f(|x|)$ must be the same. This is option C.
41. D Answer follows from the Fundamental Theorem of Calculus.
42. B This is an example of exponential growth. We know from pre-calculus that $w = 2\left(\frac{3.5}{2}\right)^{\frac{t}{2}}$ is an exponential function that meets the two given conditions. When $t = 3$, $w = 4.630$. Using calculus the student may translate the statement “increasing at a rate proportional to its weight” to mean exponential growth and write the equation $w = 2e^{kt}$. Using the given conditions, $3.5 = 2e^{2k}$; $\ln(1.75) = 2k$; $k = \frac{\ln(1.75)}{2}$; $w = 2e^{t \cdot \frac{\ln(1.75)}{2}}$. When $t = 3$, $w = 4.630$.
43. B Use the technique of antiderivative by parts, which is no longer in the AB Course Description. The formula is $\int u dv = uv - \int v du$. Let $u = f(x)$ and $dv = x dx$. This leads to
$$\int x f(x) dx = \frac{1}{2}x^2 f(x) - \frac{1}{2} \int x^2 f'(x) dx.$$
44. C $f'(x) = \ln x + x \cdot \frac{1}{x}$; $f'(x)$ changes sign from negative to positive only at $x = e^{-1}$.
 $f(e^{-1}) = -e^{-1} = -\frac{1}{e}$.

45. B Let $f(x) = x^3 + x - 1$. Then Newton's method (which is no longer part of the AP Course Description) gives

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} = x_n - \frac{x_n^3 + x_n - 1}{3x_n^2 + 1}$$

$$x_2 = 1 - \frac{1+1-1}{3+1} = \frac{3}{4}$$

$$x_3 = \frac{3}{4} - \frac{\left(\frac{3}{4}\right)^3 + \frac{3}{4} - 1}{3\left(\frac{3}{4}\right)^2 + 1} = \frac{59}{86} = 0.686$$

1. A $\int_0^1 (x - x^2) dx = \left(\frac{1}{2}x^2 - \frac{1}{3}x^3 \right) \Big|_0^1 = \frac{1}{2} - \frac{1}{3} = \frac{1}{6}$
2. C $\lim_{x \rightarrow 0} \frac{2x^2 + 1 - 1}{x^2} = 2$
3. E $Q'(x) = p(x) \Rightarrow$ degree of Q is $n + 1$
4. B If $x = 2$ then $y = 5$. $x \frac{dy}{dt} + y \frac{dx}{dt} = 0$; $2(3) + 5 \cdot \frac{dx}{dt} = 0 \Rightarrow \frac{dx}{dt} = -\frac{6}{5}$
5. D $r = 2 \sec \theta$; $r \cos \theta = 2 \Rightarrow x = 2$. This is a vertical line through the point $(2, 0)$.
6. A $\frac{dx}{dt} = 2t$, $\frac{dy}{dt} = 3t^2$ thus $\frac{dy}{dx} = \frac{3t^2}{2t} = \frac{3}{2}t$; $\frac{d^2y}{dx^2} = \frac{d\left(\frac{dy}{dx}\right)}{dx} = \frac{\frac{d\left(\frac{dy}{dx}\right)}{dt}}{\frac{dx}{dt}} = \frac{\frac{3}{2}}{2t} = \frac{3}{4t}$
7. A $\int_0^1 x^3 e^{x^4} dx = \frac{1}{4} \int_0^1 e^{x^4} (4x^3 dx) = \frac{1}{4} e^{x^4} \Big|_0^1 = \frac{1}{4}(e - 1)$
8. B $f(x) = \ln e^{2x} = 2x$, $f'(x) = 2$
9. D $f'(x) = \frac{2}{3} \cdot \frac{1}{x^{1/3}}$. This does not exist at $x = 0$. D is false, all others are true.
10. E I. $\ln x$ is continuous for $x > 0$
 II. e^x is continuous for all x
 III. $\ln(e^x - 1)$ is continuous for $x > 0$.
11. E $\int_4^\infty \frac{-2x}{\sqrt[3]{9-x^2}} dx = \lim_{b \rightarrow \infty} \frac{3}{2} (9-x^2)^{2/3} \Big|_4^b$. This limit diverges. Another way to see this without doing the integration is to observe that the denominator behaves like $x^{2/3}$ which has a smaller degree than the degree of the numerator. This would imply that the integral will diverge.

12. E $v(t) = 2 \cos 2t + 3 \sin 3t$, $a(t) = -4 \sin 2t + 9 \cos 3t$, $a(\pi) = -9$.

13. C $\frac{dy}{y} = x^2 dx$, $\ln|y| = \frac{1}{3}x^3 + C_1$, $y = Ce^{\frac{1}{3}x^3}$. Only C is of this form.

14. B The only place that $f'(x)$ changes sign from positive to negative is at $x = -3$.

15. D $f'(x) = e^{\tan^2 x} \cdot \frac{d(\tan^2 x)}{dx} = 2 \tan x \cdot \sec^2 x \cdot e^{\tan^2 x}$

16. A I. Compare with p -series, $p = 2$

II. Geometric series with $r = \frac{6}{7}$

III. Alternating harmonic series

17. A Using implicit differentiation, $\frac{y+xy'}{xy} = 1$. When $x = 1$, $\frac{y+y'}{y} = 1 \Rightarrow y' = 0$.


Alternatively, $xy = e^x$, $y = \frac{e^x}{x}$, $y' = \frac{xe^x - e^x}{x^2} = \frac{e^x(x-1)}{x^2}$. $y'(1) = 0$

18. B $f'(x) \cdot e^{f(x)} = 2x \Rightarrow f'(x) = \frac{2x}{e^{f(x)}} = \frac{2x}{1+x^2}$

19. B Use cylindrical shells which is no longer part of the AP Course Description. Each shell is of the form $2\pi rh\Delta x$ where $r = x$ and $h = kx - x^2$. Solve the equation

$$10 = 2\pi \int_0^k x(kx - x^2) dx = 2\pi \left(\frac{kx^3}{3} - \frac{x^4}{4} \right) \Big|_0^k = 2\pi \cdot \frac{k^4}{12}.$$

$$k = \sqrt[4]{\frac{60}{\pi}} \approx 2.0905.$$



20. E $v(t) = -\frac{1}{2}e^{-2t} + 3$ and $x(t) = \frac{1}{4}e^{-2t} + 3t + 4$

21. A Use logarithms.

$$\ln y = \frac{1}{3} \ln(x^2 + 8) - \frac{1}{4} \ln(2x + 1); \frac{y'}{y} = \frac{2x}{3(x^2 + 8)} - \frac{2}{4(2x + 1)}; \text{ at } (0, 2), y' = -1.$$

22. B $f'(x) = x^2 e^x + 2x e^x = x e^x (x + 2)$; $f'(x) < 0$ for $-2 < x < 0$

23. D $L = \int_0^4 \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = \int_0^4 \sqrt{4t^2 + 1} dt$

24. C This is L'Hôpital's Rule.

25. D At $t = 3$, slope $= \frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{-te^{-t} + e^{-t}}{e^t} = \frac{1-t}{e^{2t}} \Big|_{t=3} = -\frac{2}{e^6} = -0.005$

26. B $\frac{2e^{2x}}{1+(e^{2x})^2} = \frac{2e^{2x}}{1+e^{4x}}$

27. C This is a geometric series with $r = \frac{x-1}{3}$. Convergence for $-1 < r < 1$. Thus the series is convergent for $-2 < x < 4$.


28. A $v = \left(\frac{2t+2}{t^2+2t}, 4t\right)$, $v(2) = \left(\frac{6}{8}, 8\right) = \left(\frac{3}{4}, 8\right)$

29. E Use the technique of antiderivatives by parts: $u = x$ and $dv = \sec^2 x dx$

$$\int x \sec^2 x dx = x \tan x - \int \tan x dx = x \tan x + \ln |\cos x| + C$$

30. C Each slice is a disk with radius $r = \sec x$ and width Δx .

$$\text{Volume} = \pi \int_0^{\pi/3} \sec^2 x dx = \pi \tan x \Big|_0^{\pi/3} = \pi\sqrt{3}$$



31. A $s_n = \frac{1}{5} \left(\frac{5+n}{4+n} \right)^{100}$, $\lim_{n \rightarrow \infty} s_n = \frac{1}{5} \cdot 1 = \frac{1}{5}$
32. B Only II is true. To see that neither I nor III must be true, let $f(x) = 1$ and let $g(x) = x^2 - \frac{128}{15}$ on the interval $[0, 5]$.
33. A The value of this integral is 2. Option A is also 2 and none of the others have a value of 2. Visualizing the graphs of $y = \sin x$ and $y = \cos x$ is a useful approach to the problem.
34. E Let $y = PR$ and $x = RQ$.
 $x^2 + y^2 = 40^2$, $2x \frac{dx}{dt} + 2y \frac{dy}{dt} = 0$, $x \cdot \frac{3}{4} \left(-\frac{dy}{dt} \right) + y \frac{dy}{dt} = 0 \Rightarrow y = \frac{3}{4}x$.
 Substitute into $x^2 + y^2 = 40^2$. $x^2 + \frac{9}{16}x^2 = 40^2$, $\frac{25}{16}x^2 = 40^2$, $x = 32$
35. A Apply the Mean Value Theorem to F . $F'(c) = \frac{F(b) - F(a)}{b - a} = \frac{0}{a} = 0$. This means that there is number in the interval (a, b) for which F' is zero. However, $F'(x) = f(x)$. So, $f(x) = 0$ for some number in the interval (a, b) .
36. E $v = \pi r^2 h$ and $h + 2\pi r = 30 \Rightarrow v = 2\pi(15r^2 - \pi r^3)$ for $0 < r < \frac{15}{\pi}$; $\frac{dv}{dr} = 6\pi r(10 - \pi r)$. The maximum volume is when $r = \frac{10}{\pi}$ because $\frac{dv}{dr} > 0$ on $\left(0, \frac{10}{\pi}\right)$ and $\frac{dv}{dr} < 0$ on $\left(\frac{10}{\pi}, \frac{15}{\pi}\right)$.
37. B $\int_0^e f(x) dx = \int_0^1 x dx + \int_1^e \frac{1}{x} dx = \frac{1}{2} + \ln e = \frac{3}{2}$
38. C $\frac{dN}{dt} = kN \Rightarrow N = Ce^{kt}$. $N(0) = 1000 \Rightarrow C = 1000$. $N(7) = 1200 \Rightarrow k = \frac{1}{7} \ln(1.2)$. Therefore $N(12) = 1000e^{\frac{12}{7} \ln(1.2)} \approx 1367$.
39. C Want $\frac{y(4) - y(1)}{4 - 1}$ where $y(x) = \ln|x| + C$. This gives $\frac{\ln 4 - \ln 1}{3} = \frac{1}{3} \ln 4 = \frac{1}{3} \ln 2^2 = \frac{2}{3} \ln 2$.
40. C The interval is $[0, 2]$, $x_0 = 0$, $x_1 = 1$, $x_2 = 2$. $S = \frac{1}{3}(0 + 4 \ln 2 + 0) = \frac{4}{3} \ln 2$. Note that Simpson's rule is no longer part of the BC Course Description.

41. C $f'(x) = (2x-3)e^{(x^2-3x)^2}$; $f' < 0$ for $x < \frac{3}{2}$ and $f' > 0$ for $x > \frac{3}{2}$.

Thus f has its absolute minimum at $x = \frac{3}{2}$.

42. E Suppose $\lim_{x \rightarrow 0} \ln\left((1+2x)^{\csc x}\right) = A$. The answer to the given question is e^A .

Use L'Hôpital's Rule: $\lim_{x \rightarrow 0} \ln\left((1+2x)^{\csc x}\right) = \lim_{x \rightarrow 0} \frac{\ln(1+2x)}{\sin x} = \lim_{x \rightarrow 0} \frac{2}{1+2x} \cdot \frac{1}{\cos x} = 2$.

43. A $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots \Rightarrow \sin x^2 = x^2 - \frac{(x^2)^3}{3!} + \frac{(x^2)^5}{5!} - \dots = x^2 - \frac{x^6}{3!} + \frac{x^{10}}{5!} - \dots$

44. E By the Intermediate Value Theorem there is a c satisfying $a < c < b$ such that $f(c)$ is equal to the average value of f on the interval $[a, b]$. But the average value is also given by $\frac{1}{b-a} \int_a^b f(x) dx$. Equating the two gives option E.

Alternatively, let $F(t) = \int_a^t f(x) dx$. By the Mean Value Theorem, there is a c satisfying

$a < c < b$ such that $\frac{F(b) - F(a)}{b - a} = F'(c)$. But $F(b) - F(a) = \int_a^b f(x) dx$, and $F'(c) = f(c)$ by the Fundamental Theorem of Calculus. This again gives option E as the answer. This result is called the Mean Value Theorem for Integrals.

45. D This is an infinite geometric series with a first term of $\sin^2 x$ and a ratio of $\sin^2 x$.

The series converges to $\frac{\sin^2 x}{1 - \sin^2 x} = \tan^2 x$ for $x \neq (2k+1)\frac{\pi}{2}$, k an integer. The answer is therefore $\tan^2 1 = 2.426$.

1997 Calculus AB Solutions: Part A

1. C $\int_1^2 (4x^3 - 6x) dx = (x^4 - 3x^2) \Big|_1^2 = (16 - 12) - (1 - 3) = 6$

2. A $f(x) = x(2x - 3)^{\frac{1}{2}}; f'(x) = (2x - 3)^{\frac{1}{2}} + x(2x - 3)^{-\frac{1}{2}} = (2x - 3)^{-\frac{1}{2}}(3x - 3) = \frac{(3x - 3)}{\sqrt{2x - 3}}$

3. C $\int_a^b (f(x) + 5) dx = \int_a^b f(x) dx + 5 \int_a^b 1 dx = a + 2b + 5(b - a) = 7b - 4a$

4. D $f(x) = -x^3 + x + \frac{1}{x}; f'(x) = -3x^2 + 1 - \frac{1}{x^2}; f'(-1) = -3(-1)^2 + 1 - \frac{1}{(-1)^2} = -3 + 1 - 1 = -3$

5. E $y = 3x^4 - 16x^3 + 24x^2 + 48; y' = 12x^3 - 48x^2 + 48x; y'' = 36x^2 - 96x + 48 = 12(3x - 2)(x - 2)$
 $y'' < 0$ for $\frac{2}{3} < x < 2$, therefore the graph is concave down for $\frac{2}{3} < x < 2$

6. C $\frac{1}{2} \int e^{\frac{t}{2}} dt = e^{\frac{t}{2}} + C$

7. D $\frac{d}{dx} \cos^2(x^3) = 2 \cos(x^3) \left(\frac{d}{dx} (\cos(x^3)) \right) = 2 \cos(x^3) (-\sin(x^3)) \left(\frac{d}{dx} (x^3) \right)$
 $= 2 \cos(x^3) (-\sin(x^3)) (3x^2)$

8. C The bug change direction when v changes sign. This happens at $t = 6$.

9. B Let A_1 be the area between the graph and t -axis for $0 \leq t \leq 6$, and let A_2 be the area between the graph and the t -axis for $6 \leq t \leq 8$. Then $A_1 = 12$ and $A_2 = 1$. The total distance is $A_1 + A_2 = 13$.

10. E $y = \cos(2x); y' = -2 \sin(2x); y' \left(\frac{\pi}{4} \right) = -2$ and $y \left(\frac{\pi}{4} \right) = 0; y = -2 \left(x - \frac{\pi}{4} \right)$

11. E Since f' is positive for $-2 < x < 2$ and negative for $x < -2$ and for $x > 2$, we are looking for a graph that is increasing for $-2 < x < 2$ and decreasing otherwise. Only option E.

12. B $y = \frac{1}{2}x^2; y' = x; \text{ We want } y' = \frac{1}{2} \Rightarrow x = \frac{1}{2}. \text{ So the point is } \left(\frac{1}{2}, \frac{1}{8} \right).$

1997 Calculus AB Solutions: Part A

13. A $f'(x) = \frac{|4-x^2|}{x-2}$; f is decreasing when $f' < 0$. Since the numerator is non-negative, this is only when the denominator is negative. Only when $x < 2$.
14. C $f(x) \approx L(x) = 2 + 5(x-3)$; $L(x) = 0$ if $0 = 5x - 13 \Rightarrow x = 2.6$
15. B Statement B is true because $\lim_{x \rightarrow a^-} f(x) = 2 = \lim_{x \rightarrow a^+} f(x)$. Also, $\lim_{x \rightarrow b} f(x)$ does not exist because the left- and right-sided limits are not equal, so neither (A), (C), nor (D) are true.
16. D The area of the region is given by $\int_{-2}^2 (5 - (x^2 + 1)) dx = 2 \left(4x - \frac{1}{3} x^3 \right) \Big|_0^2 = 2 \left(8 - \frac{8}{3} \right) = \frac{32}{3}$
17. A $x^2 + y^2 = 25$; $2x + 2y \cdot y' = 0$; $x + y \cdot y' = 0$; $y'(4, 3) = -\frac{4}{3}$;
 $x + y \cdot y' = 0 \Rightarrow 1 + y \cdot y'' + y' \cdot y' = 0$; $1 + (3)y'' + \left(-\frac{4}{3}\right) \cdot \left(-\frac{4}{3}\right) = 0$; $y'' = -\frac{25}{27}$
18. C $\int_0^{\frac{\pi}{4}} \frac{e^{\tan x}}{\cos^2 x} dx$ is of the form $\int e^u du$ where $u = \tan x$.
 $\int_0^{\frac{\pi}{4}} \frac{e^{\tan x}}{\cos^2 x} dx = e^{\tan x} \Big|_0^{\frac{\pi}{4}} = e^1 - e^0 = e - 1$
19. D $f(x) = \ln|x^2 - 1|$; $f'(x) = \frac{1}{x^2 - 1} \cdot \frac{d}{dx}(x^2 - 1) = \frac{2x}{x^2 - 1}$
20. E $\frac{1}{8} \int_{-3}^5 \cos x dx = \frac{1}{8} (\sin 5 - \sin(-3)) = \frac{1}{8} (\sin 5 + \sin 3)$; Note: Since the sine is an odd function, $\sin(-3) = -\sin(3)$.
21. E $\lim_{x \rightarrow 1} \frac{x}{\ln x}$ is nonexistent since $\lim_{x \rightarrow 1} \ln x = 0$ and $\lim_{x \rightarrow 1} x \neq 0$.
22. D $f(x) = (x^2 - 3)e^{-x}$; $f'(x) = e^{-x}(-x^2 + 2x + 3) = -e^{-x}(x-3)(x+1)$; $f'(x) > 0$ for $-1 < x < 3$
23. A Disks where $r = x$. $V = \pi \int_0^2 x^2 dy = \pi \int_0^2 y^4 dy = \frac{\pi}{5} y^5 \Big|_0^2 = \frac{32\pi}{5}$

1997 Calculus AB Solutions: Part A

24. B Let $[0,1]$ be divided into 50 subintervals. $\Delta x = \frac{1}{50}$; $x_1 = \frac{1}{50}, x_2 = \frac{2}{50}, x_3 = \frac{3}{50}, \dots, x_{50} = 1$

Using $f(x) = \sqrt{x}$, the right Riemann sum $\sum_{i=1}^{50} f(x_i)\Delta x$ is an approximation for $\int_0^1 \sqrt{x} dx$.

25. A Use the technique of antiderivatives by parts, which was removed from the AB Course Description in 1998.

$$u = x \quad dv = \sin 2x dx$$

$$du = dx \quad v = -\frac{1}{2} \cos 2x$$

$$\int x \sin(2x) dx = -\frac{1}{2} x \cos(2x) + \int \frac{1}{2} \cos(2x) dx = -\frac{1}{2} x \cos(2x) + \frac{1}{4} \sin(2x) + C$$

76. E $f(x) = \frac{e^{2x}}{2x}$; $f'(x) = \frac{2e^{2x} \cdot 2x - 2e^{2x}}{4x^2} = \frac{e^{2x}(2x-1)}{2x^2}$

77. D $y = x^3 + 6x^2 + 7x - 2 \cos x$. Look at the graph of $y'' = 6x + 12 + 2 \cos x$ in the window $[-3, -1]$ since that domain contains all the option values. y'' changes sign at $x = -1.89$.

78. D $F(3) - F(0) = \int_0^3 f(x) dx = \int_0^1 f(x) dx + \int_1^3 f(x) dx = 2 + 2.3 = 4.3$

(Count squares for $\int_0^1 f(x) dx$)

79. C The stem of the questions means $f'(2) = 5$. Thus f is differentiable at $x = 2$ and therefore continuous at $x = 2$. We know nothing of the continuity of f' . I and II only.


80. A $f(x) = 2e^{4x^2}$; $f'(x) = 16xe^{4x^2}$; We want $16xe^{4x^2} = 3$. Graph the derivative function and the function $y = 3$, then find the intersection to get $x = 0.168$.

81. A Let x be the distance of the train from the crossing. Then $\frac{dx}{dt} = 60$.

$$S^2 = x^2 + 70^2 \Rightarrow 2S \frac{dS}{dt} = 2x \frac{dx}{dt} \Rightarrow \frac{dS}{dt} = \frac{x}{S} \frac{dx}{dt}$$

After 4 seconds, $x = 240$ and so $S = 250$.

Therefore $\frac{dS}{dt} = \frac{240}{250}(60) = 57.6$



82. B $P(x) = 2x^2 - 8x$; $P'(x) = 4x - 8$; P' changes from negative to positive at $x = 2$. $P(2) = -8$

83. C $\cos x = x$ at $x = 0.739085$. Store this in A . $\int_0^A (\cos x - x) dx = 0.400$

84. C Cross sections are squares with sides of length y .

$$\text{Volume} = \int_1^e y^2 dx = \int_1^e \ln x dx = (x \ln x - x) \Big|_1^e = (e \ln e - e) - (0 - 1) = 1$$

85. C Look at the graph of f' and locate where the y changes from positive to negative. $x = 0.91$

86. A $f(x) = \sqrt{x}$; $f'(x) = \frac{1}{2\sqrt{x}}$; $\frac{1}{2\sqrt{c}} = 2 \cdot \frac{1}{2\sqrt{1}} \Rightarrow c = \frac{1}{4}$

1997 Calculus AB Solutions: Part B

87. B $a(t) = t + \sin t$ and $v(0) = -2 \Rightarrow v(t) = \frac{1}{2}t^2 - \cos t - 1$; $v(t) = 0$ at $t = 1.48$
88. E $f(x) = \int_a^x h(x)dx \Rightarrow f(a) = 0$, therefore only (A) or (E) are possible. But $f'(x) = h(x)$ and therefore f is differentiable at $x = b$. This is true for the graph in option (E) but not in option (A) where there appears to be a corner in the graph at $x = b$. Also, Since h is increasing at first, the graph of f must start out concave up. This is also true in (E) but not (A).
89. B $T = \frac{1}{2} \cdot \frac{1}{2} (3 + 2 \cdot 3 + 2 \cdot 5 + 2 \cdot 8 + 13) = 12$
90. D $F(x) = \frac{1}{2} \sin^2 x$ $F'(x) = \sin x \cos x$ Yes
- $F(x) = \frac{1}{2} \cos^2 x$ $F'(x) = -\cos x \sin x$ No
- $F(x) = -\frac{1}{4} \cos(2x)$ $F'(x) = \frac{1}{2} \sin(2x) = \sin x \cos x$ Yes

1997 Calculus BC Solutions: Part A

1. C $\int_0^1 \sqrt{x}(x+1) dx = \int_0^1 x^{\frac{3}{2}} + x^{\frac{1}{2}} dx = \frac{2}{5}x^{\frac{5}{2}} + \frac{2}{3}x^{\frac{3}{2}} \Big|_0^1 = \frac{16}{15}$
2. E $x = e^{2t}, y = \sin(2t); \frac{dy}{dx} = \frac{2 \cos(2t)}{2e^{2t}} = \frac{\cos(2t)}{e^{2t}}$
3. A $f(x) = 3x^5 - 4x^3 - 3x; f'(x) = 15x^4 - 12x^2 - 3 = 3(5x^2 + 1)(x^2 - 1) = 3(5x^2 + 1)(x+1)(x-1);$
 f' changes from positive to negative only at $x = -1$.
4. C $e^{\ln x^2} = x^2; \text{ so } xe^{\ln x^2} = x^3 \text{ and } \frac{d}{dx}(x^3) = 3x^2$
5. C $f(x) = (x-1)^{\frac{3}{2}} + \frac{1}{2}e^{x-2}; f'(x) = \frac{3}{2}(x-1)^{\frac{1}{2}} + \frac{1}{2}e^{x-2}; f'(2) = \frac{3}{2} + \frac{1}{2} = 2$
6. A $y = (16-x)^{\frac{1}{2}}; y' = -\frac{1}{2}(16-x)^{-\frac{1}{2}}; y'(0) = -\frac{1}{8};$ The slope of the normal line is 8.
7. C The slope at $x = 3$ is 2. The equation of the tangent line is $y - 5 = 2(x - 3)$.
8. E Points of inflection occur where f' changes from increasing to decreasing, or from decreasing to increasing. There are six such points.
9. A f increases for $0 \leq x \leq 6$ and decreases for $6 \leq x \leq 8$. By comparing areas it is clear that f increases more than it decreases, so the absolute minimum must occur at the left endpoint, $x = 0$.
10. B $y = xy + x^2 + 1; y' = xy' + y + 2x; \text{ at } x = -1, y = 1; y' = -y' + 1 - 2 \Rightarrow y' = -\frac{1}{2}$
11. C $\int_1^{\infty} x(1+x^2)^{-2} dx = \lim_{L \rightarrow \infty} -\frac{1}{2}(1+x^2)^{-1} \Big|_1^L = \lim_{L \rightarrow \infty} \frac{1}{4} - \frac{1}{2(1+L^2)} = \frac{1}{4}$
12. A f' changes from positive to negative once and from negative to positive twice. Thus one relative maximum and two relative minimums.
13. B $a(t) = 2t - 7$ and $v(0) = 6; \text{ so } v(t) = t^2 - 7t + 6 = (t-1)(t-6).$ Movement is right then left with the particle changing direction at $t = 1, 6$, therefore it will be farthest to the right at $t = 1$.

1997 Calculus BC Solutions: Part A

14. C Geometric Series. $r = \frac{3}{8} < 1 \Rightarrow$ convergence. $a = \frac{3}{2}$ so the sum will be $S = \frac{\frac{3}{2}}{1 - \frac{3}{8}} = 2.4$

15. D $x = \cos^3 t, y = \sin^3 t$ for $0 \leq t \leq \frac{\pi}{2}$. $L = \int_0^{\pi/2} \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$

$$L = \int_0^{\pi/2} \sqrt{(-3\cos^2 t \sin t)^2 + (3\sin^2 t \cos t)^2} dt = \int_0^{\pi/2} \sqrt{9\cos^4 t \sin^2 t + 9\sin^4 t \cos^2 t} dt$$

16. B $\lim_{h \rightarrow 0} \frac{e^h - 1}{2h} = \frac{1}{2} \lim_{h \rightarrow 0} \frac{e^h - e^0}{h} = \frac{1}{2} f'(0)$, where $f(x) = e^x$ and $f'(0) = 1$. $\lim_{h \rightarrow 0} \frac{e^h - 1}{2h} = \frac{1}{2}$

17. B $f(x) = \ln(3-x); f'(x) = \frac{1}{x-3}, f''(x) = -\frac{1}{(x-3)^2}, f'''(x) = \frac{2}{(x-3)^3};$

$$f(2) = 0, f'(2) = -1, f''(2) = -1, f'''(2) = -2; a_0 = 0, a_1 = -1, a_2 = -\frac{1}{2}, a_3 = -\frac{1}{3}$$

$$f(x) \approx -(x-2) - \frac{(x-2)^2}{2} - \frac{(x-2)^3}{3}$$


18. C $x = t^3 - t^2 - 1, y = t^4 + 2t^2 - 8t; \frac{dy}{dx} = \frac{4t^3 + 4t - 8}{3t^2 - 2t} = \frac{4t^3 + 4t - 8}{t(3t - 2)}$. Vertical tangents at $t = 0, \frac{2}{3}$

19. D $\int_{-4}^4 f(x) dx - 2 \int_{-1}^4 f(x) dx = (A_1 - A_2) - 2(-A_2) = A_1 + A_2$

20. E $\sum_{n=1}^{\infty} \frac{(x-2)^n}{n \cdot 3^n}$. The endpoints of the interval of convergence are when $(x-2) = \pm 3; x = -1, 5$.

Check endpoints: $x = -1$ gives the alternating harmonic series which converges. $x = 5$ gives the harmonic series which diverges. Therefore the interval is $-1 \leq x < 5$.

21. A Area = $2 \cdot \frac{1}{2} \int_0^{\pi/2} ((2\cos\theta)^2 - \cos^2\theta) d\theta = \int_0^{\pi/2} 3\cos^2\theta d\theta$



22. C $g'(x) = f(x)$. The only critical value of g on (a, d) is at $x = c$. Since g' changes from positive to negative at $x = c$, the absolute maximum for g occurs at this relative maximum.

1997 Calculus BC Solutions: Part A

23. E $x = 5 \sin \theta$; $\frac{dx}{dt} = 5 \cos \theta \cdot \frac{d\theta}{dt}$; When $x = 3$, $\cos \theta = \frac{4}{5}$; $\frac{dx}{dt} = 5 \left(\frac{4}{5} \right) (3) = 12$


24. D $f'(x) = \sin(x^2) = x^2 - \frac{(x^2)^3}{3!} + \dots = x^2 - \frac{1}{6}x^6 + \dots \Rightarrow f(x) = \frac{1}{3}x^3 - \frac{1}{42}x^7 + \dots$ The coefficient of x^7 is $-\frac{1}{42}$.

25. A This is the limit of a right Riemann sum of the function $f(x) = \sqrt{x}$ on the interval $[a, b]$, so

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \sqrt{x_i} \Delta x = \int_a^b \sqrt{x} \, dx = \frac{2}{3} x^{\frac{3}{2}} \Big|_a^b = \frac{2}{3} \left(b^{\frac{3}{2}} - a^{\frac{3}{2}} \right)$$

1997 Calculus BC Solutions: Part B

76. D Sequence I $\rightarrow \frac{5}{2}$; sequence II $\rightarrow \infty$; sequence III $\rightarrow 1$. Therefore I and III only.
77. E Use shells (which is no longer part of the AP Course Description.)



$$\sum 2\pi r h \Delta x \text{ where } r = x \text{ and}$$

$$h = 4x - x^2 - x$$

$$\text{Volume} =$$

$$2\pi \int_0^3 x(4x - x^2 - x) dx = 2\pi \int_0^3 (3x^2 - x^3) dx$$

78. A $\lim_{h \rightarrow 0} \frac{\ln(e+h) - 1}{h} = \lim_{h \rightarrow 0} \frac{\ln(e+h) - \ln e}{h} = f'(e)$ where $f(x) = \ln x$
79. D Count the number of places where the graph of $y(t)$ has a horizontal tangent line. Six places.
80. B Find the first turning point on the graph of $y = f'(x)$. Occurs at $x = 0.93$.
81. D f assumes every value between -1 and 3 on the interval $(-3, 6)$. Thus $f(c) = 1$ at least once.
82. B $\int_0^x (t^2 - 2t) dt \geq \int_2^x t dt$; $\frac{1}{3}x^3 - x^2 \geq \frac{1}{2}x^2 - 2$. Using the calculator, the greatest x value on the interval $[0, 4]$ that satisfies this inequality is found to occur at $x = 1.3887$.
83. E $\frac{dy}{y} = (1 + \ln x) dx$; $\ln|y| = x + x \ln x - x + k = x \ln x + k$; $|y| = e^k e^{x \ln x} \Rightarrow y = C e^{x \ln x}$. Since $y = 1$ when $x = 1$, $C = 1$. Hence $y = e^{x \ln x}$.

84. C $\int x^2 \sin x dx$; Use the technique of antiderivatives by parts with $u = x^2$ and $dv = \sin x dx$. It will take 2 iterations with a different choice of u and dv for the second iteration.

$$\begin{aligned}\int x^2 \sin x dx &= -x^2 \cos x + \int 2x \cos x dx \\ &= -x^2 \cos x + \left(2x \sin x - \int 2 \sin x dx\right) \\ &= -x^2 \cos x + 2x \sin x + 2 \cos x + C\end{aligned}$$

85. D I. Average rate of change of f is $\frac{f(3) - f(1)}{3 - 1} = \frac{5}{2}$. True
 II. Not enough information to determine the average value of f . False
 III. Average value of f' is the average rate of change of f . True

86. A Use partial fractions. $\frac{1}{(x-1)(x+3)} = \frac{A}{x-1} + \frac{B}{x+3}$; $1 = A(x+3) + B(x-1)$

Choose $x = 1 \Rightarrow A = \frac{1}{4}$ and choose $x = -3 \Rightarrow B = -\frac{1}{4}$.

$$\int \frac{1}{(x-1)(x+3)} dx = \frac{1}{4} \left[\int \frac{1}{x-1} dx - \int \frac{1}{x+3} dx \right] = \frac{1}{4} \ln \left| \frac{x-1}{x+3} \right| + C$$

87. B Squares with sides of length x . Volume $= \int_0^2 x^2 dy = \int_0^2 (2-y) dy$

88. C $f(x) = \int_0^{x^2} \sin t dt$; $f'(x) = 2x \sin(x^2)$; For the average rate of change of f we need to determine $f(0)$ and $f(\sqrt{\pi})$. $f(0) = 0$ and $f(\sqrt{\pi}) = \int_0^{\pi} \sin t dt = 2$. The average rate of change of f on the interval is $\frac{2}{\sqrt{\pi}}$. See how many points of intersection there are for the graphs of $y = 2x \sin(x^2)$ and $y = \frac{2}{\sqrt{\pi}}$ on the interval $[0, \sqrt{\pi}]$. There are two.

$$89. \quad D \quad f(x) = \int_1^x \frac{t^2}{1+t^5} dt; \quad f(4) = \int_1^4 \frac{t^2}{1+t^5} dt = 0.376$$

$$\text{Or, } f(4) = f(1) + \int_1^4 \frac{x^2}{1+x^5} dx = 0.376$$

Both statements follow from the Fundamental Theorem of Calculus.

$$90. \quad B \quad F(x) = kx; \quad 10 = 4k \Rightarrow k = \frac{5}{2}; \quad \text{Work} = \int_0^6 F(x) dx = \int_0^6 \frac{5}{2} x dx = \frac{5}{4} x^2 \Big|_0^6 = 45 \text{ inch-lbs}$$

1. D $y' = x^2 + 10x$; $y'' = 2x + 10$; y'' changes sign at $x = -5$
2. B
$$\int_{-1}^4 f(x) dx = \int_{-1}^2 f(x) dx + \int_2^4 f(x) dx$$

 $= \text{Area of trapezoid(1)} - \text{Area of trapezoid(2)} = 4 - 1.5 = 2.5$
3. C
$$\int_1^2 \frac{1}{x^2} dx = \int_1^2 x^{-2} dx = -x^{-1} \Big|_1^2 = \frac{1}{2}$$
4. B This would be false if f was a linear function with non-zero slope.
5. E
$$\int_0^x \sin t dt = -\cos t \Big|_0^x = -\cos x - (-\cos 0) = -\cos x + 1 = 1 - \cos x$$
6. A Substitute $x = 2$ into the equation to find $y = 3$. Taking the derivative implicitly gives

$$\frac{d}{dx}(x^2 + xy) = 2x + xy' + y = 0$$
. Substitute for x and y and solve for y' .

$$4 + 2y' + 3 = 0; y' = -\frac{7}{2}$$
7. E
$$\int_1^e \frac{x^2 - 1}{x} dx = \int_1^e x - \frac{1}{x} dx = \left(\frac{1}{2}x^2 - \ln x \right) \Big|_1^e = \left(\frac{1}{2}e^2 - 1 \right) - \left(\frac{1}{2} - 0 \right) = \frac{1}{2}e^2 - \frac{3}{2}$$
8. E $h(x) = f(x)g(x)$ so, $h'(x) = f'(x)g(x) + f(x)g'(x)$. It is given that $h'(x) = f(x)g'(x)$.
 Thus, $f'(x)g(x) = 0$. Since $g(x) > 0$ for all x , $f'(x) = 0$. This means that f is constant. It is given that $f(0) = 1$, therefore $f(x) = 1$.
9. D Let $r(t)$ be the rate of oil flow as given by the graph, where t is measured in hours. The total number of barrels is given by $\int_0^{24} r(t) dt$. This can be approximated by counting the squares below the curve and above the horizontal axis. There are approximately five squares with area 600 barrels. Thus the total is about 3,000 barrels.
10. D
$$f'(x) = \frac{(x-1)(2x) - (x^2-2)(1)}{(x-1)^2}; f'(2) = \frac{(2-1)(4) - (4-2)(1)}{(2-1)^2} = 2$$
11. A Since f is linear, its second derivative is zero. The integral gives the area of a rectangle with zero height and width $(b-a)$. This area is zero.

1998 Calculus AB Solutions: Part A

12. E $\lim_{x \rightarrow 2^-} f(x) = \ln 2 \neq 4 \ln 2 = \lim_{x \rightarrow 2^+} f(x)$. Therefore the limit does not exist.
13. B At $x = 0$ and $x = 2$ only. The graph has a non-vertical tangent line at every other point in the interval and so has a derivative at each of these other x 's.
14. C $v(t) = 2t - 6$; $v(t) = 0$ for $t = 3$
15. D By the Fundamental Theorem of Calculus, $F'(x) = \sqrt{x^3 + 1}$, thus $F'(2) = \sqrt{2^3 + 1} = \sqrt{9} = 3$.
16. E $f'(x) = \cos(e^{-x}) \cdot \frac{d}{dx}(e^{-x}) = \cos(e^{-x}) \left(e^{-x} \cdot \frac{d}{dx}(-x) \right) = -e^{-x} \cos(e^{-x})$
17. D From the graph $f(1) = 0$. Since $f'(1)$ represents the slope of the graph at $x = 1$, $f'(1) > 0$. Also, since $f''(1)$ represents the concavity of the graph at $x = 1$, $f''(1) < 0$.
18. B $y' = 1 - \sin x$ so $y'(0) = 1$ and the line with slope 1 containing the point $(0, 1)$ is $y = x + 1$.
19. C Points of inflection occur where f'' changes sign. This is only at $x = 0$ and $x = -1$. There is no sign change at $x = 2$.
20. A $\int_{-3}^k x^2 dx = \frac{1}{3} x^3 \Big|_{-3}^k = \frac{1}{3} (k^3 - (-3)^3) = \frac{1}{3} (k^3 + 27) = 0$ only when $k = -3$.
21. B The solution to this differential equation is known to be of the form $y = y(0) \cdot e^{kt}$. Option (B) is the only one of this form. If you do not remember the form of the solution, then separate the variables and antidifferentiate.

$$\frac{dy}{y} = k dt; \ln |y| = kt + c_1; |y| = e^{kt+c_1} = e^{kt} e^{c_1}; y = ce^{kt}.$$
22. C f is increasing on any interval where $f'(x) > 0$. $f'(x) = 4x^3 + 2x = 2x(2x^2 + 1) > 0$. Since $(x^2 + 1) > 0$ for all x , $f'(x) > 0$ whenever $x > 0$.
23. A The graph shows that f is increasing on an interval (a, c) and decreasing on the interval (c, b) , where $a < c < b$. This means the graph of the derivative of f is positive on the interval (a, c) and negative on the interval (c, b) , so the answer is (A) or (E). The derivative is not (E), however, since then the graph of f would be concave down for the entire interval.

1998 Calculus AB Solutions: Part A

24. D The maximum acceleration will occur when its derivative changes from positive to negative or at an endpoint of the interval. $a(t) = v'(t) = 3t^2 - 6t + 12 = 3(t^2 - 2t + 4)$ which is always positive. Thus the acceleration is always increasing. The maximum must occur at $t = 3$ where $a(3) = 21$
25. D The area is given by $\int_0^2 x^2 - (-x) dx = \left(\frac{1}{3}x^3 + \frac{1}{2}x^2 \right) \Big|_0^2 = \frac{8}{3} + 2 = \frac{14}{3}$.
26. A Any value of k less than $1/2$ will require the function to assume the value of $1/2$ at least twice because of the Intermediate Value Theorem on the intervals $[0,1]$ and $[1,2]$. Hence $k = 0$ is the only option.
27. A $\frac{1}{2} \int_0^2 x^2 \sqrt{x^3 + 1} dx = \frac{1}{2} \int_0^2 (x^3 + 1)^{\frac{1}{2}} \left(\frac{1}{3} \cdot 3x^2 \right) dx = \frac{1}{2} \cdot \frac{1}{3} \cdot \frac{2}{3} (x^3 + 1)^{\frac{3}{2}} \Big|_0^2 = \frac{1}{9} \left(9^{\frac{3}{2}} - 1^{\frac{3}{2}} \right) = \frac{26}{9}$
28. E $f'(x) = \sec^2(2x) \cdot \frac{d}{dx}(2x) = 2 \sec^2(2x)$; $f'\left(\frac{\pi}{6}\right) = 2 \sec^2\left(\frac{\pi}{3}\right) = 2(4) = 8$

76. A From the graph it is clear that f is not continuous at $x = a$. All others are true.
77. C Parallel tangents will occur when the slopes of f and g are equal. $f'(x) = 6e^{2x}$ and $g'(x) = 18x^2$. The graphs of these derivatives reveal that they are equal only at $x = -0.391$.
78. B $A = \pi r^2 \Rightarrow \frac{dA}{dt} = 2\pi r \frac{dr}{dt}$. However, $C = 2\pi r$ and $\frac{dr}{dt} = -0.1$. Thus $\frac{dA}{dt} = -0.1C$.
79. A The graph of the derivative would have to change from positive to negative. This is only true for the graph of f' .
80. B Look at the graph of $f'(x)$ on the interval $(0,10)$ and count the number of x -intercepts in the interval.
81. D Only II is false since the graph of the absolute value function has a sharp corner at $x = 0$.
82. E Since F is an antiderivative of f , $\int_1^3 f(2x) dx = \frac{1}{2} F(2x) \Big|_1^3 = \frac{1}{2} (F(6) - F(2))$
83. B $\lim_{x \rightarrow a} \frac{x^2 - a^2}{x^4 - a^4} = \lim_{x \rightarrow a} \frac{x^2 - a^2}{(x^2 - a^2)(x^2 + a^2)} = \lim_{x \rightarrow a} \frac{1}{x^2 + a^2} = \frac{1}{2a^2}$
84. A A known solution to this differential equation is $y(t) = y(0)e^{kt}$. Use the fact that the population is $2y(0)$ when $t = 10$. Then $2y(0) = y(0)e^{k(10)} \Rightarrow e^{10k} = 2 \Rightarrow k = (0.1) \ln 2 = 0.069$
85. C There are 3 trapezoids. $\frac{1}{2} \cdot 3(f(2) + f(5)) + \frac{1}{2} \cdot 2(f(5) + f(7)) + \frac{1}{2} \cdot 1(f(7) + f(8))$
86. C Each cross section is a semicircle with a diameter of y . The volume would be given by $\int_0^8 \frac{1}{2} \pi \left(\frac{y}{2}\right)^2 dx = \frac{\pi}{8} \int_0^8 \left(\frac{8-x}{2}\right)^2 dx = 16.755$
87. D Find the x for which $f'(x) = 1$. $f'(x) = 4x^3 + 4x = 1$ only for $x = 0.237$. Then $f(0.237) = 0.115$. So the equation is $y - 0.115 = x - 0.237$. This is equivalent to option (D).

88. C $F(9) - F(1) = \int_1^9 \frac{(\ln t)^3}{t} dt = 5.827$ using a calculator. Since $F(1) = 0$, $F(9) = 5.827$.

Or solve the differential equation with an initial condition by finding an antiderivative for $\frac{(\ln x)^3}{x}$. This is of the form $u^3 du$ where $u = \ln x$. Hence $F(x) = \frac{1}{4}(\ln x)^4 + C$ and since $F(1) = 0$, $C = 0$. Therefore $F(9) = \frac{1}{4}(\ln 9)^4 = 5.827$

89. B The graph of $y = x^2 - 4$ is a parabola that changes from positive to negative at $x = -2$ and from negative to positive at $x = 2$. Since g is always negative, f' changes sign opposite to the way $y = x^2 - 4$ does. Thus f has a relative minimum at $x = -2$ and a relative maximum at $x = 2$.

90. D The area of a triangle is given by $A = \frac{1}{2}bh$. Taking the derivative with respect to t of both sides of the equation yields $\frac{dA}{dt} = \frac{1}{2}\left(\frac{db}{dt} \cdot h + b \cdot \frac{dh}{dt}\right)$. Substitute the given rates to get $\frac{dA}{dt} = \frac{1}{2}(3h - 3b) = \frac{3}{2}(h - b)$. The area will be decreasing whenever $\frac{dA}{dt} < 0$. This is true whenever $b > h$.

91. E I. True. Apply the Intermediate Value Theorem to each of the intervals $[2, 5]$ and $[5, 9]$.

II. True. Apply the Mean Value Theorem to the interval $[2, 9]$.

III. True. Apply the Intermediate Value Theorem to the interval $[2, 5]$.

92. D $\int_k^{\frac{\pi}{2}} \cos x dx = 0.1 \Rightarrow \sin\left(\frac{\pi}{2}\right) - \sin k = 0.1 \Rightarrow \sin k = 0.9$. Therefore $k = \sin^{-1}(0.9) = 1.120$.

1998 Calculus BC Solutions: Part A

1. C f will be increasing when its derivative is positive.
 $f'(x) = 3x^2 + 6x - 9 = 3(x^2 + 2x - 3)$ $f'(x) = 3(x+3)(x-1) > 0$ for $x < -3$ or $x > 1$.
2. A $\frac{dx}{dt} = 5$ and $\frac{dy}{dt} = 3 \Rightarrow \frac{dy}{dx} = \frac{3}{5}$
3. D Find the derivative implicitly and substitute. $2y \cdot y' + 3(xy + 1)^2(x \cdot y' + y) = 0$;
 $2(-1) \cdot y' + 3((2)(-1) + 1)^2((2) \cdot y' + (-1)) = 0$; $-2y' + 6 \cdot y' - 3 = 0$; $y' = \frac{3}{4}$
4. A Use partial fractions. $\frac{1}{x^2 - 6x + 8} = \frac{1}{(x-4)(x-2)} = \frac{1}{2} \left(\frac{1}{x-4} - \frac{1}{x-2} \right)$
 $\int \frac{1}{x^2 - 6x + 8} dx = \frac{1}{2} (\ln|x-4| - \ln|x-2|) + C = \frac{1}{2} \ln \left| \frac{x-4}{x-2} \right| + C$
5. A $h'(x) = f'(g(x)) \cdot g'(x)$; $h''(x) = f''(g(x)) \cdot g'(x) \cdot g'(x) + f'(g(x)) \cdot g''(x)$
 $h''(x) = f''(g(x)) \cdot (g'(x))^2 + f'(g(x)) \cdot g''(x)$
6. E The graph of h has 2 turning points and one point of inflection. The graph of h' will have 2 x -intercepts and one turning point. Only (C) and (E) are possible answers. Since the first turning point on the graph of h is a relative maximum, the first zero of h' must be a place where the sign changes from positive to negative. This is option (E).
7. E $\int_1^e \frac{x^2 - 1}{x} dx = \int_1^e x - \frac{1}{x} dx = \left(\frac{1}{2}x^2 - \ln x \right) \Big|_1^e = \left(\frac{1}{2}e^2 - 1 \right) - \left(\frac{1}{2} - 0 \right) = \frac{1}{2}e^2 - \frac{3}{2}$
8. B $y(x) = -\frac{1}{3}(\cos x)^3 + C$; Let $x = \frac{\pi}{2}$, $0 = -\frac{1}{3} \left(\cos \frac{\pi}{2} \right)^3 + C \Rightarrow C = 0$. $y(0) = -\frac{1}{3}(\cos 0)^3 = -\frac{1}{3}$
9. D Let $r(t)$ be the rate of oil flow as given by the graph, where t is measured in hours. The total number of barrels is given by $\int_0^{24} r(t) dt$. This can be approximated by counting the squares below the curve and above the horizontal axis. There are approximately five squares with area 600 barrels. Thus the total is about 3,000 barrels.
10. E $v(t) = (3t^2 - 1, 6(2t - 1)^2)$ and $a(t) = (6t, 24(2t - 1)) \Rightarrow a(1) = (6, 24)$

1998 Calculus BC Solutions: Part A

11. A Since f is linear, its second derivative is zero and the integral gives the area of a rectangle with zero height and width $(b-a)$. This area is zero.

12. E $\lim_{x \rightarrow 2^-} f(x) = \ln 2 \neq 4 \ln 2 = \lim_{x \rightarrow 2^+} f(x)$. Therefore the limit does not exist.

13. B At $x = 0$ and $x = 2$ only. The graph has a non-vertical tangent line at every other point in the interval and so has a derivative at each of these other x 's.

14. E $\sin x \approx x - \frac{x^3}{3!} + \frac{x^5}{5!}$; $\sin 1 \approx 1 - \frac{1^3}{3!} + \frac{1^5}{5!} = 1 - \frac{1}{6} + \frac{1}{120}$

15. B Use the technique of antiderivatives by parts. Let $u = x$ and $dv = \cos x \, dx$.

$$\int x \cos x \, dx = x \sin x - \int \sin x \, dx = x \sin x + \cos x + C$$

16. C Inflection point will occur when f'' changes sign. $f'(x) = 15x^4 - 20x^3$.
 $f''(x) = 60x^3 - 60x^2 = 60x^2(x-1)$. The only sign change is at $x = 1$.

17. D From the graph $f(1) = 0$. Since $f'(1)$ represents the slope of the graph at $x = 1$, $f'(1) > 0$. Also, since $f''(1)$ represents the concavity of the graph at $x = 1$, $f''(1) < 0$.

18. B
 I. Divergent. The limit of the n th term is not zero.
 II. Convergent. This is the same as the alternating harmonic series.
 III. Divergent. This is the harmonic series.

19. D Find the points of intersection of the two curves to determine the limits of integration.

$$4 \sin \theta = 2 \text{ when } \sin \theta = 0.5; \text{ this is at } \theta = \frac{\pi}{6} \text{ and } \frac{5\pi}{6}. \text{ Area} = \frac{1}{2} \int_{\frac{\pi}{6}}^{\frac{5\pi}{6}} \left((4 \sin \theta)^2 - (2)^2 \right) d\theta$$

20. E $\left. \frac{d(\sqrt[3]{x})}{dt} \right|_{x=8} = \frac{1}{3} x^{-\frac{2}{3}} \cdot \frac{dx}{dt} \Big|_{x=8} = \frac{1}{3} (8)^{-\frac{2}{3}} \cdot \frac{dx}{dt} = \frac{1}{3} \cdot \frac{1}{4} \cdot \frac{dx}{dt} = \frac{1}{12} \cdot \frac{dx}{dt} \Rightarrow k = 12$

21. C The length of this parametric curve is given by $\int_0^1 \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = \int_0^1 \sqrt{(t^2)^2 + t^2} dt$.

22. A This is the integral test applied to the series in (A). Thus the series in (A) converges. None of the others must be true.

23. E I. False. The relative maximum could be at a cusp.
 II. True. There is a critical point at $x = c$ where $f'(c)$ exists
 III. True. If $f''(c) > 0$, then there would be a relative minimum, not maximum
24. C All slopes along the diagonal $y = -x$ appear to be 0. This is consistent only with option (C). For each of the others you can see why they do not work. Option (A) does not work because all slopes at points with the same x coordinate would have to be equal. Option (B) does not work because all slopes would have to be positive. Option (D) does not work because all slopes in the third quadrant would have to be positive. Option (E) does not work because there would only be slopes for $y > 0$.
25. C
$$\int_0^{\infty} x^2 e^{-x^3} dx = \lim_{b \rightarrow \infty} \int_0^b x^2 e^{-x^3} dx = \lim_{b \rightarrow \infty} \left. -\frac{1}{3} e^{-x^3} \right|_0^b = \frac{1}{3}.$$
26. E As $\lim_{t \rightarrow \infty} \frac{dP}{dt} = 0$ for a population satisfying a logistic differential equation, this means that $P \rightarrow 10,000$.
27. D If $f(x) = \sum_{n=0}^{\infty} a_n x^n$, then $f'(x) = \sum_{n=0}^{\infty} n a_n x^{n-1} = \sum_{n=1}^{\infty} n a_n x^{n-1}$.

$$f'(1) = \sum_{n=1}^{\infty} n a_n 1^{n-1} = \sum_{n=1}^{\infty} n a_n$$
28. C Apply L'Hôpital's rule.
$$\lim_{x \rightarrow 1} \frac{\int_1^x e^{t^2} dt}{x^2 - 1} = \lim_{x \rightarrow 1} \frac{e^{x^2}}{2x} = \frac{e}{2}$$

1998 Calculus BC Solutions: Part B

76. D The first series is either the harmonic series or the alternating harmonic series depending on whether k is odd or even. It will converge if k is odd. The second series is geometric and will converge if $k < 4$.
77. E $f'(t) = (-e^{-t}, -\sin t)$; $f''(t) = (e^{-t}, -\cos t)$.
78. B $A = \pi r^2 \Rightarrow \frac{dA}{dt} = 2\pi r \frac{dr}{dt}$. However, $C = 2\pi r$ and $\frac{dr}{dt} = -0.1$. Thus $\frac{dA}{dt} = -0.1C$.
79. A None. For every positive value of a the denominator will be zero for some value of x .
80. B The area is given by $\int_{-\frac{2}{3}}^{\frac{2}{3}} (1 + \ln(\cos^4 x)) dx = 0.919$
81. B $\frac{dy}{dx} = \sqrt{1-y^2}$; $\frac{d^2y}{dx^2} = \frac{d}{dx} \left((1-y^2)^{\frac{1}{2}} \right) = \frac{1}{2} (1-y^2)^{-\frac{1}{2}} \cdot (-2y) \cdot \frac{dy}{dx} = -y$
82. B $\int_3^5 [f(x) + g(x)] dx = \int_3^5 [2g(x) + 7] dx = 2 \int_3^5 g(x) dx + (7)(2) = 2 \int_3^5 g(x) dx + 14$
83. C Use a calculator. The maximum for $\left| \ln x - \left(\frac{(x-1)}{1} - \frac{(x-1)^2}{2} + \frac{(x-1)^3}{3} \right) \right|$ on the interval $0.3 \leq x \leq 1.7$ occurs at $x = 0.3$.
84. B You may use the ratio test. However, the series will converge if the numerator is $(-1)^n$ and diverge if the numerator is 1^n . Any value of x for which $|x+2| > 1$ in the numerator will make the series diverge. Hence the interval is $-3 \leq x < -1$.
85. C There are 3 trapezoids. $\frac{1}{2} \cdot 3(f(2) + f(5)) + \frac{1}{2} \cdot 2(f(5) + f(7)) + \frac{1}{2} \cdot 1(f(7) + f(8))$
86. C Each cross section is a semicircle with a diameter of y . The volume would be given by $\int_0^8 \frac{1}{2} \pi \left(\frac{y}{2} \right)^2 dx = \frac{\pi}{8} \int_0^8 \left(\frac{8-x}{2} \right)^2 dx = 16.755$

1998 Calculus BC Solutions: Part B

87. D Find the x for which $f'(x) = 1$. $f'(x) = 4x^3 + 4x = 1$ only for $x = 0.237$. Then $f(0.237) = 0.115$. So the equation is $y - 0.115 = x - 0.237$. This is equivalent to option (D).
88. C From the given information, f is the derivative of g . We want a graph for f that represents the slopes of the graph g . The slope of g is zero at a and b . Also the slope of g changes from positive to negative at one point between a and b . This is true only for figure (C).
89. A The series is the Maclaurin expansion of e^{-x} . Use the calculator to solve $e^{-x} = x^3$.
90. A Constant acceleration means linear velocity which in turn leads to quadratic position. Only the graph in (A) is quadratic with initial $s = 2$.
91. E $v(t) = 11 + \int_0^t a(x) dx \approx 11 + [2 \cdot 5 + 2 \cdot 2 + 2 \cdot 8] = 41$ ft/sec.
92. D $f'(x) = 2x - 2$, $f'(2) = 2$, and $f(2) = 3$, so an equation for the tangent line is $y = 2x - 1$. The difference between the function and the tangent line is represented by $(x - 2)^2$. Solve $(x - 2)^2 < 0.5$. This inequality is satisfied for all x such that $2 - \sqrt{0.5} < x < 2 + \sqrt{0.5}$. This is the same as $1.293 < x < 2.707$. Thus the largest value in the list that satisfies the inequality is 2.7.