

Indefinite Integrals

73 1. $\int (x^3 - 3x) dx =$

(A) $3x^2 - 3 + C$

(B) $4x^4 - 6x^2 + C$

(C) $\frac{x^4}{3} - 3x^2 + C$

(D) $\frac{x^4}{4} - 3x + C$

(E) $\frac{x^4}{4} - \frac{3x^2}{2} + C$

93 17. $\int (x^2 + 1)^2 dx =$

(A) $\frac{(x^2 + 1)^3}{3} + C$

(B) $\frac{(x^2 + 1)^3}{6x} + C$

(C) $\left(\frac{x^3}{3} + x\right)^2 + C$

(D) $\frac{2x(x^2 + 1)^3}{3} + C$

(E) $\frac{x^5}{5} + \frac{2x^3}{3} + x + C$

97 6. $\frac{1}{2} \int e^{\frac{t}{2}} dt =$

(A) $e^{-t} + C$

(B) $e^{\frac{t}{2}} + C$

(C) $e^{\frac{t}{2}} + C$

(D) $2e^{\frac{t}{2}} + C$

(E) $e^t + C$

69 38. $\int \frac{x^2}{e^{x^3}} dx =$

(A) $-\frac{1}{3} \ln e^{x^3} + C$

(B) $-\frac{e^{x^3}}{3} + C$

(C) $-\frac{1}{3e^{x^3}} + C$

(D) $\frac{1}{3} \ln e^{x^3} + C$

(E) $\frac{x^3}{3e^{x^3}} + C$

88 19. $\int_2^3 \frac{x}{x^2+1} dx =$

(A) $\frac{1}{2} \ln \frac{3}{2}$

(B) $\frac{1}{2} \ln 2$

(C) $\ln 2$

(D) $2 \ln 2$

(E) $\frac{1}{2} \ln 5$

73
82 20. $\int x\sqrt{4-x^2} dx =$

(A) $\frac{(4-x^2)^{3/2}}{3} + C$

(B) $-(4-x^2)^{3/2} + C$

(C) $\frac{x^2(4-x^2)^{3/2}}{3} + C$

(D) $-\frac{x^2(4-x^2)^{3/2}}{3} + C$

(E) $-\frac{(4-x^2)^{3/2}}{3} + C$

88 7. $\int \frac{x dx}{\sqrt{3x^2+5}} =$

(A) $\frac{1}{9}(3x^2+5)^{\frac{3}{2}} + C$

(B) $\frac{1}{4}(3x^2+5)^{\frac{3}{2}} + C$

(C) $\frac{1}{12}(3x^2+5)^{\frac{1}{2}} + C$

(D) $\frac{1}{3}(3x^2+5)^{\frac{1}{2}} + C$

(E) $\frac{3}{2}(3x^2+5)^{\frac{1}{2}} + C$

93 14. $\int \frac{3x^2}{\sqrt{x^3+1}} dx =$

(A) $2\sqrt{x^3+1} + C$

(B) $\frac{3}{2}\sqrt{x^3+1} + C$

(C) $\sqrt{x^3+1} + C$

(D) $\ln\sqrt{x^3+1} + C$

(E) $\ln(x^3+1) + C$

69 43. $\int \sin(2x+3) dx =$

(A) $\frac{1}{2}\cos(2x+3) + C$

(B) $\cos(2x+3) + C$

(C) $-\cos(2x+3) + C$

(D) $-\frac{1}{2}\cos(2x+3) + C$

(E) $-\frac{1}{5}\cos(2x+3) + C$

97 25. $\int x \sin(2x) dx =$

(A) $-\frac{x}{2} \cos(2x) + \frac{1}{4} \sin(2x) + C$

(B) $-\frac{x}{2} \cos(2x) - \frac{1}{4} \sin(2x) + C$

(C) $\frac{x}{2} \cos(2x) - \frac{1}{4} \sin(2x) + C$

(D) $-2x \cos(2x) + \sin(2x) + C$

(E) $-2x \cos(2x) - 4 \sin(2x) + C$

93 38. If the second derivative of f is given by $f''(x) = 2x - \cos x$, which of the following could be $f(x)$?

(A) $\frac{x^3}{3} + \cos x - x + 1$

(B) $\frac{x^3}{3} - \cos x - x + 1$

(C) $x^3 + \cos x - x + 1$

(D) $x^2 - \sin x + 1$

(E) $x^2 + \sin x + 1$

85 30. $\int \tan(2x) dx =$

(A) $-2 \ln |\cos(2x)| + C$

(B) $-\frac{1}{2} \ln |\cos(2x)| + C$

(C) $\frac{1}{2} \ln |\cos(2x)| + C$

(D) $2 \ln |\cos(2x)| + C$

(E) $\frac{1}{2} \sec(2x) \tan(2x) + C$

88 5. $\int \sec^2 x dx =$

(A) $\tan x + C$

(B) $\csc^2 x + C$

(C) $\cos^2 x + C$

(D) $\frac{\sec^3 x}{3} + C$

(E) $2 \sec^2 x \tan x + C$

69
86

27. If $\frac{dy}{dx} = \tan x$, then $y =$

(A) $\frac{1}{2}\tan^2 x + C$

(B) $\sec^2 x + C$

(C) $\ln|\sec x| + C$

(D) $\ln|\cos x| + C$

(E) $\sec x \tan x + C$

Definite Integrals

93
66

44. If f is continuous on the interval $[a, b]$, then there exists c such that $a < c < b$ and $\int_a^b f(x) dx =$

- (A) $\frac{f(c)}{b-a}$ (B) $\frac{f(b)-f(a)}{b-a}$ (C) $f(b)-f(a)$ (D) $f'(c)(b-a)$ (E) $f(c)(b-a)$
-

73

20. If F and f are continuous functions such that $F'(x) = f(x)$ for all x , then $\int_a^b f(x) dx$ is

- (A) $F'(a) - F'(b)$
(B) $F'(b) - F'(a)$
(C) $F(a) - F(b)$
(D) $F(b) - F(a)$
(E) none of the above

97

1. $\int_1^2 (4x^3 - 6x) dx =$

- (A) 2
(B) 4
(C) 6
(D) 36
(E) 42
-

88
8C

2. $\int_0^1 x(x^2 + 2)^2 dx =$

- (A) $\frac{19}{2}$ (B) $\frac{19}{3}$ (C) $\frac{9}{2}$ (D) $\frac{19}{6}$ (E) $\frac{1}{6}$
-

98

3. $\int_1^2 \frac{1}{x^2} dx =$

- (A) $-\frac{1}{2}$ (B) $\frac{7}{24}$ (C) $\frac{1}{2}$ (D) 1 (E) $2 \ln 2$

85

1. $\int_1^2 x^{-3} dx =$

- (A) $-\frac{7}{8}$ (B) $-\frac{3}{4}$ (C) $\frac{15}{64}$ (D) $\frac{3}{8}$ (E) $\frac{15}{16}$

85
BC

36. $\int_{-1}^1 \frac{3}{x^2} dx$ is

- (A) -6 (B) -3 (C) 0 (D) 6 (E) nonexistent
-

85

22. $\int_1^2 \frac{x^2-1}{x+1} dx =$

- (A) $\frac{1}{2}$ (B) 1 (C) 2 (D) $\frac{5}{2}$ (E) $\ln 3$
-

69
AB

26. $\int_0^1 \sqrt{x^2-2x+1} dx$ is

- (A) -1
(B) $-\frac{1}{2}$
(C) $\frac{1}{2}$
(D) 1
(E) none of the above
-

73
BC

2. $\int_0^3 (x+1)^{1/2} dx =$

- (A) $\frac{21}{2}$ (B) 7 (C) $\frac{16}{3}$ (D) $\frac{14}{3}$ (E) $-\frac{1}{4}$
-

69
AB

4. $\int_0^8 \frac{dx}{\sqrt{1+x}} =$

- (A) 1 (B) $\frac{3}{2}$ (C) 2 (D) 4 (E) 6
-

88

17. $\int_0^1 (3x-2)^2 dx =$

- (A) $-\frac{7}{3}$ (B) $-\frac{7}{9}$ (C) $\frac{1}{9}$ (D) 1 (E) 3
-

73 27. $\int_0^{1/2} \frac{2x}{\sqrt{1-x^2}} dx =$

- (A) $1 - \frac{\sqrt{3}}{2}$ (B) $\frac{1}{2} \ln \frac{3}{4}$ (C) $\frac{\pi}{6}$ (D) $\frac{\pi}{6} - 1$ (E) $2 - \sqrt{3}$

73 25. $\int_0^{\pi/4} \tan^2 x dx =$

- (A) $\frac{\pi}{4} - 1$ (B) $1 - \frac{\pi}{4}$ (C) $\frac{1}{3}$ (D) $\sqrt{2} - 1$ (E) $\frac{\pi}{4} + 1$

85 32. $\int_0^{\pi} \sin(3x) dx =$

- (A) -2 (B) $-\frac{2}{3}$ (C) 0 (D) $\frac{2}{3}$ (E) 2

93 33. Which of the following is equal to $\int_0^{\pi} \sin x dx$?

- (A) $\int_{\frac{\pi}{2}}^{\frac{\pi}{2}} \cos x dx$ (B) $\int_0^{\pi} \cos x dx$ (C) $\int_{-\pi}^0 \sin x dx$
(D) $\int_{\frac{\pi}{2}}^{\frac{\pi}{2}} \sin x dx$ (E) $\int_{\pi}^{2\pi} \sin x dx$

98 5. $\int_0^x \sin t dt =$

- (A) $\sin x$ (B) $-\cos x$ (C) $\cos x$ (D) $\cos x - 1$ (E) $1 - \cos x$

88 14. $\int_0^{\frac{\pi}{2}} \frac{\cos \theta}{\sqrt{1 + \sin \theta}} d\theta =$

- (A) $-2(\sqrt{2} - 1)$ (B) $-2\sqrt{2}$ (C) $2\sqrt{2}$
(D) $2(\sqrt{2} - 1)$ (E) $2(\sqrt{2} + 1)$

99
98 29. $\int_{\pi/4}^{\pi/2} \frac{\cos x}{\sin x} dx =$

- (A) $\ln \sqrt{2}$ (B) $\ln \frac{\pi}{4}$ (C) $\ln \sqrt{3}$ (D) $\ln \frac{\sqrt{3}}{2}$ (E) $\ln e$

97 18. $\int_0^{\pi/4} \frac{e^{\tan x}}{\cos^2 x} dx$ is

- (A) 0 (B) 1 (C) $e-1$ (D) e (E) $e+1$

85 7. Which of the following is equal to $\ln 4$?

- (A) $\ln 3 + \ln 1$ (B) $\frac{\ln 8}{\ln 2}$ (C) $\int_1^4 e^t dt$ (D) $\int_1^4 \ln x dx$ (E) $\int_1^4 \frac{1}{t} dt$

85 17. $\int_0^1 x e^{-x} dx =$

- (A) $1-2e$ (B) -1 (C) $1-2e^{-1}$ (D) 1 (E) $2e-1$

88 38. For $x > 0$, $\int \left(\frac{1}{x} \int_1^x \frac{du}{u} \right) dx =$

- (A) $\frac{1}{x^3} + C$ (B) $\frac{8}{x^4} - \frac{2}{x^2} + C$ (C) $\ln(\ln x) + C$
 (D) $\frac{\ln(x^2)}{2} + C$ (E) $\frac{(\ln x)^2}{2} + C$

73 30. $\int_1^2 \frac{x-4}{x^2} dx =$

- (A) $-\frac{1}{2}$ (B) $\ln 2 - 2$ (C) $\ln 2$ (D) 2 (E) $\ln 2 + 2$

98
BC 7. $\int_1^e \left(\frac{x^2-1}{x} \right) dx =$

- (A) $e - \frac{1}{e}$ (B) $e^2 - e$ (C) $\frac{e^2}{2} - e + \frac{1}{2}$ (D) $e^2 - 2$ (E) $\frac{e^2}{2} - \frac{3}{2}$

85 BC 3. $\int_1^2 \frac{x+1}{x^2+2x} dx =$

- (A) $\ln 8 - \ln 3$ (B) $\frac{\ln 8 - \ln 3}{2}$ (C) $\ln 8$ (D) $\frac{3 \ln 2}{2}$ (E) $\frac{3 \ln 2 + 2}{2}$
-

93 BC 7. $\int_0^1 x^3 e^{x^4} dx =$

- (A) $\frac{1}{4}(e-1)$ (B) $\frac{1}{4}e$ (C) $e-1$ (D) e (E) $4(e-1)$
-

98 BC 25. $\int_0^{\infty} x^2 e^{-x^3} dx$ is

- (A) $-\frac{1}{3}$ (B) 0 (C) $\frac{1}{3}$ (D) 1 (E) divergent
-

98 7. $\int_1^e \left(\frac{x^2-1}{x} \right) dx =$

- (A) $e - \frac{1}{e}$ (B) $e^2 - e$ (C) $\frac{e^2}{2} - e + \frac{1}{2}$ (D) $e^2 - 2$ (E) $\frac{e^2}{2} - \frac{3}{2}$
-

73 21. $\int_0^1 (x+1)e^{x^2+2x} dx =$

- (A) $\frac{e^3}{2}$ (B) $\frac{e^3-1}{2}$ (C) $\frac{e^4-e}{2}$ (D) e^3-1 (E) e^4-e
-

85 9. If $\int_{-1}^1 e^{-x^2} dx = k$, then $\int_{-1}^0 e^{-x^2} dx =$

- (A) $-2k$ (B) $-k$ (C) $-\frac{k}{2}$ (D) $\frac{k}{2}$ (E) $2k$
-

85 24. If $\int_{-2}^2 (x^7 + k) dx = 16$, then $k =$

- (A) -12 (B) -4 (C) 0 (D) 4 (E) 12
-

- 98 20. What are all values of k for which $\int_{-3}^k x^2 dx = 0$?
- (A) -3 (B) 0 (C) 3 (D) -3 and 3 (E) $-3, 0,$ and 3

- 88 10. If $\int_0^k (2kx - x^2) dx = 18$, then $k =$
- (A) -9 (B) -3 (C) 3 (D) 9 (E) 18

- 88 39. If $\int_1^{10} f(x) dx = 4$ and $\int_{10}^3 f(x) dx = 7$, then $\int_1^3 f(x) dx =$
- (A) -3 (B) 0 (C) 3 (D) 10 (E) 11

- 85 27. $\int_0^3 |x-1| dx =$
- (A) 0 (B) $\frac{3}{2}$ (C) 2 (D) $\frac{5}{2}$ (E) 6

- 88 28. $\int_1^4 |x-3| dx =$
- (A) $-\frac{3}{2}$ (B) $\frac{3}{2}$ (C) $\frac{5}{2}$ (D) $\frac{9}{2}$ (E) 5

Definite Integrals - WITH CALCULATOR

93 28. $\int_1^{500} (13^x - 11^x) dx + \int_2^{500} (11^x - 13^x) dx =$

- (A) 0.000 (B) 14.946 (C) 34.415 (D) 46.000 (E) 136.364

97
92 89. If f is the antiderivative of $\frac{x^2}{1+x^5}$ such that $f(1) = 0$, then $f(4) =$

- (A) -0.012 (B) 0 (C) 0.016 (D) 0.376 (E) 0.629

98 88. Let $F(x)$ be an antiderivative of $\frac{(\ln x)^3}{x}$. If $F(1) = 0$, then $F(9) =$

- (A) 0.048 (B) 0.144 (C) 5.827 (D) 23.308 (E) 1,640.250

97 90. Which of the following are antiderivatives of $f(x) = \sin x \cos x$?

I. $F(x) = \frac{\sin^2 x}{2}$

II. $F(x) = \frac{\cos^2 x}{2}$

III. $F(x) = \frac{-\cos(2x)}{4}$

- (A) I only
(B) II only
(C) III only
(D) I and III only
(E) II and III only

1^{no} Fundamental Theorem of Calculus

85 42. $\frac{d}{dx} \int_2^x \sqrt{1+t^2} dt =$

(A) $\frac{x}{\sqrt{1+x^2}}$

(B) $\sqrt{1+x^2} - 5$

(C) $\sqrt{1+x^2}$

(D) $\frac{x}{\sqrt{1+x^2}} - \frac{1}{\sqrt{5}}$

(E) $\frac{1}{2\sqrt{1+x^2}} - \frac{1}{2\sqrt{5}}$

93 41. $\frac{d}{dx} \int_0^x \cos(2\pi u) du$ is

(A) 0

(B) $\frac{1}{2\pi} \sin x$

(C) $\frac{1}{2\pi} \cos(2\pi x)$

(D) $\cos(2\pi x)$

(E) $2\pi \cos(2\pi x)$

18 15. If $F(x) = \int_0^x \sqrt{t^3+1} dt$, then $F'(2) =$

(A) -3

(B) -2

(C) 2

(D) 3

(E) 18

88 25. For all $x > 1$, if $f(x) = \int_1^x \frac{1}{t} dt$, then $f'(x) =$

(A) 1

(B) $\frac{1}{x}$

(C) $\ln x - 1$

(D) $\ln x$

(E) e^x

69 12. If $F(x) = \int_0^x e^{-t^2} dt$, then $F'(x) =$

(A) $2xe^{-x^2}$

(B) $-2xe^{-x^2}$

(C) $\frac{e^{-x^2+1}}{-x^2+1} - e$

(D) $e^{-x^2} - 1$

(E) e^{-x^2}

88 14. If $F(x) = \int_1^{x^2} \sqrt{1+t^3} dt$, then $F'(x) =$


(A) $2x\sqrt{1+x^6}$

(B) $2x\sqrt{1+x^3}$






(C) $\sqrt{1+x^6}$

(D) $\sqrt{1+x^3}$

(E) $\int_1^{x^2} \frac{3t^2}{2\sqrt{1+t^3}} dt$




- 97 88. Let $f(x) = \int_a^x h(t) dt$, where h has the graph shown above. Which of the following could be the graph of f ?









- 97 88. Let $f(x) = \int_0^{x^2} \sin t dt$. At how many points in the closed interval $[0, \sqrt{\pi}]$ does the instantaneous rate of change of f equal the average rate of change of f on that interval?

- (A) Zero
 (B) One
 (C) Two
 (D) Three
 (E) Four



98
BC

88. Let $g(x) = \int_a^x f(t) dt$, where $a \leq x \leq b$. The figure above shows the graph of g on $[a, b]$. Which of the following could be the graph of f on $[a, b]$?




97
BC


22. The graph of f is shown in the figure above. If $g(x) = \int_a^x f(t) dt$, for what value of x does $g(x)$ have a maximum?

- (A) a
- (B) b
- (C) c
- (D) d
- (E) It cannot be determined from the information given.


Graphical Interpretation of 2FTC



- 98 2. The graph of a piecewise-linear function f , for $-1 \leq x \leq 4$, is shown above. What is the value of $\int_{-1}^4 f(x) dx$?
- (A) 1 (B) 2.5 (C) 4 (D) 5.5 (E) 8




- 93 2. Which of the following represents the area of the shaded region in the figure above?
- (A) $\int_c^d f(y) dy$ (B) $\int_a^b (d - f(x)) dx$ (C) $f'(b) - f'(a)$
- (D) $(b - a)[f(b) - f(a)]$ (E) $(d - c)[f(b) - f(a)]$



- 97 BC 19. The graph of $y = f(x)$ is shown in the figure above. If A_1 and A_2 are positive numbers that represent the areas of the shaded regions, then in terms of A_1 and A_2 ,


$$\int_{-4}^4 f(x) dx - 2 \int_{-1}^4 f(x) dx =$$

- (A) A_1 (B) $A_1 - A_2$ (C) $2A_1 - A_2$ (D) $A_1 + A_2$ (E) $A_1 + 2A_2$



- 97 78. The graph of f is shown in the figure above. If $\int_1^3 f(x) dx = 2.3$ and $F'(x) = f(x)$, then $F(3) - F(0) =$


- (A) 0.3 (B) 1.3 (C) 3.3 (D) 4.3 (E) 5.3



- 98 BC 9. The flow of oil, in barrels per hour, through a pipeline on July 9 is given by the graph shown above. Of the following, which best approximates the total number of barrels of oil that passed through the pipeline that day?

- (A) 500 (B) 600 (C) 2,400 (D) 3,000 (E) 4,800


Graph Analysis w/ Second FTC



Graph of f

11-A


4. The continuous function f is defined on the interval $-4 \leq x \leq 3$. The graph of f consists of two quarter circles and one line segment, as shown in the figure above. Let $g(x) = 2x + \int_0^x f(t) dt$.
- Find $g(-3)$. Find $g'(x)$ and evaluate $g'(-3)$.
 - Determine the x -coordinate of the point at which g has an absolute maximum on the interval $-4 \leq x \leq 3$. Justify your answer.
 - Find all values of x on the interval $-4 < x < 3$ for which the graph of g has a point of inflection. Give a reason for your answer.
 - Find the average rate of change of f on the interval $-4 \leq x \leq 3$. There is no point c , $-4 < c < 3$, for which $f'(c)$ is equal to that average rate of change. Explain why this statement does not contradict the Mean Value Theorem.



Graph of g


11-B

6. Let g be the piecewise-linear function defined on $[-2\pi, 4\pi]$ whose graph is given above, and let $f(x) = g(x) - \cos\left(\frac{x}{2}\right)$.
- Find $\int_{-2\pi}^{4\pi} f(x) dx$. Show the computations that lead to your answer.
 - Find all x -values in the open interval $(-2\pi, 4\pi)$ for which f has a critical point.
 - Let $h(x) = \int_0^{3x} g(t) dt$. Find $h'\left(-\frac{\pi}{3}\right)$.




6-A

3. The graph of the function f shown above consists of six line segments. Let g be the function given by $g(x) = \int_0^x f(t) dt$.
- Find $g(4)$, $g'(4)$, and $g''(4)$.
 - Does g have a relative minimum, a relative maximum, or neither at $x = 1$? Justify your answer.
 - Suppose that f is defined for all real numbers x and is periodic with a period of length 5. The graph above shows two periods of f . Given that $g(5) = 2$, find $g(10)$ and write an equation for the line tangent to the graph of g at $x = 108$.



5-B


4. The graph of the function f above consists of three line segments.
- Let g be the function given by $g(x) = \int_{-4}^x f(t) dt$. For each of $g(-1)$, $g'(-1)$, and $g''(-1)$, find the value or state that it does not exist.
 - For the function g defined in part (a), find the x -coordinate of each point of inflection of the graph of g on the open interval $-4 < x < 3$. Explain your reasoning.
 - Let h be the function given by $h(x) = \int_x^3 f(t) dt$. Find all values of x in the closed interval $-4 \leq x \leq 3$ for which $h(x) = 0$.
 - For the function h defined in part (c), find all intervals on which h is decreasing. Explain your reasoning.



4-A

5. The graph of the function f shown above consists of a semicircle and three line segments. Let g be the function given by $g(x) = \int_{-3}^x f(t) dt$.


- Find $g(0)$ and $g'(0)$.
- Find all values of x in the open interval $(-5, 4)$ at which g attains a relative maximum. Justify your answer.
- Find the absolute minimum value of g on the closed interval $[-5, 4]$. Justify your answer.
- Find all values of x in the open interval $(-5, 4)$ at which the graph of g has a point of inflection.



3-B

5. Let f be a function defined on the closed interval $[0, 7]$. The graph of f , consisting of four line segments, is shown above. Let g be the function given by $g(x) = \int_2^x f(t) dt$.

- Find $g(3)$, $g'(3)$, and $g''(3)$.
- Find the average rate of change of g on the interval $0 \leq x \leq 3$.
- For how many values c , where $0 < c < 3$, is $g'(c)$ equal to the average rate found in part (b)? Explain your reasoning.
- Find the x -coordinate of each point of inflection of the graph of g on the interval $0 < x < 7$. Justify your answer.




Graph of f

2-B

4. The graph of a differentiable function f on the closed interval $[-3, 15]$ is shown in the figure above. The graph of f has a horizontal tangent line at $x = 6$. Let $g(x) = 5 + \int_6^x f(t) dt$ for $-3 \leq x \leq 15$.

- Find $g(6)$, $g'(6)$, and $g''(6)$.
- On what intervals is g decreasing? Justify your answer.
- On what intervals is the graph of g concave down? Justify your answer.
- Find a trapezoidal approximation of $\int_{-3}^{15} f(t) dt$ using six subintervals of length $\Delta t = 3$.




Graph of f

2-A


4. The graph of the function f shown above consists of two line segments. Let g be the function given by $g(x) = \int_0^x f(t) dt$.

- Find $g(-1)$, $g'(-1)$, and $g''(-1)$.
- For what values of x in the open interval $(-2, 2)$ is g increasing? Explain your reasoning.
- For what values of x in the open interval $(-2, 2)$ is the graph of g concave down? Explain your reasoning.
- On the axes provided, sketch the graph of g on the closed interval $[-2, 2]$.


(Note: The axes are provided in the pink test booklet only.)



- 10-A 5. The function g is defined and differentiable on the closed interval $[-7, 5]$ and satisfies $g(0) = 5$. The graph of $y = g'(x)$, the derivative of g , consists of a semicircle and three line segments, as shown in the figure above.
- Find $g(3)$ and $g(-2)$.
 - Find the x -coordinate of each point of inflection of the graph of $y = g(x)$ on the interval $-7 < x < 5$. Explain your reasoning.
 - The function h is defined by $h(x) = g(x) - \frac{1}{2}x^2$. Find the x -coordinate of each critical point of h , where $-7 < x < 5$, and classify each critical point as the location of a relative minimum, relative maximum, or neither a minimum nor a maximum. Explain your reasoning.



- 9-B 3. A continuous function f is defined on the closed interval $-4 \leq x \leq 6$. The graph of f consists of a line segment and a curve that is tangent to the x -axis at $x = 3$, as shown in the figure above. On the interval $0 < x < 6$, the function f is twice differentiable, with $f''(x) > 0$.
- Is f differentiable at $x = 0$? Use the definition of the derivative with one-sided limits to justify your answer.
 - For how many values of a , $-4 \leq a < 6$, is the average rate of change of f on the interval $[a, 6]$ equal to 0? Give a reason for your answer.
 - Is there a value of a , $-4 \leq a < 6$, for which the Mean Value Theorem, applied to the interval $[a, 6]$, guarantees a value c , $a < c < 6$, at which $f'(c) = \frac{1}{3}$? Justify your answer.
 - The function g is defined by $g(x) = \int_0^x f(t) dt$ for $-4 \leq x \leq 6$. On what intervals contained in $[-4, 6]$ is the graph of g concave up? Explain your reasoning.



Graph of f

- 12-A 3. Let f be the continuous function defined on $[-4, 3]$ whose graph, consisting of three line segments and a semicircle centered at the origin, is given above. Let g be the function given by $g(x) = \int_1^x f(t) dt$.
- Find the values of $g(2)$ and $g(-2)$.
 - For each of $g'(-3)$ and $g''(-3)$, find the value or state that it does not exist.
 - Find the x -coordinate of each point at which the graph of g has a horizontal tangent line. For each of these points, determine whether g has a relative minimum, relative maximum, or neither a minimum nor a maximum at the point. Justify your answers.
 - For $-4 < x < 3$, find all values of x for which the graph of g has a point of inflection. Explain your reasoning.


**AP[®] CALCULUS AB
2011 SCORING GUIDELINES**

Question 4

The continuous function f is defined on the interval $-4 \leq x \leq 3$. The graph of f consists of two quarter circles and one line segment, as shown in the figure above.

Let $g(x) = 2x + \int_0^x f(t) dt$.

- (a) Find $g(-3)$. Find $g'(x)$ and evaluate $g'(-3)$.
- (b) Determine the x -coordinate of the point at which g has an absolute maximum on the interval $-4 \leq x \leq 3$. Justify your answer.
- (c) Find all values of x on the interval $-4 < x < 3$ for which the graph of g has a point of inflection. Give a reason for your answer.
- (d) Find the average rate of change of f on the interval $-4 \leq x \leq 3$. There is no point c , $-4 < c < 3$, for which $f'(c)$ is equal to that average rate of change. Explain why this statement does not contradict the Mean Value Theorem.



Graph of f

(a) $g(-3) = 2(-3) + \int_0^{-3} f(t) dt = -6 - \frac{9\pi}{4}$
 $g'(x) = 2 + f(x)$
 $g'(-3) = 2 + f(-3) = 2$

3 : $\begin{cases} 1 : g(-3) \\ 1 : g'(x) \\ 1 : g'(-3) \end{cases}$

(b) $g'(x) = 0$ when $f(x) = -2$. This occurs at $x = \frac{5}{2}$.
 $g'(x) > 0$ for $-4 < x < \frac{5}{2}$ and $g'(x) < 0$ for $\frac{5}{2} < x < 3$.

3 : $\begin{cases} 1 : \text{considers } g'(x) = 0 \\ 1 : \text{identifies interior candidate} \\ 1 : \text{answer with justification} \end{cases}$

Therefore g has an absolute maximum at $x = \frac{5}{2}$.

(c) $g''(x) = f'(x)$ changes sign only at $x = 0$. Thus the graph of g has a point of inflection at $x = 0$.

1 : answer with reason

(d) The average rate of change of f on the interval $-4 \leq x \leq 3$ is $\frac{f(3) - f(-4)}{3 - (-4)} = -\frac{2}{7}$.


2 : $\begin{cases} 1 : \text{average rate of change} \\ 1 : \text{explanation} \end{cases}$

To apply the Mean Value Theorem, f must be differentiable at each point in the interval $-4 < x < 3$. However, f is not differentiable at $x = -3$ and $x = 0$.

AP[®] CALCULUS AB
2011 SCORING GUIDELINES (Form B)

Question 6

Let g be the piecewise-linear function defined on $[-2\pi, 4\pi]$ whose graph is given above, and let $f(x) = g(x) - \cos\left(\frac{x}{2}\right)$.



Graph of g

- (a) Find $\int_{-2\pi}^{4\pi} f(x) dx$. Show the computations that lead to your answer.
- (b) Find all x -values in the open interval $(-2\pi, 4\pi)$ for which f has a critical point.
- (c) Let $h(x) = \int_0^{3x} g(t) dt$. Find $h'\left(-\frac{\pi}{3}\right)$.

$$\begin{aligned} \text{(a)} \quad \int_{-2\pi}^{4\pi} f(x) dx &= \int_{-2\pi}^{4\pi} \left(g(x) - \cos\left(\frac{x}{2}\right) \right) dx \\ &= 6\pi^2 - \left[2\sin\left(\frac{x}{2}\right) \right]_{x=-2\pi}^{x=4\pi} \\ &= 6\pi^2 \end{aligned}$$

2 : $\begin{cases} 1 : \text{antiderivative} \\ 1 : \text{answer} \end{cases}$

$$\text{(b)} \quad f'(x) = g'(x) + \frac{1}{2}\sin\left(\frac{x}{2}\right) = \begin{cases} 1 + \frac{1}{2}\sin\left(\frac{x}{2}\right) & \text{for } -2\pi < x < 0 \\ -\frac{1}{2} + \frac{1}{2}\sin\left(\frac{x}{2}\right) & \text{for } 0 < x < 4\pi \end{cases}$$

4 : $\begin{cases} 1 : \frac{d}{dx}\left(\cos\left(\frac{x}{2}\right)\right) \\ 1 : g'(x) \\ 1 : x = 0 \\ 1 : x = \pi \end{cases}$

$f'(x)$ does not exist at $x = 0$.

For $-2\pi < x < 0$, $f'(x) \neq 0$.

For $0 < x < 4\pi$, $f'(x) = 0$ when $x = \pi$.

f has critical points at $x = 0$ and $x = \pi$.

$$\begin{aligned} \text{(c)} \quad h'(x) &= g(3x) \cdot 3 \\ h'\left(-\frac{\pi}{3}\right) &= 3g(-\pi) = 3\pi \end{aligned}$$

3 : $\begin{cases} 2 : h'(x) \\ 1 : \text{answer} \end{cases}$


**AP[®] CALCULUS AB
2006 SCORING GUIDELINES**

Question 3

The graph of the function f shown above consists of six line segments. Let g be the function given by

$$g(x) = \int_0^x f(t) dt.$$

- (a) Find $g(4)$, $g'(4)$, and $g''(4)$.
 (b) Does g have a relative minimum, a relative maximum, or neither at $x = 1$? Justify your answer.



Graph of f

- (c) Suppose that f is defined for all real numbers x and is periodic with a period of length 5. The graph above shows two periods of f . Given that $g(5) = 2$, find $g(10)$ and write an equation for the line tangent to the graph of g at $x = 108$.

(a) $g(4) = \int_0^4 f(t) dt = 3$

$$g'(4) = f(4) = 0$$

$$g''(4) = f'(4) = -2$$

- (b) g has a relative minimum at $x = 1$ because $g' = f$ changes from negative to positive at $x = 1$.

- (c) $g(0) = 0$ and the function values of g increase by 2 for every increase of 5 in x .

$$g(10) = 2g(5) = 4$$

$$\begin{aligned} g(108) &= \int_0^{105} f(t) dt + \int_{105}^{108} f(t) dt \\ &= 21g(5) + g(3) = 44 \end{aligned}$$

$$g'(108) = f(108) = f(3) = 2$$

An equation for the line tangent to the graph of g at $x = 108$ is $y - 44 = 2(x - 108)$.

$$3 : \begin{cases} 1 : g(4) \\ 1 : g'(4) \\ 1 : g''(4) \end{cases}$$

$$2 : \begin{cases} 1 : \text{answer} \\ 1 : \text{reason} \end{cases}$$

$$4 : \begin{cases} 1 : g(10) \\ 3 : \begin{cases} 1 : g(108) \\ 1 : g'(108) \\ 1 : \text{equation of tangent line} \end{cases} \end{cases}$$


138

138

AP[®] CALCULUS AB
2005 SCORING GUIDELINES (Form B)

Question 4

The graph of the function f above consists of three line segments.



(a) Let g be the function given by $g(x) = \int_{-4}^x f(t) dt$.

For each of $g(-1)$, $g'(-1)$, and $g''(-1)$, find the value or state that it does not exist.

(b) For the function g defined in part (a), find the x -coordinate of each point of inflection of the graph of g on the open interval $-4 < x < 3$. Explain your reasoning.

(c) Let h be the function given by $h(x) = \int_x^3 f(t) dt$. Find all values of x in the closed interval $-4 \leq x \leq 3$ for which $h(x) = 0$.

(d) For the function h defined in part (c), find all intervals on which h is decreasing. Explain your reasoning.

(a) $g(-1) = \int_{-4}^{-1} f(t) dt = -\frac{1}{2}(3)(5) = -\frac{15}{2}$
 $g'(-1) = f(-1) = -2$
 $g''(-1)$ does not exist because f is not differentiable at $x = -1$.

3 : $\begin{cases} 1 : g(-1) \\ 1 : g'(-1) \\ 1 : g''(-1) \end{cases}$

(b) $x = 1$
 $g' = f$ changes from increasing to decreasing at $x = 1$.

2 : $\begin{cases} 1 : x = 1 \text{ (only)} \\ 1 : \text{reason} \end{cases}$

(c) $x = -1, 1, 3$

2 : correct values
 $\langle -1 \rangle$ each missing or extra value

(d) h is decreasing on $[0, 2]$
 $h' = -f < 0$ when $f > 0$

2 : $\begin{cases} 1 : \text{interval} \\ 1 : \text{reason} \end{cases}$


**AP[®] CALCULUS AB
2004 SCORING GUIDELINES**

Question 5

The graph of the function f shown above consists of a semicircle and three line segments. Let g be the function

given by $g(x) = \int_{-3}^x f(t) dt$.

- (a) Find $g(0)$ and $g'(0)$.
- (b) Find all values of x in the open interval $(-5, 4)$ at which g attains a relative maximum. Justify your answer.
- (c) Find the absolute minimum value of g on the closed interval $[-5, 4]$. Justify your answer.
- (d) Find all values of x in the open interval $(-5, 4)$ at which the graph of g has a point of inflection.



(a) $g(0) = \int_{-3}^0 f(t) dt = \frac{1}{2}(3)(2+1) = \frac{9}{2}$
 $g'(0) = f(0) = 1$

2: $\begin{cases} 1: g(0) \\ 1: g'(0) \end{cases}$

- (b) g has a relative maximum at $x = 3$.
 This is the only x -value where $g' = f$ changes from positive to negative.

2: $\begin{cases} 1: x = 3 \\ 1: \text{justification} \end{cases}$

- (c) The only x -value where f changes from negative to positive is $x = -4$. The other candidates for the location of the absolute minimum value are the endpoints.

3: $\begin{cases} 1: \text{identifies } x = -4 \text{ as a candidate} \\ 1: g(-4) = -1 \\ 1: \text{justification and answer} \end{cases}$

$g(-5) = 0$

$g(-4) = \int_{-3}^{-4} f(t) dt = -1$

$g(4) = \frac{9}{2} + \left(2 - \frac{\pi}{2}\right) = \frac{13 - \pi}{2}$

So the absolute minimum value of g is -1 .


- (d) $x = -3, 1, 2$

2: correct values
 $\langle -1 \rangle$ each missing or extra value

AP[®] CALCULUS AB
2003 SCORING GUIDELINES (Form B)

Question 5

Let f be a function defined on the closed interval $[0, 7]$. The graph of f , consisting of four line segments, is shown above. Let g be the function given by $g(x) = \int_2^x f(t) dt$.



Graph of f

- (a) Find $g(3)$, $g'(3)$, and $g''(3)$.
- (b) Find the average rate of change of g on the interval $0 \leq x \leq 3$.
- (c) For how many values c , where $0 < c < 3$, is $g'(c)$ equal to the average rate found in part (b)? Explain your reasoning.
- (d) Find the x -coordinate of each point of inflection of the graph of g on the interval $0 < x < 7$. Justify your answer.

(a) $g(3) = \int_2^3 f(t) dt = \frac{1}{2}(4 + 2) = 3$
 $g'(3) = f(3) = 2$
 $g''(3) = f'(3) = \frac{0 - 4}{4 - 2} = -2$

3 : $\left\{ \begin{array}{l} 1 : g(3) \\ 1 : g'(3) \\ 1 : g''(3) \end{array} \right.$

(b) $\frac{g(3) - g(0)}{3} = \frac{1}{3} \int_0^3 f(t) dt$
 $= \frac{1}{3} \left(\frac{1}{2}(2)(4) + \frac{1}{2}(4 + 2) \right) = \frac{7}{3}$

2 : $\left\{ \begin{array}{l} 1 : g(3) - g(0) = \int_0^3 f(t) dt \\ 1 : \text{answer} \end{array} \right.$

(c) There are two values of c .
 We need $\frac{7}{3} = g'(c) = f(c)$

2 : $\left\{ \begin{array}{l} 1 : \text{answer of 2} \\ 1 : \text{reason} \end{array} \right.$

The graph of f intersects the line $y = \frac{7}{3}$ at two places between 0 and 3.

Note: 1/2 if answer is 1 by MVT

(d) $x = 2$ and $x = 5$
 because $g' = f$ changes from increasing to decreasing at $x = 2$, and from decreasing to increasing at $x = 5$.

2 : $\left\{ \begin{array}{l} 1 : x = 2 \text{ and } x = 5 \text{ only} \\ 1 : \text{justification} \\ \quad (\text{ignore discussion at } x = 4) \end{array} \right.$


141

141

AP[®] CALCULUS AB
2002 SCORING GUIDELINES (Form B)

Question 4

The graph of a differentiable function f on the closed interval $[-3, 15]$ is shown in the figure above. The graph of f has a horizontal tangent line at $x = 6$. Let



$$g(x) = 5 + \int_6^x f(t) dt \text{ for } -3 \leq x \leq 15.$$

- (a) Find $g(6)$, $g'(6)$, and $g''(6)$.
 (b) On what intervals is g decreasing? Justify your answer.
 (c) On what intervals is the graph of g concave down? Justify your answer.
 (d) Find a trapezoidal approximation of $\int_{-3}^{15} f(t) dt$ using six subintervals of length $\Delta t = 3$.

(a) $g(6) = 5 + \int_6^6 f(t) dt = 5$

$$g'(6) = f(6) = 3$$

$$g''(6) = f'(6) = 0$$

$$3 \begin{cases} 1 : g(6) \\ 1 : g'(6) \\ 1 : g''(6) \end{cases}$$

(b) g is decreasing on $[-3, 0]$ and $[12, 15]$ since

$$g'(x) = f(x) < 0 \text{ for } x < 0 \text{ and } x > 12.$$

$$3 \begin{cases} 1 : [-3, 0] \\ 1 : [12, 15] \\ 1 : \text{justification} \end{cases}$$

(c) The graph of g is concave down on $(6, 15)$ since

$$g' = f \text{ is decreasing on this interval.}$$

$$2 \begin{cases} 1 : \text{interval} \\ 1 : \text{justification} \end{cases}$$


(d) $\frac{3}{2}(-1 + 2(0 + 1 + 3 + 1 + 0) - 1)$
 $= 12$

1 : trapezoidal method

AP[®] CALCULUS AB 2002 SCORING GUIDELINES

Question 4

The graph of the function f shown above consists of two line segments. Let g be the function given by $g(x) = \int_0^x f(t) dt$.



- (a) Find $g(-1)$, $g'(-1)$, and $g''(-1)$.
- (b) For what values of x in the open interval $(-2, 2)$ is g increasing? Explain your reasoning.
- (c) For what values of x in the open interval $(-2, 2)$ is the graph of g concave down? Explain your reasoning.
- (d) On the axes provided, sketch the graph of g on the closed interval $[-2, 2]$.

(a) $g(-1) = \int_0^{-1} f(t) dt = -\int_{-1}^0 f(t) dt = -\frac{3}{2}$
 $g'(-1) = f(-1) = 0$
 $g''(-1) = f'(-1) = 3$

3 $\left\{ \begin{array}{l} 1: g(-1) \\ 1: g'(-1) \\ 1: g''(-1) \end{array} \right.$


(b) g is increasing on $-1 < x < 1$ because $g'(x) = f(x) > 0$ on this interval.

2 $\left\{ \begin{array}{l} 1: \text{interval} \\ 1: \text{reason} \end{array} \right.$

(c) The graph of g is concave down on $0 < x < 2$ because $g''(x) = f'(x) < 0$ on this interval.
 or
 because $g'(x) = f(x)$ is decreasing on this interval.

2 $\left\{ \begin{array}{l} 1: \text{interval} \\ 1: \text{reason} \end{array} \right.$


(d)



2 $\left\{ \begin{array}{l} 1: g(-2) = g(0) = g(2) = 0 \\ 1: \text{appropriate increasing/decreasing} \\ \text{and concavity behavior} \\ < -1 > \text{vertical asymptote} \end{array} \right.$

AP[®] CALCULUS AB
2010 SCORING GUIDELINES

Question 5



The function g is defined and differentiable on the closed interval $[-7, 5]$ and satisfies $g(0) = 5$. The graph of $y = g'(x)$, the derivative of g , consists of a semicircle and three line segments, as shown in the figure above.

- (a) Find $g(3)$ and $g(-2)$.
- (b) Find the x -coordinate of each point of inflection of the graph of $y = g(x)$ on the interval $-7 < x < 5$. Explain your reasoning.
- (c) The function h is defined by $h(x) = g(x) - \frac{1}{2}x^2$. Find the x -coordinate of each critical point of h , where $-7 < x < 5$, and classify each critical point as the location of a relative minimum, relative maximum, or neither a minimum nor a maximum. Explain your reasoning.

(a) $g(3) = 5 + \int_0^3 g'(x) dx = 5 + \frac{\pi \cdot 2^2}{4} + \frac{3}{2} = \frac{13}{2} + \pi$
 $g(-2) = 5 + \int_0^{-2} g'(x) dx = 5 - \pi$

3 : $\begin{cases} 1 : \text{uses } g(0) = 5 \\ 1 : g(3) \\ 1 : g(-2) \end{cases}$

- (b) The graph of $y = g(x)$ has points of inflection at $x = 0$, $x = 2$, and $x = 3$ because g' changes from increasing to decreasing at $x = 0$ and $x = 3$, and g' changes from decreasing to increasing at $x = 2$.

2 : $\begin{cases} 1 : \text{identifies } x = 0, 2, 3 \\ 1 : \text{explanation} \end{cases}$


- (c) $h'(x) = g'(x) - x = 0 \Rightarrow g'(x) = x$
 On the interval $-2 \leq x \leq 2$, $g'(x) = \sqrt{4 - x^2}$.
 On this interval, $g'(x) = x$ when $x = \sqrt{2}$.
 The only other solution to $g'(x) = x$ is $x = 3$.
 $h'(x) = g'(x) - x > 0$ for $0 \leq x < \sqrt{2}$
 $h'(x) = g'(x) - x \leq 0$ for $\sqrt{2} < x \leq 5$

4 : $\begin{cases} 1 : h'(x) \\ 1 : \text{identifies } x = \sqrt{2}, 3 \\ 1 : \text{answer for } \sqrt{2} \text{ with analysis} \\ 1 : \text{answer for } 3 \text{ with analysis} \end{cases}$

Therefore h has a relative maximum at $x = \sqrt{2}$, and h has neither a minimum nor a maximum at $x = 3$.

AP[®] CALCULUS AB
2009 SCORING GUIDELINES (Form B)

Question 3



Graph of f

A continuous function f is defined on the closed interval $-4 \leq x \leq 6$. The graph of f consists of a line segment and a curve that is tangent to the x -axis at $x = 3$, as shown in the figure above. On the interval $0 < x < 6$, the function f is twice differentiable, with $f''(x) > 0$.

- (a) Is f differentiable at $x = 0$? Use the definition of the derivative with one-sided limits to justify your answer.
- (b) For how many values of a , $-4 \leq a < 6$, is the average rate of change of f on the interval $[a, 6]$ equal to 0? Give a reason for your answer.
- (c) Is there a value of a , $-4 \leq a < 6$, for which the Mean Value Theorem, applied to the interval $[a, 6]$, guarantees a value c , $a < c < 6$, at which $f'(c) = \frac{1}{3}$? Justify your answer.
- (d) The function g is defined by $g(x) = \int_0^x f(t) dt$ for $-4 \leq x \leq 6$. On what intervals contained in $[-4, 6]$ is the graph of g concave up? Explain your reasoning.

(a)
$$\lim_{h \rightarrow 0^-} \frac{f(h) - f(0)}{h} = \frac{2}{3}$$

$$\lim_{h \rightarrow 0^+} \frac{f(h) - f(0)}{h} < 0$$

Since the one-sided limits do not agree, f is not differentiable at $x = 0$.

2 : $\left\{ \begin{array}{l} 1 : \text{sets up difference quotient at } x = 0 \\ 1 : \text{answer with justification} \end{array} \right.$

(b)
$$\frac{f(6) - f(a)}{6 - a} = 0 \text{ when } f(a) = f(6). \text{ There are two values of } a \text{ for which this is true.}$$

(c) Yes, $a = 3$. The function f is differentiable on the interval $3 < x < 6$ and continuous on $3 \leq x \leq 6$.

Also,
$$\frac{f(6) - f(3)}{6 - 3} = \frac{1 - 0}{6 - 3} = \frac{1}{3}.$$

By the Mean Value Theorem, there is a value c ,

$$3 < c < 6, \text{ such that } f'(c) = \frac{1}{3}.$$

(d)
$$g'(x) = f(x), \quad g''(x) = f'(x)$$

$$g''(x) > 0 \text{ when } f'(x) > 0$$

This is true for $-4 < x < 0$ and $3 < x < 6$.

2 : $\left\{ \begin{array}{l} 1 : \text{expression for average rate of change} \\ 1 : \text{answer with reason} \end{array} \right.$

2 : $\left\{ \begin{array}{l} 1 : \text{answers "yes" and identifies } a = 3 \\ 1 : \text{justification} \end{array} \right.$

3 : $\left\{ \begin{array}{l} 1 : g'(x) = f(x) \\ 1 : \text{considers } g''(x) > 0 \\ 1 : \text{answer} \end{array} \right.$

Integrals - Piecewise Functions & Inverse Trig

69 41. If $\begin{cases} f(x) = 8 - x^2 & \text{for } -2 \leq x \leq 2, \\ f(x) = x^2 & \text{elsewhere,} \end{cases}$ then $\int_{-1}^3 f(x) dx$ is a number between

- (A) 0 and 8 (B) 8 and 16 (C) 16 and 24 (D) 24 and 32 (E) 32 and 40

93 37. If $f(x) = \begin{cases} x & \text{for } x \leq 1 \\ \frac{1}{x} & \text{for } x > 1, \end{cases}$ then $\int_0^e f(x) dx =$

- (A) 0 (B) $\frac{3}{2}$ (C) 2 (D) e (E) $e + \frac{1}{2}$

73 41. Given $f(x) = \begin{cases} x+1 & \text{for } x < 0, \\ \cos \pi x & \text{for } x \geq 0, \end{cases}$ $\int_{-1}^1 f(x) dx =$

- (A) $\frac{1}{2} + \frac{1}{\pi}$ (B) $-\frac{1}{2}$ (C) $\frac{1}{2} - \frac{1}{\pi}$ (D) $\frac{1}{2}$ (E) $-\frac{1}{2} + \pi$

93 32. $\int_0^{\sqrt{3}} \frac{dx}{\sqrt{4-x^2}} =$

- (A) $\frac{\pi}{3}$ (B) $\frac{\pi}{4}$ (C) $\frac{\pi}{6}$ (D) $\frac{1}{2} \ln 2$ (E) $-\ln 2$

85 7. Which of the following is equal to $\int \frac{1}{\sqrt{25-x^2}} dx$?

- (A) $\arcsin \frac{x}{5} + C$ (B) $\arcsin x + C$ (C) $\frac{1}{5} \arcsin \frac{x}{5} + C$
 (D) $\sqrt{25-x^2} + C$ (E) $2\sqrt{25-x^2} + C$

73 32. $\int \frac{5}{1+x^2} dx =$

- (A) $\frac{-10x}{(1+x^2)^2} + C$ (B) $\frac{5}{2x} \ln(1+x^2) + C$ (C) $5x - \frac{5}{x} + C$
 (D) $5 \arctan x + C$ (E) $5 \ln(1+x^2) + C$

73
BC

43. $\int \arcsin x \, dx =$

(A) $\sin x - \int \frac{x \, dx}{\sqrt{1-x^2}}$

(B) $\frac{(\arcsin x)^2}{2} + C$

(C) $\arcsin x + \int \frac{dx}{\sqrt{1-x^2}}$

(D) $x \arccos x - \int \frac{x \, dx}{\sqrt{1-x^2}}$

(E) $x \arcsin x - \int \frac{x \, dx}{\sqrt{1-x^2}}$

Miscellaneous Integrals

- 88
x
42. If $\int_1^4 f(x) dx = 6$, what is the value of $\int_1^4 f(5-x) dx$?
- (A) 6 (B) 3 (C) 0 (D) -1 (E) -6

- 73
38. If $\int_1^2 f(x-c) dx = 5$ where c is a constant, then $\int_{1-c}^{2-c} f(x) dx =$
- (A) $5+c$ (B) 5 (C) $5-c$ (D) $c-5$ (E) -5

- 97
3. If $\int_a^b f(x) dx = a+2b$, then $\int_a^b (f(x)+5) dx =$
- (A) $a+2b+5$ (B) $5b-5a$ (C) $7b-4a$ (D) $7b-5a$ (E) $7b-6a$

- 85
40. Let f be a continuous function on the closed interval $[0,2]$. If $2 \leq f(x) \leq 4$, then the greatest possible value of $\int_0^2 f(x) dx$ is
- (A) 0 (B) 2 (C) 4 (D) 8 (E) 16

- 85
38. Let f and g have continuous first and second derivatives everywhere. If $f(x) \leq g(x)$ for all real x , which of the following must be true?

- I. $f'(x) \leq g'(x)$ for all real x
II. $f''(x) \leq g''(x)$ for all real x
III. $\int_0^1 f(x) dx \leq \int_0^1 g(x) dx$

- (A) None (B) I only (C) III only (D) I and II only (E) I, II, and III

- 18
11. If f is a linear function and $0 < a < b$, then $\int_a^b f''(x) dx =$

- (A) 0 (B) 1 (C) $\frac{ab}{2}$ (D) $b-a$ (E) $\frac{b^2-a^2}{2}$

93 43. $\int x f(x) dx =$

(A) $x f(x) - \int x f'(x) dx$

(B) $\frac{x^2}{2} f(x) - \int \frac{x^2}{2} f'(x) dx$

(C) $x f(x) - \frac{x^2}{2} f(x) + C$

(D) $x f(x) - \int f'(x) dx$

(E) $\frac{x^2}{2} \int f(x) dx$

93 12. If f and g are continuous functions, and if $f(x) \geq 0$ for all real numbers x , which of the following must be true?

I. $\int_a^b f(x)g(x)dx = \left(\int_a^b f(x)dx\right)\left(\int_a^b g(x)dx\right)$

II. $\int_a^b (f(x) + g(x))dx = \int_a^b f(x)dx + \int_a^b g(x)dx$

III. $\int_a^b \sqrt{f(x)} dx = \sqrt{\int_a^b f(x)dx}$

(A) I only

(B) II only

(C) III only

(D) II and III only

(E) I, II, and III

98 82. If f is a continuous function and if $F'(x) = f(x)$ for all real numbers x , then $\int_1^3 f(2x) dx =$

(A) $2F(3) - 2F(1)$

(B) $\frac{1}{2}F(3) - \frac{1}{2}F(1)$

(C) $2F(6) - 2F(2)$

(D) $F(6) - F(2)$

(E) $\frac{1}{2}F(6) - \frac{1}{2}F(2)$

Average Value

88 BC 21. The average value of $\frac{1}{x}$ on the closed interval $[1, 3]$ is

- (A) $\frac{1}{2}$ (B) $\frac{2}{3}$ (C) $\frac{\ln 2}{2}$ (D) $\frac{\ln 3}{2}$ (E) $\ln 3$
-

88 36. What is the average value of y for the part of the curve $y = 3x - x^2$ which is in the first quadrant?

- (A) -6 (B) -2 (C) $\frac{3}{2}$ (D) $\frac{9}{4}$ (E) $\frac{9}{2}$
-

97 BC 85. Let f be a twice differentiable function such that $f(1) = 2$ and $f(3) = 7$. Which of the following must be true for the function f on the interval $1 \leq x \leq 3$?

- I. The average rate of change of f is $\frac{5}{2}$.
II. The average value of f is $\frac{9}{2}$.
III. The average value of f' is $\frac{5}{2}$.
- (A) None
(B) I only
(C) III only
(D) I and III only
(E) II and III only
-

88 BC 24. If c is the number that satisfies the conclusion of the Mean Value Theorem for $f(x) = x^3 - 2x^2$ on the interval $0 \leq x \leq 2$, then $c =$

- (A) 0 (B) $\frac{1}{2}$ (C) 1 (D) $\frac{4}{3}$ (E) 2
-

73 34. The average value of \sqrt{x} over the interval $0 \leq x \leq 2$ is

- (A) $\frac{1}{3}\sqrt{2}$ (B) $\frac{1}{2}\sqrt{2}$ (C) $\frac{2}{3}\sqrt{2}$ (D) 1 (E) $\frac{4}{3}\sqrt{2}$
-

85 44. The average value of $f(x) = x^2\sqrt{x^3+1}$ on the closed interval $[0,2]$ is

- (A) $\frac{26}{9}$ (B) $\frac{13}{3}$ (C) $\frac{26}{3}$ (D) 13 (E) 26
-

97 20. The average value of $\cos x$ on the interval $[-3,5]$ is

- (A) $\frac{\sin 5 - \sin 3}{8}$
(B) $\frac{\sin 5 - \sin 3}{2}$
(C) $\frac{\sin 3 - \sin 5}{2}$
(D) $\frac{\sin 3 + \sin 5}{2}$
(E) $\frac{\sin 3 + \sin 5}{8}$
-

69 33. What is the average (mean) value of $3t^3 - t^2$ over the interval $-1 \leq t \leq 2$?

- A
(A) $\frac{11}{4}$ (B) $\frac{7}{2}$ (C) 8 (D) $\frac{33}{4}$ (E) 16
-

Straightforward

2-A

2. The rate at which people enter an amusement park on a given day is modeled by the function E defined by

$$E(t) = \frac{15600}{(t^2 - 24t + 160)}$$

The rate at which people leave the same amusement park on the same day is modeled by the function L defined by

$$L(t) = \frac{9890}{(t^2 - 38t + 370)}$$

Both $E(t)$ and $L(t)$ are measured in people per hour and time t is measured in hours after midnight. These functions are valid for $9 \leq t \leq 23$, the hours during which the park is open. At time $t = 9$, there are no people in the park.

- (a) How many people have entered the park by 5:00 P.M. ($t = 17$)? Round your answer to the nearest whole number.
- (b) The price of admission to the park is \$15 until 5:00 P.M. ($t = 17$). After 5:00 P.M., the price of admission to the park is \$11. How many dollars are collected from admissions to the park on the given day? Round your answer to the nearest whole number.
- (c) Let $H(t) = \int_9^t (E(x) - L(x)) dx$ for $9 \leq t \leq 23$. The value of $H(17)$ to the nearest whole number is 3725. Find the value of $H'(17)$, and explain the meaning of $H(17)$ and $H'(17)$ in the context of the amusement park.
- (d) At what time t , for $9 \leq t \leq 23$, does the model predict that the number of people in the park is a maximum?

2-B

2. The number of gallons, $P(t)$, of a pollutant in a lake changes at the rate $P'(t) = 1 - 3e^{-0.2\sqrt{t}}$ gallons per day, where t is measured in days. There are 50 gallons of the pollutant in the lake at time $t = 0$. The lake is considered to be safe when it contains 40 gallons or less of pollutant.

- (a) Is the amount of pollutant increasing at time $t = 9$? Why or why not?
- (b) For what value of t will the number of gallons of pollutant be at its minimum? Justify your answer.
- (c) Is the lake safe when the number of gallons of pollutant is at its minimum? Justify your answer.
- (d) An investigator uses the tangent line approximation to $P(t)$ at $t = 0$ as a model for the amount of pollutant in the lake. At what time t does this model predict that the lake becomes safe?

3-8

2. A tank contains 125 gallons of heating oil at time $t = 0$. During the time interval $0 \leq t \leq 12$ hours, heating oil is pumped into the tank at the rate

$$H(t) = 2 + \frac{10}{(1 + \ln(t + 1))} \text{ gallons per hour.}$$

During the same time interval, heating oil is removed from the tank at the rate

$$R(t) = 12 \sin\left(\frac{t^2}{47}\right) \text{ gallons per hour.}$$

- How many gallons of heating oil are pumped into the tank during the time interval $0 \leq t \leq 12$ hours?
- Is the level of heating oil in the tank rising or falling at time $t = 6$ hours? Give a reason for your answer.
- How many gallons of heating oil are in the tank at time $t = 12$ hours?
- At what time t , for $0 \leq t \leq 12$, is the volume of heating oil in the tank the least? Show the analysis that leads to your conclusion.

5-3

2. A water tank at Camp Newton holds 1200 gallons of water at time $t = 0$. During the time interval $0 \leq t \leq 18$ hours, water is pumped into the tank at the rate

$$W(t) = 95\sqrt{t} \sin^2\left(\frac{t}{6}\right) \text{ gallons per hour.}$$

During the same time interval, water is removed from the tank at the rate

$$R(t) = 275 \sin^2\left(\frac{t}{3}\right) \text{ gallons per hour.}$$

- Is the amount of water in the tank increasing at time $t = 15$? Why or why not?
- To the nearest whole number, how many gallons of water are in the tank at time $t = 18$?
- At what time t , for $0 \leq t \leq 18$, is the amount of water in the tank at an absolute minimum? Show the work that leads to your conclusion.
- For $t > 18$, no water is pumped into the tank, but water continues to be removed at the rate $R(t)$ until the tank becomes empty. Let k be the time at which the tank becomes empty. Write, but do not solve, an equation involving an integral expression that can be used to find the value of k .

5-A

2. The tide removes sand from Sandy Point Beach at a rate modeled by the function R , given by

$$R(t) = 2 + 5 \sin\left(\frac{4\pi t}{25}\right).$$

A pumping station adds sand to the beach at a rate modeled by the function S , given by

$$S(t) = \frac{15t}{1 + 3t}.$$

Both $R(t)$ and $S(t)$ have units of cubic yards per hour and t is measured in hours for $0 \leq t \leq 6$. At time $t = 0$, the beach contains 2500 cubic yards of sand.

- How much sand will the tide remove from the beach during this 6-hour period? Indicate units of measure.
- Write an expression for $Y(t)$, the total number of cubic yards of sand on the beach at time t .
- Find the rate at which the total amount of sand on the beach is changing at time $t = 4$.
- For $0 \leq t \leq 6$, at what time t is the amount of sand on the beach a minimum? What is the minimum value? Justify your answers.

4-A

1. Traffic flow is defined as the rate at which cars pass through an intersection, measured in cars per minute. The traffic flow at a particular intersection is modeled by the function F defined by


$$F(t) = 82 + 4 \sin\left(\frac{t}{2}\right) \text{ for } 0 \leq t \leq 30,$$

where $F(t)$ is measured in cars per minute and t is measured in minutes.

- To the nearest whole number, how many cars pass through the intersection over the 30-minute period?
- Is the traffic flow increasing or decreasing at $t = 7$? Give a reason for your answer.
- What is the average value of the traffic flow over the time interval $10 \leq t \leq 15$? Indicate units of measure.
- What is the average rate of change of the traffic flow over the time interval $10 \leq t \leq 15$? Indicate units of measure.

9-A

2. The rate at which people enter an auditorium for a rock concert is modeled by the function R given by $R(t) = 1380t^2 - 675t^3$ for $0 \leq t \leq 2$ hours; $R(t)$ is measured in people per hour. No one is in the auditorium at time $t = 0$, when the doors open. The doors close and the concert begins at time $t = 2$.
- How many people are in the auditorium when the concert begins?
 - Find the time when the rate at which people enter the auditorium is a maximum. Justify your answer.
 - The total wait time for all the people in the auditorium is found by adding the time each person waits, starting at the time the person enters the auditorium and ending when the concert begins. The function w models the total wait time for all the people who enter the auditorium before time t . The derivative of w is given by $w'(t) = (2 - t)R(t)$. Find $w(2) - w(1)$, the total wait time for those who enter the auditorium after time $t = 1$.
 - On average, how long does a person wait in the auditorium for the concert to begin? Consider all people who enter the auditorium after the doors open, and use the model for total wait time from part (c).



6-A

2. At an intersection in Thomasville, Oregon, cars turn left at the rate $L(t) = 60\sqrt{t} \sin^2\left(\frac{t}{3}\right)$ cars per hour over the time interval $0 \leq t \leq 18$ hours. The graph of $y = L(t)$ is shown above.
- To the nearest whole number, find the total number of cars turning left at the intersection over the time interval $0 \leq t \leq 18$ hours.
 - Traffic engineers will consider turn restrictions when $L(t) \geq 150$ cars per hour. Find all values of t for which $L(t) \geq 150$ and compute the average value of L over this time interval. Indicate units of measure.
 - Traffic engineers will install a signal if there is any two-hour time interval during which the product of the total number of cars turning left and the total number of oncoming cars traveling straight through the intersection is greater than 200,000. In every two-hour time interval, 500 oncoming cars travel straight through the intersection. Does this intersection require a traffic signal? Explain the reasoning that leads to your conclusion.

9-5

1. At a certain height, a tree trunk has a circular cross section. The radius $R(t)$ of that cross section grows at a rate modeled by the function

$$\frac{dR}{dt} = \frac{1}{16}(3 + \sin(t^2)) \text{ centimeters per year}$$

for $0 \leq t \leq 3$, where time t is measured in years. At time $t = 0$, the radius is 6 centimeters. The area of the cross section at time t is denoted by $A(t)$.

- Write an expression, involving an integral, for the radius $R(t)$ for $0 \leq t \leq 3$. Use your expression to find $R(3)$.
- Find the rate at which the cross-sectional area $A(t)$ is increasing at time $t = 3$ years. Indicate units of measure.
- Evaluate $\int_0^3 A'(t) dt$. Using appropriate units, interpret the meaning of that integral in terms of cross-sectional area.

Critical Thinking

- 118 1. A cylindrical can of radius 10 millimeters is used to measure rainfall in Stormville. The can is initially empty, and rain enters the can during a 60-day period. The height of water in the can is modeled by the function S , where $S(t)$ is measured in millimeters and t is measured in days for $0 \leq t \leq 60$. The rate at which the height of the water is rising in the can is given by $S'(t) = 2 \sin(0.03t) + 1.5$.
- According to the model, what is the height of the water in the can at the end of the 60-day period?
 - According to the model, what is the average rate of change in the height of water in the can over the 60-day period? Show the computations that lead to your answer. Indicate units of measure.
-
- Assuming no evaporation occurs, at what rate is the volume of water in the can changing at time $t = 7$? Indicate units of measure.
 - During the same 60-day period, rain on Monsoon Mountain accumulates in a can identical to the one in Stormville. The height of the water in the can on Monsoon Mountain is modeled by the function M , where $M(t) = \frac{1}{400}(3t^3 - 30t^2 + 330t)$. The height $M(t)$ is measured in millimeters, and t is measured in days for $0 \leq t \leq 60$. Let $D(t) = M'(t) - S'(t)$. Apply the Intermediate Value Theorem to the function D on the interval $0 \leq t \leq 60$ to justify that there exists a time t , $0 < t < 60$, at which the heights of water in the two cans are changing at the same rate.

7-8

3. The wind chill is the temperature, in degrees Fahrenheit ($^{\circ}\text{F}$), a human feels based on the air temperature, in degrees Fahrenheit, and the wind velocity v , in miles per hour (mph). If the air temperature is 32°F , then the wind chill is given by $W(v) = 55.6 - 22.1v^{0.16}$ and is valid for $5 \leq v \leq 60$.
- Find $W'(20)$. Using correct units, explain the meaning of $W'(20)$ in terms of the wind chill.
 - Find the average rate of change of W over the interval $5 \leq v \leq 60$. Find the value of v at which the instantaneous rate of change of W is equal to the average rate of change of W over the interval $5 \leq v \leq 60$.
 - Over the time interval $0 \leq t \leq 4$ hours, the air temperature is a constant 32°F . At time $t = 0$, the wind velocity is $v = 20$ mph. If the wind velocity increases at a constant rate of 5 mph per hour, what is the rate of change of the wind chill with respect to time at $t = 3$ hours? Indicate units of measure.

9-8

2. A storm washed away sand from a beach, causing the edge of the water to get closer to a nearby road. The rate at which the distance between the road and the edge of the water was changing during the storm is modeled by $f(t) = \sqrt{t} + \cos t - 3$ meters per hour, t hours after the storm began. The edge of the water was 35 meters from the road when the storm began, and the storm lasted 5 hours. The derivative of $f(t)$ is $f'(t) = \frac{1}{2\sqrt{t}} - \sin t$.
- What was the distance between the road and the edge of the water at the end of the storm?
 - Using correct units, interpret the value $f'(4) = 1.007$ in terms of the distance between the road and the edge of the water.
 - At what time during the 5 hours of the storm was the distance between the road and the edge of the water decreasing most rapidly? Justify your answer.
 - After the storm, a machine pumped sand back onto the beach so that the distance between the road and the edge of the water was growing at a rate of $g(p)$ meters per day, where p is the number of days since pumping began. Write an equation involving an integral expression whose solution would give the number of days that sand must be pumped to restore the original distance between the road and the edge of the water.

8-8

2. For time $t \geq 0$ hours, let $r(t) = 120(1 - e^{-10t^2})$ represent the speed, in kilometers per hour, at which a car travels along a straight road. The number of liters of gasoline used by the car to travel x kilometers is modeled by $g(x) = 0.05x(1 - e^{-x/2})$.

- How many kilometers does the car travel during the first 2 hours?
- Find the rate of change with respect to time of the number of liters of gasoline used by the car when $t = 2$ hours. Indicate units of measure.
- How many liters of gasoline have been used by the car when it reaches a speed of 80 kilometers per hour?

10A

1. There is no snow on Janet's driveway when snow begins to fall at midnight. From midnight to 9 A.M., snow accumulates on the driveway at a rate modeled by $f(t) = 7te^{\cos t}$ cubic feet per hour, where t is measured in hours since midnight. Janet starts removing snow at 6 A.M. ($t = 6$). The rate $g(t)$, in cubic feet per hour, at which Janet removes snow from the driveway at time t hours after midnight is modeled by


$$g(t) = \begin{cases} 0 & \text{for } 0 \leq t < 6 \\ 125 & \text{for } 6 \leq t < 7 \\ 108 & \text{for } 7 \leq t \leq 9. \end{cases}$$

- How many cubic feet of snow have accumulated on the driveway by 6 A.M.?
 - Find the rate of change of the volume of snow on the driveway at 8 A.M.
-
- Let $h(t)$ represent the total amount of snow, in cubic feet, that Janet has removed from the driveway at time t hours after midnight. Express h as a piecewise-defined function with domain $0 \leq t \leq 9$.
 - How many cubic feet of snow are on the driveway at 9 A.M.?

4-8

2. For $0 \leq t \leq 31$, the rate of change of the number of mosquitoes on Tropical Island at time t days is modeled by $R(t) = 5\sqrt{t} \cos\left(\frac{t}{5}\right)$ mosquitoes per day. There are 1000 mosquitoes on Tropical Island at time $t = 0$.

- (a) Show that the number of mosquitoes is increasing at time $t = 6$.
- (b) At time $t = 6$, is the number of mosquitoes increasing at an increasing rate, or is the number of mosquitoes increasing at a decreasing rate? Give a reason for your answer.
- (c) According to the model, how many mosquitoes will be on the island at time $t = 31$? Round your answer to the nearest whole number.
- (d) To the nearest whole number, what is the maximum number of mosquitoes for $0 \leq t \leq 31$? Show the analysis that leads to your conclusion.



7-A

2. The amount of water in a storage tank, in gallons, is modeled by a continuous function on the time interval $0 \leq t \leq 7$, where t is measured in hours. In this model, rates are given as follows:


- (i) The rate at which water enters the tank is $f(t) = 100t^2 \sin(\sqrt{t})$ gallons per hour for $0 \leq t \leq 7$.
- (ii) The rate at which water leaves the tank is

$$g(t) = \begin{cases} 250 & \text{for } 0 \leq t < 3 \\ 2000 & \text{for } 3 < t \leq 7 \end{cases} \text{ gallons per hour.}$$

The graphs of f and g , which intersect at $t = 1.617$ and $t = 5.076$, are shown in the figure above. At time $t = 0$, the amount of water in the tank is 5000 gallons.


- (a) How many gallons of water enter the tank during the time interval $0 \leq t \leq 7$? Round your answer to the nearest gallon.
- (b) For $0 \leq t \leq 7$, find the time intervals during which the amount of water in the tank is decreasing. Give a reason for each answer.
- (c) For $0 \leq t \leq 7$, at what time t is the amount of water in the tank greatest? To the nearest gallon, compute the amount of water at this time. Justify your answer.

Accumulation → Non-Calculator



6-B

4. The rate, in calories per minute, at which a person using an exercise machine burns calories is modeled by the function f . In the figure above, $f(t) = -\frac{1}{4}t^3 + \frac{3}{2}t^2 + 1$ for $0 \leq t \leq 4$ and f is piecewise linear for $4 \leq t \leq 24$.
- Find $f'(22)$. Indicate units of measure.
 - For the time interval $0 \leq t \leq 24$, at what time t is f increasing at its greatest rate? Show the reasoning that supports your answer.
 - Find the total number of calories burned over the time interval $6 \leq t \leq 18$ minutes.
 - The setting on the machine is now changed so that the person burns $f(t) + c$ calories per minute. For this setting, find c so that an average of 15 calories per minute is burned during the time interval $6 \leq t \leq 18$.



10-A

3. There are 700 people in line for a popular amusement-park ride when the ride begins operation in the morning. Once it begins operation, the ride accepts passengers until the park closes 8 hours later. While there is a line, people move onto the ride at a rate of 800 people per hour. The graph above shows the rate, $r(t)$, at which people arrive at the ride throughout the day. Time t is measured in hours from the time the ride begins operation.
- How many people arrive at the ride between $t = 0$ and $t = 3$? Show the computations that lead to your answer.
 - Is the number of people waiting in line to get on the ride increasing or decreasing between $t = 2$ and $t = 3$? Justify your answer.
 - At what time t is the line for the ride the longest? How many people are in line at that time? Justify your answers.
 - Write, but do not solve, an equation involving an integral expression of r whose solution gives the earliest time t at which there is no longer a line for the ride.

- 11B 2. A 12,000-liter tank of water is filled to capacity. At time $t = 0$, water begins to drain out of the tank at a rate modeled by $r(t)$, measured in liters per hour, where r is given by the piecewise-defined function

$$r(t) = \begin{cases} \frac{600t}{t+3} & \text{for } 0 \leq t \leq 5 \\ 1000e^{-0.2t} & \text{for } t > 5 \end{cases}$$

- Is r continuous at $t = 5$? Show the work that leads to your answer.
- Find the average rate at which water is draining from the tank between time $t = 0$ and time $t = 8$ hours.
- Find $r'(3)$. Using correct units, explain the meaning of that value in the context of this problem.
- Write, but do not solve, an equation involving an integral to find the time A when the amount of water in the tank is 9000 liters.

88

4. Water is pumped into an underground tank at a constant rate of 8 gallons per minute. Water leaks out of the tank at the rate of $\sqrt{t+1}$ gallons per minute, for $0 \leq t \leq 120$ minutes. At time $t = 0$, the tank contains 30 gallons of water.
- (a) How many gallons of water leak out of the tank from time $t = 0$ to $t = 3$ minutes?
 - (b) How many gallons of water are in the tank at time $t = 3$ minutes?
 - (c) Write an expression for $A(t)$, the total number of gallons of water in the tank at time t .
 - (d) At what time t , for $0 \leq t \leq 120$, is the amount of water in the tank a maximum? Justify your answer.

AP[®] CALCULUS AB 2002 SCORING GUIDELINES

Question 2

The rate at which people enter an amusement park on a given day is modeled by the function E defined by

$$E(t) = \frac{15600}{(t^2 - 24t + 160)}$$

The rate at which people leave the same amusement park on the same day is modeled by the function L defined by

$$L(t) = \frac{9890}{(t^2 - 38t + 370)}$$

Both $E(t)$ and $L(t)$ are measured in people per hour and time t is measured in hours after midnight. These functions are valid for $9 \leq t \leq 23$, the hours during which the park is open. At time $t = 9$, there are no people in the park.

- (a) How many people have entered the park by 5:00 P.M. ($t = 17$)? Round answer to the nearest whole number.
- (b) The price of admission to the park is \$15 until 5:00 P.M. ($t = 17$). After 5:00 P.M., the price of admission to the park is \$11. How many dollars are collected from admissions to the park on the given day? Round your answer to the nearest whole number.
- (c) Let $H(t) = \int_0^t (E(x) - L(x)) dx$ for $9 \leq t \leq 23$. The value of $H(17)$ to the nearest whole number is 3725. Find the value of $H'(17)$ and explain the meaning of $H(17)$ and $H'(17)$ in the context of the park.
- (d) At what time t , for $9 \leq t \leq 23$, does the model predict that the number of people in the park is a maximum?

(a) $\int_9^{17} E(t) dt = 6004.270$
6004 people entered the park by 5 pm.

(b) $15 \int_9^{17} E(t) dt + 11 \int_{17}^{23} E(t) dt = 104048.165$
The amount collected was \$104,048.

or
 $\int_{17}^{23} E(t) dt = 1271.283$
1271 people entered the park between 5 pm and 11 pm, so the amount collected was
 $\$15 \cdot (6004) + \$11 \cdot (1271) = \$104,041.$

(c) $H'(17) = E(17) - L(17) = -380.281$
There were 3725 people in the park at $t = 17$.
The number of people in the park was decreasing at the rate of approximately 380 people/hr at time $t = 17$.

(d) $H'(t) = E(t) - L(t) = 0$
 $t = 15.794$ or 15.795

$$\left\{ \begin{array}{l} 1 : \text{limits} \\ 3 : \text{integrand} \\ 1 : \text{answer} \end{array} \right.$$

1 : setup

$$\left\{ \begin{array}{l} 1 : \text{value of } H'(17) \\ 2 : \text{meanings} \\ 3 : \begin{array}{l} 1 : \text{meaning of } H(17) \\ 1 : \text{meaning of } H'(17) \\ < -1 > \text{ if no reference to } t = 17 \end{array} \end{array} \right.$$

$$\left\{ \begin{array}{l} 1 : E(t) - L(t) = 0 \\ 2 : \text{answer} \end{array} \right.$$

164

164

AP[®] CALCULUS AB
2002 SCORING GUIDELINES (Form B)

Question 2

The number of gallons, $P(t)$, of a pollutant in a lake changes at the rate $P'(t) = 1 - 3e^{-0.2\sqrt{t}}$ gallons per day, where t is measured in days. There are 50 gallons of the pollutant in the lake at time $t = 0$. The lake is considered to be safe when it contains 40 gallons or less of pollutant.

- (a) Is the amount of pollutant increasing at time $t = 9$? Why or why not?
- (b) For what value of t will the number of gallons of pollutant be at its minimum? Justify your answer.
- (c) Is the lake safe when the number of gallons of pollutant is at its minimum? Justify your answer.
- (d) An investigator uses the tangent line approximation to $P(t)$ at $t = 0$ as a model for the amount of pollutant in the lake. At what time t does this model predict that the lake becomes safe?

<p>(a) $P'(9) = 1 - 3e^{-0.6} = -0.646 < 0$ so the amount is not increasing at this time.</p>	<p>1 : answer with reason</p>
<p>(b) $P'(t) = 1 - 3e^{-0.2\sqrt{t}} = 0$ $t = (5 \ln 3)^2 = 30.174$ $P'(t)$ is negative for $0 < t < (5 \ln 3)^2$ and positive for $t > (5 \ln 3)^2$. Therefore there is a minimum at $t = (5 \ln 3)^2$.</p>	<p>3 {</p> <ul style="list-style-type: none"> 1 : sets $P'(t) = 0$ 1 : solves for t 1 : justification
<p>(c) $P(30.174) = 50 + \int_0^{30.174} (1 - 3e^{-0.2\sqrt{t}}) dt$ $= 35.104 < 40$, so the lake is safe.</p>	<p>3 {</p> <ul style="list-style-type: none"> 1 : integrand 1 : limits 1 : conclusion with reason based on integral of $P'(t)$
<p>(d) $P'(0) = 1 - 3 = -2$. The lake will become safe when the amount decreases by 10. A linear model predicts this will happen when $t = 5$.</p>	<p>2 {</p> <ul style="list-style-type: none"> 1 : slope of tangent line 1 : answer

AP[®] CALCULUS AB
2003 SCORING GUIDELINES (Form B)

Question 2

A tank contains 125 gallons of heating oil at time $t = 0$. During the time interval $0 \leq t \leq 12$ hours, heating oil is pumped into the tank at the rate

$$H(t) = 2 + \frac{10}{(1 + \ln(t + 1))} \text{ gallons per hour.}$$

During the same time interval, heating oil is removed from the tank at the rate

$$R(t) = 12 \sin\left(\frac{t^2}{47}\right) \text{ gallons per hour.}$$

- (a) How many gallons of heating oil are pumped into the tank during the time interval $0 \leq t \leq 12$ hours?
 (b) Is the level of heating oil in the tank rising or falling at time $t = 6$ hours? Give a reason for your answer.
 (c) How many gallons of heating oil are in the tank at time $t = 12$ hours?
 (d) At what time t , for $0 \leq t \leq 12$, is the volume of heating oil in the tank the least? Show the analysis that leads to your conclusion.

(a) $\int_0^{12} H(t) dt = 70.570$ or 70.571

2 : $\left\{ \begin{array}{l} 1 : \text{integral} \\ 1 : \text{answer} \end{array} \right.$

(b) $H(6) - R(6) = -2.924$,
 so the level of heating oil is falling at $t = 6$.

1 : answer with reason

(c) $125 + \int_0^{12} (H(t) - R(t)) dt = 122.025$ or 122.026

3 : $\left\{ \begin{array}{l} 1 : \text{limits} \\ 1 : \text{integrand} \\ 1 : \text{answer} \end{array} \right.$

(d) The absolute minimum occurs at a critical point or an endpoint.

$H(t) - R(t) = 0$ when $t = 4.790$ and $t = 11.318$.

3 : $\left\{ \begin{array}{l} 1 : \text{sets } H(t) - R(t) = 0 \\ 1 : \text{volume is least at} \\ t = 11.318 \end{array} \right.$

The volume increases until $t = 4.790$, then decreases until $t = 11.318$, then increases, so the absolute minimum will be at $t = 0$ or at $t = 11.318$.

1 : analysis for absolute minimum

$125 + \int_0^{11.318} (H(t) - R(t)) dt = 120.738$

Since the volume is 125 at $t = 0$, the volume is least at $t = 11.318$.

AP[®] CALCULUS AB
2005 SCORING GUIDELINES (Form B)

Question 2

A water tank at Camp Newton holds 1200 gallons of water at time $t = 0$. During the time interval $0 \leq t \leq 18$ hours, water is pumped into the tank at the rate

$$W(t) = 95\sqrt{t} \sin^2\left(\frac{t}{6}\right) \text{ gallons per hour.}$$

During the same time interval, water is removed from the tank at the rate

$$R(t) = 275 \sin^2\left(\frac{t}{3}\right) \text{ gallons per hour.}$$

- (a) Is the amount of water in the tank increasing at time $t = 15$? Why or why not?
 (b) To the nearest whole number, how many gallons of water are in the tank at time $t = 18$?
 (c) At what time t , for $0 \leq t \leq 18$, is the amount of water in the tank at an absolute minimum? Show the work that leads to your conclusion.
 (d) For $t > 18$, no water is pumped into the tank, but water continues to be removed at the rate $R(t)$ until the tank becomes empty. Let k be the time at which the tank becomes empty. Write, but do not solve, an equation involving an integral expression that can be used to find the value of k .

- (a) No; the amount of water is not increasing at $t = 15$ since $W(15) - R(15) = -121.09 < 0$.

1 : answer with reason

- (b) $1200 + \int_0^{18} (W(t) - R(t)) dt = 1309.788$
 1310 gallons

3 : { 1 : limits
 1 : integrand
 1 : answer

- (c) $W(t) - R(t) = 0$
 $t = 0, 6.4948, 12.9748$

3 : { 1 : interior critical points
 1 : amount of water is least at
 $t = 6.494$ or 6.495
 1 : analysis for absolute minimum

t (hours)	gallons of water
0	1200
6.495	525
12.975	1697
18	1310

The values at the endpoints and the critical points show that the absolute minimum occurs when $t = 6.494$ or 6.495 .

- (d) $\int_{18}^k R(t) dt = 1310$

2 : { 1 : limits
 1 : equation

**AP[®] CALCULUS AB
2005 SCORING GUIDELINES**

Question 2

The tide removes sand from Sandy Point Beach at a rate modeled by the function R , given by

$$R(t) = 2 + 5 \sin\left(\frac{4\pi t}{25}\right).$$

A pumping station adds sand to the beach at a rate modeled by the function S , given by

$$S(t) = \frac{15t}{1 + 3t}.$$

Both $R(t)$ and $S(t)$ have units of cubic yards per hour and t is measured in hours for $0 \leq t \leq 6$. At time $t = 0$, the beach contains 2500 cubic yards of sand.

- (a) How much sand will the tide remove from the beach during this 6-hour period? Indicate units of measure.
 (b) Write an expression for $Y(t)$, the total number of cubic yards of sand on the beach at time t .
 (c) Find the rate at which the total amount of sand on the beach is changing at time $t = 4$.
 (d) For $0 \leq t \leq 6$, at what time t is the amount of sand on the beach a minimum? What is the minimum value?
 Justify your answers.

(a) $\int_0^6 R(t) dt = 31.815$ or 31.816 yd^3

2 : { 1 : integral
1 : answer with units

(b) $Y(t) = 2500 + \int_0^t (S(x) - R(x)) dx$

3 : { 1 : integrand
1 : limits
1 : answer

(c) $Y'(t) = S(t) - R(t)$
 $Y'(4) = S(4) - R(4) = -1.908$ or $-1.909 \text{ yd}^3/\text{hr}$

1 : answer

- (d) $Y'(t) = 0$ when $S(t) - R(t) = 0$.
 The only value in $[0, 6]$ to satisfy $S(t) = R(t)$
 is $a = 5.117865$.

3 : { 1 : sets $Y'(t) = 0$
1 : critical t -value
1 : answer with justification

t	$Y(t)$
0	2500
a	2492.3694
6	2493.2766

The amount of sand is a minimum when $t = 5.117$ or 5.118 hours. The minimum value is 2492.369 cubic yards.

**AP[®] CALCULUS AB
2004 SCORING GUIDELINES**

Question 1

Traffic flow is defined as the rate at which cars pass through an intersection, measured in cars per minute. The traffic flow at a particular intersection is modeled by the function F defined by

$$F(t) = 82 + 4 \sin\left(\frac{t}{2}\right) \text{ for } 0 \leq t \leq 30,$$

where $F(t)$ is measured in cars per minute and t is measured in minutes.

- (a) To the nearest whole number, how many cars pass through the intersection over the 30-minute period?
- (b) Is the traffic flow increasing or decreasing at $t = 7$? Give a reason for your answer.
- (c) What is the average value of the traffic flow over the time interval $10 \leq t \leq 15$? Indicate units of measure.
- (d) What is the average rate of change of the traffic flow over the time interval $10 \leq t \leq 15$? Indicate units of measure.

(a) $\int_0^{30} F(t) dt = 2474$ cars

3 : $\begin{cases} 1 : \text{limits} \\ 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(b) $F'(7) = -1.872$ or -1.873
Since $F'(7) < 0$, the traffic flow is decreasing at $t = 7$.

1 : answer with reason

(c) $\frac{1}{5} \int_{10}^{15} F(t) dt = 81.899$ cars/min

3 : $\begin{cases} 1 : \text{limits} \\ 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(d) $\frac{F(15) - F(10)}{15 - 10} = 1.517$ or 1.518 cars/min²

1 : answer

Units of cars/min in (c) and cars/min² in (d)

1 : units in (c) and (d)

**AP[®] CALCULUS AB
2009 SCORING GUIDELINES**

Question 2

The rate at which people enter an auditorium for a rock concert is modeled by the function R given by $R(t) = 1380t^2 - 675t^3$ for $0 \leq t \leq 2$ hours; $R(t)$ is measured in people per hour. No one is in the auditorium at time $t = 0$, when the doors open. The doors close and the concert begins at time $t = 2$.

- (a) How many people are in the auditorium when the concert begins?
- (b) Find the time when the rate at which people enter the auditorium is a maximum. Justify your answer.
- (c) The total wait time for all the people in the auditorium is found by adding the time each person waits, starting at the time the person enters the auditorium and ending when the concert begins. The function w models the total wait time for all the people who enter the auditorium before time t . The derivative of w is given by $w'(t) = (2 - t)R(t)$. Find $w(2) - w(1)$, the total wait time for those who enter the auditorium after time $t = 1$.
- (d) On average, how long does a person wait in the auditorium for the concert to begin? Consider all people who enter the auditorium after the doors open, and use the model for total wait time from part (c).

(a) $\int_0^2 R(t) dt = 980$ people

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b) $R'(t) = 0$ when $t = 0$ and $t = 1.36296$
The maximum rate may occur at 0, $a = 1.36296$, or 2.

$R(0) = 0$
 $R(a) = 854.527$
 $R(2) = 120$

The maximum rate occurs when $t = 1.362$ or 1.363 .

3 : $\begin{cases} 1 : \text{considers } R'(t) = 0 \\ 1 : \text{interior critical point} \\ 1 : \text{answer and justification} \end{cases}$

(c) $w(2) - w(1) = \int_1^2 w'(t) dt = \int_1^2 (2 - t)R(t) dt = 387.5$

The total wait time for those who enter the auditorium after time $t = 1$ is 387.5 hours.

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(d) $\frac{1}{980}w(2) = \frac{1}{980} \int_0^2 (2 - t)R(t) dt = 0.77551$


On average, a person waits 0.775 or 0.776 hour.

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

**AP[®] CALCULUS AB
2006 SCORING GUIDELINES**

Question 2

At an intersection in Thomasville, Oregon, cars turn left at the rate $L(t) = 60\sqrt{t} \sin^2\left(\frac{t}{3}\right)$ cars per hour over the time interval $0 \leq t \leq 18$ hours. The graph of $y = L(t)$ is shown above.



- (a) To the nearest whole number, find the total number of cars turning left at the intersection over the time interval $0 \leq t \leq 18$ hours.
- (b) Traffic engineers will consider turn restrictions when $L(t) \geq 150$ cars per hour. Find all values of t for which $L(t) \geq 150$ and compute the average value of L over this time interval. Indicate units of measure.
- (c) Traffic engineers will install a signal if there is any two-hour time interval during which the product of the total number of cars turning left and the total number of oncoming cars traveling straight through the intersection is greater than 200,000. In every two-hour time interval, 500 oncoming cars travel straight through the intersection. Does this intersection require a traffic signal? Explain the reasoning that leads to your conclusion.

(a) $\int_0^{18} L(t) dt \approx 1658$ cars

2 : { 1 : setup
1 : answer

(b) $L(t) = 150$ when $t = 12.42831, 16.12166$
Let $R = 12.42831$ and $S = 16.12166$
 $L(t) \geq 150$ for t in the interval $[R, S]$

$$\frac{1}{S - R} \int_R^S L(t) dt = 199.426 \text{ cars per hour}$$

3 : { 1 : t -interval when $L(t) \geq 150$
1 : average value integral
1 : answer with units

- (c) For the product to exceed 200,000, the number of cars turning left in a two-hour interval must be greater than 400.

$$\int_{13}^{15} L(t) dt = 431.931 > 400$$

OR

The number of cars turning left will be greater than 400 on a two-hour interval if $L(t) \geq 200$ on that interval.

$$L(t) \geq 200 \text{ on any two-hour subinterval of } [13.25304, 15.32386].$$

Yes, a traffic signal is required.

4 : { 1 : considers 400 cars
1 : valid interval $[h, h + 2]$
1 : value of $\int_h^{h+2} L(t) dt$
1 : answer and explanation

OR

4 : { 1 : considers 200 cars per hour
1 : solves $L(t) \geq 200$
1 : discusses 2 hour interval
1 : answer and explanation

AP[®] CALCULUS AB
2009 SCORING GUIDELINES (Form B)

Question 1

At a certain height, a tree trunk has a circular cross section. The radius $R(t)$ of that cross section grows at a rate modeled by the function

$$\frac{dR}{dt} = \frac{1}{16}(3 + \sin(t^2)) \text{ centimeters per year}$$

for $0 \leq t \leq 3$, where time t is measured in years. At time $t = 0$, the radius is 6 centimeters. The area of the cross section at time t is denoted by $A(t)$.

- (a) Write an expression, involving an integral, for the radius $R(t)$ for $0 \leq t \leq 3$. Use your expression to find $R(3)$.
- (b) Find the rate at which the cross-sectional area $A(t)$ is increasing at time $t = 3$ years. Indicate units of measure.
- (c) Evaluate $\int_0^3 A'(t) dt$. Using appropriate units, interpret the meaning of that integral in terms of cross-sectional area.

(a) $R(t) = 6 + \int_0^t \frac{1}{16}(3 + \sin(x^2)) dx$
 $R(3) = 6.610$ or 6.611

3 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{expression for } R(t) \\ 1 : R(3) \end{cases}$

(b) $A(t) = \pi(R(t))^2$
 $A'(t) = 2\pi R(t)R'(t)$
 $A'(3) = 8.858 \text{ cm}^2/\text{year}$

3 : $\begin{cases} 1 : \text{expression for } A(t) \\ 1 : \text{expression for } A'(t) \\ 1 : \text{answer with units} \end{cases}$

(c) $\int_0^3 A'(t) dt = A(3) - A(0) = 24.200$ or 24.201

From time $t = 0$ to $t = 3$ years, the cross-sectional area grows by 24.201 square centimeters.

3 : $\begin{cases} 1 : \text{uses Fundamental Theorem of Calculus} \\ 1 : \text{value of } \int_0^3 A'(t) dt \\ 1 : \text{meaning of } \int_0^3 A'(t) dt \end{cases}$

AP[®] CALCULUS AB
2011 SCORING GUIDELINES (Form B)

Question 1

A cylindrical can of radius 10 millimeters is used to measure rainfall in Stormville. The can is initially empty, and rain enters the can during a 60-day period. The height of water in the can is modeled by the function S , where $S(t)$ is measured in millimeters and t is measured in days for $0 \leq t \leq 60$. The rate at which the height of the water is rising in the can is given by $S'(t) = 2\sin(0.03t) + 1.5$.

- (a) According to the model, what is the height of the water in the can at the end of the 60-day period?
- (b) According to the model, what is the average rate of change in the height of water in the can over the 60-day period? Show the computations that lead to your answer. Indicate units of measure.
- (c) Assuming no evaporation occurs, at what rate is the volume of water in the can changing at time $t = 7$? Indicate units of measure.
- (d) During the same 60-day period, rain on Monsoon Mountain accumulates in a can identical to the one in Stormville. The height of the water in the can on Monsoon Mountain is modeled by the function M , where $M(t) = \frac{1}{400}(3t^3 - 30t^2 + 330t)$. The height $M(t)$ is measured in millimeters, and t is measured in days for $0 \leq t \leq 60$. Let $D(t) = M'(t) - S'(t)$. Apply the Intermediate Value Theorem to the function D on the interval $0 \leq t \leq 60$ to justify that there exists a time t , $0 < t < 60$, at which the heights of water in the two cans are changing at the same rate.

(a) $S(60) = \int_0^{60} S'(t) dt = 171.813 \text{ mm}$

3 : $\begin{cases} 1 : \text{limits} \\ 1 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(b) $\frac{S(60) - S(0)}{60} = 2.863 \text{ or } 2.864 \text{ mm/day}$

1 : answer

(c) $V(t) = 100\pi S(t)$
 $V'(7) = 100\pi S'(7) = 602.218$

The volume of water in the can is increasing at a rate of $602.218 \text{ mm}^3/\text{day}$.

2 : $\begin{cases} 1 : \text{relationship between } V \text{ and } S \\ 1 : \text{answer} \end{cases}$

(d) $D(0) = -0.675 < 0$ and $D(60) = 69.37730 > 0$

Because D is continuous, the Intermediate Value Theorem implies that there is a time t , $0 < t < 60$, at which $D(t) = 0$. At this time, the heights of water in the two cans are changing at the same rate.

2 : $\begin{cases} 1 : \text{considers } D(0) \text{ and } D(60) \\ 1 : \text{justification} \end{cases}$

1 : units in (b) or (c)

AP[®] CALCULUS AB
2007 SCORING GUIDELINES (Form B)

Question 3

The wind chill is the temperature, in degrees Fahrenheit ($^{\circ}\text{F}$), a human feels based on the air temperature, in degrees Fahrenheit, and the wind velocity v , in miles per hour (mph). If the air temperature is 32°F , then the wind chill is given by $W(v) = 55.6 - 22.1v^{0.16}$ and is valid for $5 \leq v \leq 60$.

- (a) Find $W'(20)$. Using correct units, explain the meaning of $W'(20)$ in terms of the wind chill.
- (b) Find the average rate of change of W over the interval $5 \leq v \leq 60$. Find the value of v at which the instantaneous rate of change of W is equal to the average rate of change of W over the interval $5 \leq v \leq 60$.
- (c) Over the time interval $0 \leq t \leq 4$ hours, the air temperature is a constant 32°F . At time $t = 0$, the wind velocity is $v = 20$ mph. If the wind velocity increases at a constant rate of 5 mph per hour, what is the rate of change of the wind chill with respect to time at $t = 3$ hours? Indicate units of measure.

(a) $W'(20) = -22.1 \cdot 0.16 \cdot 20^{-0.84} = -0.285$ or -0.286

When $v = 20$ mph, the wind chill is decreasing at $0.286^{\circ}\text{F}/\text{mph}$.

2 : { 1 : value
1 : explanation

(b) The average rate of change of W over the interval $5 \leq v \leq 60$ is $\frac{W(60) - W(5)}{60 - 5} = -0.253$ or -0.254 .

$W'(v) = \frac{W(60) - W(5)}{60 - 5}$ when $v = 23.011$.

3 : { 1 : average rate of change
1 : $W'(v) =$ average rate of change
1 : value of v

(c) $\left. \frac{dW}{dt} \right|_{t=3} = \left(\frac{dW}{dv} \cdot \frac{dv}{dt} \right) \Big|_{t=3} = W'(35) \cdot 5 = -0.892^{\circ}\text{F}/\text{hr}$

OR

$W = 55.6 - 22.1(20 + 5t)^{0.16}$

$\left. \frac{dW}{dt} \right|_{t=3} = -0.892^{\circ}\text{F}/\text{hr}$

3 : { 1 : $\frac{dv}{dt} = 5$
1 : uses $v(3) = 35$,
or
uses $v(t) = 20 + 5t$
1 : answer

Units of $^{\circ}\text{F}/\text{mph}$ in (a) and $^{\circ}\text{F}/\text{hr}$ in (c)

1 : units in (a) and (c)

AP[®] CALCULUS AB
2009 SCORING GUIDELINES (Form B)

Question 2

A storm washed away sand from a beach, causing the edge of the water to get closer to a nearby road. The rate at which the distance between the road and the edge of the water was changing during the storm is modeled by $f(t) = \sqrt{t} + \cos t - 3$ meters per hour, t hours after the storm began. The edge of the water was 35 meters from the road when the storm began, and the storm lasted 5 hours. The derivative of $f(t)$ is $f'(t) = \frac{1}{2\sqrt{t}} - \sin t$.

- What was the distance between the road and the edge of the water at the end of the storm?
- Using correct units, interpret the value $f'(4) = 1.007$ in terms of the distance between the road and the edge of the water.
- At what time during the 5 hours of the storm was the distance between the road and the edge of the water decreasing most rapidly? Justify your answer.
- After the storm, a machine pumped sand back onto the beach so that the distance between the road and the edge of the water was growing at a rate of $g(p)$ meters per day, where p is the number of days since pumping began. Write an equation involving an integral expression whose solution would give the number of days that sand must be pumped to restore the original distance between the road and the edge of the water.

(a) $35 + \int_0^5 f(t) dt = 26.494$ or 26.495 meters

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b) Four hours after the storm began, the rate of change of the distance between the road and the edge of the water is increasing at a rate of 1.007 meters/hours².

2 : $\begin{cases} 1 : \text{interpretation of } f'(4) \\ 1 : \text{units} \end{cases}$

(c) $f'(t) = 0$ when $t = 0.66187$ and $t = 2.84038$
 The minimum of f for $0 \leq t \leq 5$ may occur at 0, 0.66187, 2.84038, or 5.

3 : $\begin{cases} 1 : \text{considers } f'(t) = 0 \\ 1 : \text{answer} \\ 1 : \text{justification} \end{cases}$

$$\begin{aligned} f(0) &= -2 \\ f(0.66187) &= -1.39760 \\ f(2.84038) &= -2.26963 \\ f(5) &= -0.48027 \end{aligned}$$

The distance between the road and the edge of the water was decreasing most rapidly at time $t = 2.840$ hours after the storm began.

(d) $-\int_0^5 f(t) dt = \int_0^x g(p) dp$

2 : $\begin{cases} 1 : \text{integral of } g \\ 1 : \text{answer} \end{cases}$

AP[®] CALCULUS AB
2008 SCORING GUIDELINES (Form B)

Question 2

For time $t \geq 0$ hours, let $r(t) = 120(1 - e^{-10t^2})$ represent the speed, in kilometers per hour, at which a car travels along a straight road. The number of liters of gasoline used by the car to travel x kilometers is modeled by $g(x) = 0.05x(1 - e^{-x/2})$.

- (a) How many kilometers does the car travel during the first 2 hours?
 (b) Find the rate of change with respect to time of the number of liters of gasoline used by the car when $t = 2$ hours. Indicate units of measure.
 (c) How many liters of gasoline have been used by the car when it reaches a speed of 80 kilometers per hour?

(a) $\int_0^2 r(t) dt = 206.370$ kilometers

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b) $\frac{dg}{dt} = \frac{dg}{dx} \cdot \frac{dx}{dt}; \quad \frac{dx}{dt} = r(t)$
 $\left. \frac{dg}{dt} \right|_{t=2} = \left. \frac{dg}{dx} \right|_{x=206.370} \cdot r(2)$
 $= (0.050)(120) = 6$ liters/hour

3 : $\begin{cases} 2 : \text{uses chain rule} \\ 1 : \text{answer with units} \end{cases}$

- (c) Let T be the time at which the car's speed reaches 80 kilometers per hour.

Then, $r(T) = 80$ or $T = 0.331453$ hours.

At time T , the car has gone

$x(T) = \int_0^T r(t) dt = 10.794097$ kilometers

and has consumed $g(x(T)) = 0.537$ liters of gasoline.

4 : $\begin{cases} 1 : \text{equation } r(t) = 80 \\ 2 : \text{distance integral} \\ 1 : \text{answer} \end{cases}$

**AP[®] CALCULUS AB
2010 SCORING GUIDELINES**

Question 1

There is no snow on Janet's driveway when snow begins to fall at midnight. From midnight to 9 A.M., snow accumulates on the driveway at a rate modeled by $f(t) = 7te^{\cos t}$ cubic feet per hour, where t is measured in hours since midnight. Janet starts removing snow at 6 A.M. ($t = 6$). The rate $g(t)$, in cubic feet per hour, at which Janet removes snow from the driveway at time t hours after midnight is modeled by

$$g(t) = \begin{cases} 0 & \text{for } 0 \leq t < 6 \\ 125 & \text{for } 6 \leq t < 7 \\ 108 & \text{for } 7 \leq t \leq 9. \end{cases}$$

- (a) How many cubic feet of snow have accumulated on the driveway by 6 A.M.?
 (b) Find the rate of change of the volume of snow on the driveway at 8 A.M.
 (c) Let $h(t)$ represent the total amount of snow, in cubic feet, that Janet has removed from the driveway at time t hours after midnight. Express h as a piecewise-defined function with domain $0 \leq t \leq 9$.
 (d) How many cubic feet of snow are on the driveway at 9 A.M.?

(a) $\int_0^6 f(t) dt = 142.274$ or 142.275 cubic feet

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b) Rate of change is $f(8) - g(8) = -59.582$ or -59.583 cubic feet per hour.

1 : answer

(c) $h(0) = 0$

For $0 < t \leq 6$, $h(t) = h(0) + \int_0^t g(s) ds = 0 + \int_0^t 0 ds = 0$.

For $6 < t \leq 7$, $h(t) = h(6) + \int_6^t g(s) ds = 0 + \int_6^t 125 ds = 125(t - 6)$.

For $7 < t \leq 9$, $h(t) = h(7) + \int_7^t g(s) ds = 125 + \int_7^t 108 ds = 125 + 108(t - 7)$.

Thus, $h(t) = \begin{cases} 0 & \text{for } 0 \leq t \leq 6 \\ 125(t - 6) & \text{for } 6 < t \leq 7 \\ 125 + 108(t - 7) & \text{for } 7 < t \leq 9 \end{cases}$

3 : $\begin{cases} 1 : h(t) \text{ for } 0 \leq t \leq 6 \\ 1 : h(t) \text{ for } 6 < t \leq 7 \\ 1 : h(t) \text{ for } 7 < t \leq 9 \end{cases}$

(d) Amount of snow is $\int_0^9 f(t) dt - h(9) = 26.334$ or 26.335 cubic feet.

3 : $\begin{cases} 1 : \text{integral} \\ 1 : h(9) \\ 1 : \text{answer} \end{cases}$

AP[®] CALCULUS AB
2004 SCORING GUIDELINES (Form B)

Question 2

For $0 \leq t \leq 31$, the rate of change of the number of mosquitoes on Tropical Island at time t days is modeled by $R(t) = 5\sqrt{t} \cos\left(\frac{t}{5}\right)$ mosquitoes per day. There are 1000 mosquitoes on Tropical Island at time $t = 0$.

- (a) Show that the number of mosquitoes is increasing at time $t = 6$.
 (b) At time $t = 6$, is the number of mosquitoes increasing at an increasing rate, or is the number of mosquitoes increasing at a decreasing rate? Give a reason for your answer.
 (c) According to the model, how many mosquitoes will be on the island at time $t = 31$? Round your answer to the nearest whole number.
 (d) To the nearest whole number, what is the maximum number of mosquitoes for $0 \leq t \leq 31$? Show the analysis that leads to your conclusion.

(a) Since $R(6) = 4.438 > 0$, the number of mosquitoes is increasing at $t = 6$.

1 : shows that $R(6) > 0$

(b) $R'(6) = -1.913$
 Since $R'(6) < 0$, the number of mosquitoes is increasing at a decreasing rate at $t = 6$.

2 : $\left\{ \begin{array}{l} 1 : \text{considers } R'(6) \\ 1 : \text{answer with reason} \end{array} \right.$

(c) $1000 + \int_0^{31} R(t) dt = 964.335$
 To the nearest whole number, there are 964 mosquitoes.

2 : $\left\{ \begin{array}{l} 1 : \text{integral} \\ 1 : \text{answer} \end{array} \right.$

(d) $R(t) = 0$ when $t = 0$, $t = 2.5\pi$, or $t = 7.5\pi$
 $R(t) > 0$ on $0 < t < 2.5\pi$
 $R(t) < 0$ on $2.5\pi < t < 7.5\pi$
 $R(t) > 0$ on $7.5\pi < t < 31$
 The absolute maximum number of mosquitoes occurs at $t = 2.5\pi$ or at $t = 31$.

$$1000 + \int_0^{2.5\pi} R(t) dt = 1039.357,$$

There are 964 mosquitoes at $t = 31$, so the maximum number of mosquitoes is 1039, to the nearest whole number.

2 : absolute maximum value
 1 : integral
 1 : answer
 4 : $\left\{ \begin{array}{l} 2 : \text{analysis} \\ 1 : \text{computes interior critical points} \\ 1 : \text{completes analysis} \end{array} \right.$

**AP[®] CALCULUS AB
2007 SCORING GUIDELINES**


Question 2

The amount of water in a storage tank, in gallons, is modeled by a continuous function on the time interval $0 \leq t \leq 7$, where t is measured in hours. In this model, rates are given as follows:

(i) The rate at which water enters the tank is $f(t) = 100t^2 \sin(\sqrt{t})$ gallons per hour for $0 \leq t \leq 7$.

(ii) The rate at which water leaves the tank is

$$g(t) = \begin{cases} 250 & \text{for } 0 \leq t < 3 \\ 2000 & \text{for } 3 < t \leq 7 \end{cases} \text{ gallons per hour.}$$



The graphs of f and g , which intersect at $t = 1.617$ and $t = 5.076$, are shown in the figure above. At time $t = 0$, the amount of water in the tank is 5000 gallons.

- (a) How many gallons of water enter the tank during the time interval $0 \leq t \leq 7$? Round your answer to the nearest gallon.
- (b) For $0 \leq t \leq 7$, find the time intervals during which the amount of water in the tank is decreasing. Give a reason for each answer.
- (c) For $0 \leq t \leq 7$, at what time t is the amount of water in the tank greatest? To the nearest gallon, compute the amount of water at this time. Justify your answer.

(a) $\int_0^7 f(t) dt \approx 8264$ gallons

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

(b) The amount of water in the tank is decreasing on the intervals $0 \leq t \leq 1.617$ and $3 \leq t \leq 5.076$ because $f(t) < g(t)$ for $0 \leq t < 1.617$ and $3 < t < 5.076$.

2 : $\begin{cases} 1 : \text{intervals} \\ 1 : \text{reason} \end{cases}$

(c) Since $f(t) - g(t)$ changes sign from positive to negative only at $t = 3$, the candidates for the absolute maximum are at $t = 0, 3$, and 7 .

5 : $\begin{cases} 1 : \text{identifies } t = 3 \text{ as a candidate} \\ 1 : \text{integrand} \\ 1 : \text{amount of water at } t = 3 \\ 1 : \text{amount of water at } t = 7 \\ 1 : \text{conclusion} \end{cases}$


t (hours)	gallons of water
0	5000
3	$5000 + \int_0^3 f(t) dt - 250(3) = 5126.591$
7	$5126.591 + \int_3^7 f(t) dt - 2000(4) = 4513.807$

The amount of water in the tank is greatest at 3 hours. At that time, the amount of water in the tank, rounded to the nearest gallon, is 5127 gallons.

**AP[®] CALCULUS AB
2006 SCORING GUIDELINES (Form B)**

Question 4

The rate, in calories per minute, at which a person using an exercise machine burns calories is modeled by the function f . In the figure above, $f(t) = -\frac{1}{4}t^3 + \frac{3}{2}t^2 + 1$ for $0 \leq t \leq 4$ and f is piecewise linear for $4 \leq t \leq 24$.



- (a) Find $f'(22)$. Indicate units of measure.
 (b) For the time interval $0 \leq t \leq 24$, at what time t is f increasing at its greatest rate? Show the reasoning that supports your answer.
 (c) Find the total number of calories burned over the time interval $6 \leq t \leq 18$ minutes.
 (d) The setting on the machine is now changed so that the person burns $f(t) + c$ calories per minute. For this setting, find c so that an average of 15 calories per minute is burned during the time interval $6 \leq t \leq 18$.

(a) $f'(22) = \frac{15 - 3}{20 - 24} = -3$ calories/min/min

(b) f is increasing on $[0, 4]$ and on $[12, 16]$.

On $(12, 16)$, $f'(t) = \frac{15 - 9}{16 - 12} = \frac{3}{2}$ since f has constant slope on this interval.

On $(0, 4)$, $f'(t) = -\frac{3}{4}t^2 + 3t$ and

$f''(t) = -\frac{3}{2}t + 3 = 0$ when $t = 2$. This is where f' has a maximum on $[0, 4]$ since $f'' > 0$ on $(0, 2)$ and $f'' < 0$ on $(2, 4)$.

On $[0, 24]$, f is increasing at its greatest rate when $t = 2$ because $f'(2) = 3 > \frac{3}{2}$.

(c) $\int_6^{18} f(t) dt = 6(9) + \frac{1}{2}(4)(9 + 15) + 2(15) = 132$ calories

(d) We want $\frac{1}{12} \int_6^{18} (f(t) + c) dt = 15$.

This means $132 + 12c = 15(12)$. So, $c = 4$.

OR

Currently, the average is $\frac{132}{12} = 11$ calories/min.

Adding c to $f(t)$ will shift the average by c .

So $c = 4$ to get an average of 15 calories/min.

1 : $f'(22)$ and units

4 : $\begin{cases} 1 : f' \text{ on } (0, 4) \\ 1 : \text{shows } f' \text{ has a max at } t = 2 \text{ on } (0, 4) \\ 1 : \text{shows for } 12 < t < 16, f'(t) < f'(2) \\ 1 : \text{answer} \end{cases}$


2 : $\begin{cases} 1 : \text{method} \\ 1 : \text{answer} \end{cases}$

2 : $\begin{cases} 1 : \text{setup} \\ 1 : \text{value of } c \end{cases}$

**AP[®] CALCULUS AB
2010 SCORING GUIDELINES**

Question 3

There are 700 people in line for a popular amusement-park ride when the ride begins operation in the morning. Once it begins operation, the ride accepts passengers until the park closes 8 hours later. While there is a line, people move onto the ride at a rate of 800 people per hour. The graph above shows the rate, $r(t)$, at which people arrive at the ride throughout the day. Time t is measured in hours from the time the ride begins operation.



- (a) How many people arrive at the ride between $t = 0$ and $t = 3$? Show the computations that lead to your answer.
- (b) Is the number of people waiting in line to get on the ride increasing or decreasing between $t = 2$ and $t = 3$? Justify your answer.
- (c) At what time t is the line for the ride the longest? How many people are in line at that time? Justify your answers.
- (d) Write, but do not solve, an equation involving an integral expression of r whose solution gives the earliest time t at which there is no longer a line for the ride.

(a) $\int_0^3 r(t) dt = 2 \cdot \frac{1000 + 1200}{2} + \frac{1200 + 800}{2} = 3200$ people

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

- (b) The number of people waiting in line is increasing because people move onto the ride at a rate of 800 people per hour and for $2 < t < 3$, $r(t) > 800$.

1 : answer with reason

- (c) $r(t) = 800$ only at $t = 3$
 For $0 \leq t < 3$, $r(t) > 800$. For $3 < t \leq 8$, $r(t) < 800$.
 Therefore, the line is longest at time $t = 3$.
 There are $700 + 3200 - 800 \cdot 3 = 1500$ people waiting in line at time $t = 3$.

3 : $\begin{cases} 1 : \text{identifies } t = 3 \\ 1 : \text{number of people in line} \\ 1 : \text{justification} \end{cases}$

(d) $0 = 700 + \int_0^t r(s) ds - 800t$

3 : $\begin{cases} 1 : 800t \\ 1 : \text{integral} \\ 1 : \text{answer} \end{cases}$

AP[®] CALCULUS AB
2011 SCORING GUIDELINES (Form B)

Question 2

A 12,000-liter tank of water is filled to capacity. At time $t = 0$, water begins to drain out of the tank at a rate modeled by $r(t)$, measured in liters per hour, where r is given by the piecewise-defined function

$$r(t) = \begin{cases} \frac{600t}{t+3} & \text{for } 0 \leq t \leq 5 \\ 1000e^{-0.2t} & \text{for } t > 5 \end{cases}$$

- (a) Is r continuous at $t = 5$? Show the work that leads to your answer.
 (b) Find the average rate at which water is draining from the tank between time $t = 0$ and time $t = 8$ hours.
 (c) Find $r'(3)$. Using correct units, explain the meaning of that value in the context of this problem.
 (d) Write, but do not solve, an equation involving an integral to find the time A when the amount of water in the tank is 9000 liters.

(a) $\lim_{t \rightarrow 5^-} r(t) = \lim_{t \rightarrow 5^-} \left(\frac{600t}{t+3} \right) = 375 = r(5)$
 $\lim_{t \rightarrow 5^+} r(t) = \lim_{t \rightarrow 5^+} (1000e^{-0.2t}) = 367.879$

Because the left-hand and right-hand limits are not equal, r is not continuous at $t = 5$.

2 : conclusion with analysis

(b) $\frac{1}{8} \int_0^8 r(t) dt = \frac{1}{8} \left(\int_0^5 \frac{600t}{t+3} dt + \int_5^8 1000e^{-0.2t} dt \right)$
 $= 258.052$ or 258.053

3 : $\begin{cases} 1 : \text{integrand} \\ 1 : \text{limits and constant} \\ 1 : \text{answer} \end{cases}$

(c) $r'(3) = 50$

The rate at which water is draining out of the tank at time $t = 3$ hours is increasing at 50 liters/hour².

2 : $\begin{cases} 1 : r'(3) \\ 1 : \text{meaning of } r'(3) \end{cases}$

(d) $12,000 - \int_0^A r(t) dt = 9000$

2 : $\begin{cases} 1 : \text{integral} \\ 1 : \text{equation} \end{cases}$

Differential Equations

98 21. If $\frac{dy}{dt} = ky$ and k is a nonzero constant, then y could be

- (A) $2e^{kty}$ (B) $2e^{kt}$ (C) $e^{kt} + 3$ (D) $ky + 5$ (E) $\frac{1}{2}ky^2 + \frac{1}{2}$

85 4. If $\frac{dy}{dx} = \cos(2x)$, then $y =$

- (A) $-\frac{1}{2}\cos(2x) + C$ (B) $-\frac{1}{2}\cos^2(2x) + C$ (C) $\frac{1}{2}\sin(2x) + C$
(D) $\frac{1}{2}\sin^2(2x) + C$ (E) $-\frac{1}{2}\sin(2x) + C$

73 36. If $y = e^{nx}$, then $\frac{d^n y}{dx^n} =$

- (A) $n^n e^{nx}$ (B) $n!e^{nx}$ (C) ne^{nx} (D) $n^n e^x$ (E) $n!e^x$

73 37. If $\frac{dy}{dx} = 4y$ and if $y = 4$ when $x = 0$, then $y =$

- (A) $4e^{4x}$ (B) e^{4x} (C) $3 + e^{4x}$ (D) $4 + e^{4x}$ (E) $2x^2 + 4$

88 BC 32. The general solution of the differential equation $y' = y + x^2$ is $y =$

- (A) Ce^x (B) $Ce^x + x^2$ (C) $-x^2 - 2x - 2 + C$
(D) $e^x - x^2 - 2x - 2 + C$ (E) $Ce^x - x^2 - 2x - 2$

85 BC 33. If $\frac{dy}{dt} = -2y$ and if $y = 1$ when $t = 0$, what is the value of t for which $y = \frac{1}{2}$?

- (A) $-\frac{\ln 2}{2}$ (B) $-\frac{1}{4}$ (C) $\frac{\ln 2}{2}$ (D) $\frac{\sqrt{2}}{2}$ (E) $\ln 2$

85 BC 37. The general solution for the equation $\frac{dy}{dx} + y = xe^{-x}$ is

(A) $y = \frac{x^2}{2}e^{-x} + Ce^{-x}$

(B) $y = \frac{x^2}{2}e^{-x} + e^{-x} + C$

(C) $y = -e^{-x} + \frac{C}{1+x}$

(D) $y = xe^{-x} + Ce^{-x}$

(E) $y = C_1e^x + C_2xe^{-x}$

88 BC 39. If $\frac{dy}{dx} = y \sec^2 x$ and $y = 5$ when $x = 0$, then $y =$

(A) $e^{\tan x} + 4$

(B) $e^{\tan x} + 5$

(C) $5e^{\tan x}$

(D) $\tan x + 5$

(E) $\tan x + 5e^x$

69 BC 23. If the graph of $y = f(x)$ contains the point $(0, 2)$, $\frac{dy}{dx} = \frac{-x}{ye^{x^2}}$ and $f(x) > 0$ for all x , then $f(x) =$

(A) $3 + e^{-x^2}$

(B) $\sqrt{3} + e^{-x}$

(C) $1 + e^{-x}$

(D) $\sqrt{3 + e^{-x^2}}$

(E) $\sqrt{3 + e^{x^2}}$

93 BC 13. If $\frac{dy}{dx} = x^2y$, then y could be

(A) $3 \ln\left(\frac{x}{3}\right)$

(B) $e^{\frac{x^3}{3} + 7}$

(C) $2e^{\frac{x^3}{3}}$

(D) $3e^{2x}$

(E) $\frac{x^3}{3} + 1$

98 BC 8. If $\frac{dy}{dx} = \sin x \cos^2 x$ and if $y = 0$ when $x = \frac{\pi}{2}$, what is the value of y when $x = 0$?

(A) -1

(B) $-\frac{1}{3}$

(C) 0

(D) $\frac{1}{3}$

(E) 1

97
BC


83. If $\frac{dy}{dx} = (1 + \ln x)y$ and if $y = 1$ when $x = 1$, then $y =$

- (A) $e^{\frac{x^2-1}{x^2}}$
(B) $1 + \ln x$
(C) $\ln x$
(D) $e^{2x+x \ln x - 2}$
(E) $e^{x \ln x}$

93
BC

39. If $\frac{dy}{dx} = \frac{1}{x}$, then the average rate of change of y with respect to x on the closed interval $[1, 4]$ is

- (A) $-\frac{1}{4}$ (B) $\frac{1}{2} \ln 2$ (C) $\frac{2}{3} \ln 2$ (D) $\frac{2}{5}$ (E) 2



98
BC

24. Shown above is a slope field for which of the following differential equations?

- (A) $\frac{dy}{dx} = 1 + x$ (B) $\frac{dy}{dx} = x^2$ (C) $\frac{dy}{dx} = x + y$ (D) $\frac{dy}{dx} = \frac{x}{y}$ (E) $\frac{dy}{dx} = \ln y$

88
BC

43. Bacteria in a certain culture increase at a rate proportional to the number present. If the number of bacteria doubles in three hours, in how many hours will the number of bacteria triple?

- (A) $\frac{3 \ln 3}{\ln 2}$ (B) $\frac{2 \ln 3}{\ln 2}$ (C) $\frac{\ln 3}{\ln 2}$ (D) $\ln\left(\frac{27}{2}\right)$ (E) $\ln\left(\frac{9}{2}\right)$
-

98
BC

26. The population $P(t)$ of a species satisfies the logistic differential equation $\frac{dP}{dt} = P\left(2 - \frac{P}{5000}\right)$, where the initial population $P(0) = 3,000$ and t is the time in years. What is $\lim_{t \rightarrow \infty} P(t)$?

- (A) 2,500 (B) 3,000 (C) 4,200 (D) 5,000 (E) 10,000
-

98

84. Population y grows according to the equation $\frac{dy}{dt} = ky$, where k is a constant and t is measured in years. If the population doubles every 10 years, then the value of k is

- (A) 0.069 (B) 0.200 (C) 0.301 (D) 3.322 (E) 5.000
-

93
BC

38. During a certain epidemic, the number of people that are infected at any time increases at a rate proportional to the number of people that are infected at that time. If 1,000 people are infected when the epidemic is first discovered, and 1,200 are infected 7 days later, how many people are infected 12 days after the epidemic is first discovered?

- (A) 343 (B) 1,343 (C) 1,367 (D) 1,400 (E) 2,057
-

93
BC

17. The number of bacteria in a culture is growing at a rate of $3,000e^{2t/5}$ per unit of time t . At $t = 0$, the number of bacteria present was 7,500. Find the number present at $t = 5$.

- (A) $1,200e^2$ (B) $3,000e^2$ (C) $7,500e^2$ (D) $7,500e^5$ (E) $\frac{15,000}{7}e^7$


Slope Fields and Diff EQs

6-A

5. Consider the differential equation $\frac{dy}{dx} = \frac{1+y}{x}$, where $x \neq 0$.

(a) On the axes provided, sketch a slope field for the given differential equation at the eight points indicated.

(Note: Use the axes provided in the pink exam booklet.)




(b) Find the particular solution $y = f(x)$ to the differential equation with the initial condition $f(-1) = 1$ and state its domain.

5-A

6. Consider the differential equation $\frac{dy}{dx} = -\frac{2x}{y}$.

(a) On the axes provided, sketch a slope field for the given differential equation at the twelve points indicated.

(Note: Use the axes provided in the pink test booklet.)



(b) Let $y = f(x)$ be the particular solution to the differential equation with the initial condition $f(1) = -1$.


Write an equation for the line tangent to the graph of f at $(1, -1)$ and use it to approximate $f(1.1)$.

(c) Find the particular solution $y = f(x)$ to the given differential equation with the initial condition $f(1) = -1$.

6-8

5. Consider the differential equation $\frac{dy}{dx} = (y - 1)^2 \cos(\pi x)$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the nine points indicated.
 (Note: Use the axes provided in the exam booklet.)




- (b) There is a horizontal line with equation $y = c$ that satisfies this differential equation. Find the value of c .
 (c) Find the particular solution $y = f(x)$ to the differential equation with the initial condition $f(1) = 0$.

4-8

5. Consider the differential equation $\frac{dy}{dx} = x^4(y - 2)$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the twelve points indicated.
 (Note: Use the axes provided in the test booklet.)




- (b) While the slope field in part (a) is drawn at only twelve points, it is defined at every point in the xy -plane. Describe all points in the xy -plane for which the slopes are negative.
 (c) Find the particular solution $y = f(x)$ to the given differential equation with the initial condition $f(0) = 0$.

108

5. Consider the differential equation $\frac{dy}{dx} = \frac{x+1}{y}$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the twelve points indicated, and for $-1 < x < 1$, sketch the solution curve that passes through the point $(0, -1)$.

(Note: Use the axes provided in the exam booklet.)




- (b) While the slope field in part (a) is drawn at only twelve points, it is defined at every point in the xy -plane for which $y \neq 0$. Describe all points in the xy -plane, $y \neq 0$, for which $\frac{dy}{dx} = -1$.
- (c) Find the particular solution $y = f(x)$ to the given differential equation with the initial condition $f(0) = -2$.

8-A

5. Consider the differential equation $\frac{dy}{dx} = \frac{y-1}{x^2}$, where $x \neq 0$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the nine points indicated.
- (Note: Use the axes provided in the exam booklet.)



- (b) Find the particular solution $y = f(x)$ to the differential equation with the initial condition $f(2) = 0$.
- (c) For the particular solution $y = f(x)$ described in part (b), find $\lim_{x \rightarrow \infty} f(x)$.


189

189

5-8

6. Consider the differential equation $\frac{dy}{dx} = \frac{-xy^2}{2}$. Let $y = f(x)$ be the particular solution to this differential equation with the initial condition $f(-1) = 2$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the twelve points indicated. (Note: Use the axes provided in the test booklet.)




- (b) Write an equation for the line tangent to the graph of f at $x = -1$.
(c) Find the solution $y = f(x)$ to the given differential equation with the initial condition $f(-1) = 2$.

4-A

6. Consider the differential equation $\frac{dy}{dx} = x^2(y - 1)$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the twelve points indicated. (Note: Use the axes provided in the pink test booklet.)




- (b) While the slope field in part (a) is drawn at only twelve points, it is defined at every point in the xy -plane. Describe all points in the xy -plane for which the slopes are positive.
(c) Find the particular solution $y = f(x)$ to the given differential equation with the initial condition $f(0) = 3$.

7-8

5. Consider the differential equation $\frac{dy}{dx} = \frac{1}{2}x + y - 1$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the nine points indicated.
(Note: Use the axes provided in the exam booklet.)



- (b) Find $\frac{d^2y}{dx^2}$ in terms of x and y . Describe the region in the xy -plane in which all solution curves to the differential equation are concave up.
- (c) Let $y = f(x)$ be a particular solution to the differential equation with the initial condition $f(0) = 1$. Does f have a relative minimum, a relative maximum, or neither at $x = 0$? Justify your answer.
- (d) Find the values of the constants m and b , for which $y = mx + b$ is a solution to the differential equation.

2-0

5. Consider the differential equation $\frac{dy}{dx} = \frac{3-x}{y}$.

- (a) Let $y = f(x)$ be the particular solution to the given differential equation for $1 < x < 5$ such that the line $y = -2$ is tangent to the graph of f . Find the x -coordinate of the point of tangency, and determine whether f has a local maximum, local minimum, or neither at this point. Justify your answer.
- (b) Let $y = g(x)$ be the particular solution to the given differential equation for $-2 < x < 8$, with the initial condition $g(6) = -4$. Find $y = g(x)$.

2011

5. At the beginning of 2010, a landfill contained 1400 tons of solid waste. The increasing function W models the total amount of solid waste stored at the landfill. Planners estimate that W will satisfy the differential equation $\frac{dW}{dt} = \frac{1}{25}(W - 300)$ for the next 20 years. W is measured in tons, and t is measured in years from the start of 2010.
- (a) Use the line tangent to the graph of W at $t = 0$ to approximate the amount of solid waste that the landfill contains at the end of the first 3 months of 2010 (time $t = \frac{1}{4}$).
 - (b) Find $\frac{d^2W}{dt^2}$ in terms of W . Use $\frac{d^2W}{dt^2}$ to determine whether your answer in part (a) is an underestimate or an overestimate of the amount of solid waste that the landfill contains at time $t = \frac{1}{4}$.
 - (c) Find the particular solution $W = W(t)$ to the differential equation $\frac{dW}{dt} = \frac{1}{25}(W - 300)$ with initial condition $W(0) = 1400$.
-

3-9

6. Let f be the function satisfying $f'(x) = x\sqrt{f(x)}$ for all real numbers x , where $f(3) = 25$.
- (a) Find $f''(3)$.
 - (b) Write an expression for $y = f(x)$ by solving the differential equation $\frac{dy}{dx} = x\sqrt{y}$ with the initial condition $f(3) = 25$.
-

98

4. Let f be a function with $f(1) = 4$ such that for all points (x, y) on the graph of f the slope is given by $\frac{3x^2 + 1}{2y}$.
- (a) Find the slope of the graph of f at the point where $x = 1$.
 - (b) Write an equation for the line tangent to the graph of f at $x = 1$ and use it to approximate $f(1.2)$.
 - (c) Find $f(x)$ by solving the separable differential equation $\frac{dy}{dx} = \frac{3x^2 + 1}{2y}$ with the initial condition $f(1) = 4$.
 - (d) Use your solution from part (c) to find $f(1.2)$.

00

6. Consider the differential equation $\frac{dy}{dx} = \frac{3x^2}{e^{2y}}$.

- (a) Find a solution $y = f(x)$ to the differential equation satisfying $f(0) = \frac{1}{2}$.
- (b) Find the domain and range of the function f found in part (a).

10-A 6. Solutions to the differential equation $\frac{dy}{dx} = xy^3$ also satisfy $\frac{d^2y}{dx^2} = y^3(1 + 3x^2y^2)$. Let $y = f(x)$ be a particular solution to the differential equation $\frac{dy}{dx} = xy^3$ with $f(1) = 2$.


- Write an equation for the line tangent to the graph of $y = f(x)$ at $x = 1$.
- Use the tangent line equation from part (a) to approximate $f(1.1)$. Given that $f(x) > 0$ for $1 < x < 1.1$, is the approximation for $f(1.1)$ greater than or less than $f(1.1)$? Explain your reasoning.
- Find the particular solution $y = f(x)$ with initial condition $f(1) = 2$.

12-A 5. The rate at which a baby bird gains weight is proportional to the difference between its adult weight and its current weight. At time $t = 0$, when the bird is first weighed, its weight is 20 grams. If $B(t)$ is the weight of the bird, in grams, at time t days after it is first weighed, then

$$\frac{dB}{dt} = \frac{1}{5}(100 - B).$$

Let $y = B(t)$ be the solution to the differential equation above with initial condition $B(0) = 20$.

- Is the bird gaining weight faster when it weighs 40 grams or when it weighs 70 grams? Explain your reasoning.
- Find $\frac{d^2B}{dt^2}$ in terms of B . Use $\frac{d^2B}{dt^2}$ to explain why the graph of B cannot resemble the following graph.




- Use separation of variables to find $y = B(t)$, the particular solution to the differential equation with initial condition $B(0) = 20$.

**AP[®] CALCULUS AB
2006 SCORING GUIDELINES**

Question 5


Consider the differential equation $\frac{dy}{dx} = \frac{1+y}{x}$, where $x \neq 0$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the eight points indicated.
(Note: Use the axes provided in the pink exam booklet.)



- (b) Find the particular solution $y = f(x)$ to the differential equation with the initial condition $f(-1) = 1$ and state its domain.

(a)



2 : sign of slope at each point and relative steepness of slope lines in rows and columns

(b) $\frac{1}{1+y} dy = \frac{1}{x} dx$

$$\ln|1+y| = \ln|x| + K$$

$$|1+y| = e^{\ln|x|+K}$$

$$1+y = C|x|$$

$$2 = C$$

$$1+y = 2|x|$$

$$y = 2|x| - 1 \text{ and } x < 0$$

or

$$y = -2x - 1 \text{ and } x < 0$$

6 : { 1 : separates variables
2 : antiderivatives
1 : constant of integration
1 : uses initial condition
1 : solves for y

7 : Note: max 3/6 [1-2-0-0-0] if no constant of integration
Note: 0/6 if no separation of variables


1 : domain

AP[®] CALCULUS AB
2005 SCORING GUIDELINES


Question 6

Consider the differential equation $\frac{dy}{dx} = -\frac{2x}{y}$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the twelve points indicated.
(Note: Use the axes provided in the pink test booklet.)
- (b) Let $y = f(x)$ be the particular solution to the differential equation with the initial condition $f(1) = -1$. Write an equation for the line tangent to the graph of f at $(1, -1)$ and use it to approximate $f(1.1)$.
- (c) Find the particular solution $y = f(x)$ to the given differential equation with the initial condition $f(1) = -1$.



(a)



2 : $\begin{cases} 1 : \text{zero slopes} \\ 1 : \text{nonzero slopes} \end{cases}$

- (b) The line tangent to f at $(1, -1)$ is $y + 1 = 2(x - 1)$.
Thus, $f(1.1)$ is approximately -0.8 .

2 : $\begin{cases} 1 : \text{equation of the tangent line} \\ 1 : \text{approximation for } f(1.1) \end{cases}$

- (c) $\frac{dy}{dx} = -\frac{2x}{y}$
 $y \, dy = -2x \, dx$
 $\frac{y^2}{2} = -x^2 + C$
 $\frac{1}{2} = -1 + C; C = \frac{3}{2}$
 $y^2 = -2x^2 + 3$
 Since the particular solution goes through $(1, -1)$,
 y must be negative.
 Thus the particular solution is $y = -\sqrt{3 - 2x^2}$.

5 : $\begin{cases} 1 : \text{separates variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } y \end{cases}$


Note: max 2/5 [1-1-0-0-0] if no constant of integration
 Note: 0/5 if no separation of variables

AP[®] CALCULUS AB
2006 SCORING GUIDELINES (Form B)

Question 5


Consider the differential equation $\frac{dy}{dx} = (y - 1)^2 \cos(\pi x)$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the nine points indicated.
(Note: Use the axes provided in the exam booklet.)



- (b) There is a horizontal line with equation $y = c$ that satisfies this differential equation. Find the value of c .
(c) Find the particular solution $y = f(x)$ to the differential equation with the initial condition $f(1) = 0$.

(a)



- (b) The line $y = 1$ satisfies the differential equation, so $c = 1$.

(c) $\frac{1}{(y-1)^2} dy = \cos(\pi x) dx$

$$-(y-1)^{-1} = \frac{1}{\pi} \sin(\pi x) + C$$

$$\frac{1}{1-y} = \frac{1}{\pi} \sin(\pi x) + C$$

$$1 = \frac{1}{\pi} \sin(\pi) + C = C$$

$$\frac{1}{1-y} = \frac{1}{\pi} \sin(\pi x) + 1$$

$$\frac{\pi}{1-y} = \sin(\pi x) + \pi$$

$$y = 1 - \frac{\pi}{\sin(\pi x) + \pi} \text{ for } -\infty < x < \infty$$

2 : $\begin{cases} 1 : \text{zero slopes} \\ 1 : \text{all other slopes} \end{cases}$

1 : $c = 1$

6 : $\begin{cases} 1 : \text{separates variables} \\ 2 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{answer} \end{cases}$

Note: max 3/6 [1-2-0-0-0] if no constant of integration



Note: 0/6 if no separation of variables

**AP[®] CALCULUS AB
2004 SCORING GUIDELINES (Form B)**

Question 5

Consider the differential equation $\frac{dy}{dx} = x^4(y - 2)$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the twelve points indicated.
(Note: Use the axes provided in the test booklet.)
- (b) While the slope field in part (a) is drawn at only twelve points, it is defined at every point in the xy -plane. Describe all points in the xy -plane for which the slopes are negative.
- (c) Find the particular solution $y = f(x)$ to the given differential equation with the initial condition $f(0) = 0$.



- 1 : zero slope at each point (x, y)
where $x = 0$ or $y = 2$
- 2 : { positive slope at each point (x, y)
where $x \neq 0$ and $y > 2$
- 1 : { negative slope at each point (x, y)
where $x \neq 0$ and $y < 2$

- (b) Slopes are negative at points (x, y)
where $x \neq 0$ and $y < 2$.

1 : description

(c) $\frac{1}{y-2} dy = x^4 dx$
 $\ln|y-2| = \frac{1}{5}x^5 + C$
 $|y-2| = e^C e^{\frac{1}{5}x^5}$
 $y-2 = Ke^{\frac{1}{5}x^5}, K = \pm e^C$
 $-2 = Ke^0 = K$
 $y = 2 - 2e^{\frac{1}{5}x^5}$

- 6 : { 1 : separates variables
2 : antiderivatives
1 : constant of integration
1 : uses initial condition
1 : solves for y
0/1 if y is not exponential

Note: max 3/6 [1-2-0-0-0] if no
constant of integration

Note: 0/6 if no separation of variables

AP[®] CALCULUS AB
2010 SCORING GUIDELINES (Form B)


Question 5

Consider the differential equation $\frac{dy}{dx} = \frac{x+1}{y}$.


- (a) On the axes provided, sketch a slope field for the given differential equation at the twelve points indicated, and for $-1 < x < 1$, sketch the solution curve that passes through the point $(0, -1)$.

(Note: Use the axes provided in the exam booklet.)

- (b) While the slope field in part (a) is drawn at only twelve points, it is defined at every point in the xy -plane for which $y \neq 0$. Describe all points in the xy -plane, $y \neq 0$, for which $\frac{dy}{dx} = -1$.
- (c) Find the particular solution $y = f(x)$ to the given differential equation with the initial condition $f(0) = -2$.



(a)



3 : { 1 : zero slopes
 1 : nonzero slopes
 1 : solution curve through $(0, -1)$

(b) $-1 = \frac{x+1}{y} \Rightarrow y = -x - 1$

$\frac{dy}{dx} = -1$ for all (x, y) with $y = -x - 1$ and $y \neq 0$

(c) $\int y \, dy = \int (x+1) \, dx$

$\frac{y^2}{2} = \frac{x^2}{2} + x + C$

$\frac{(-2)^2}{2} = \frac{0^2}{2} + 0 + C \Rightarrow C = 2$

$y^2 = x^2 + 2x + 4$

Since the solution goes through $(0, -2)$, y must be negative. Therefore $y = -\sqrt{x^2 + 2x + 4}$.

1 : description

5 : { 1 : separates variables
 1 : antiderivatives
 1 : constant of integration
 1 : uses initial condition
 1 : solves for y

Note: max 2/5 [1-1-0-0-0] if no constant of integration

Note: 0/5 if no separation of variables

**AP[®] CALCULUS AB
2008 SCORING GUIDELINES**

Question 5


Consider the differential equation $\frac{dy}{dx} = \frac{y-1}{x^2}$, where $x \neq 0$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the nine points indicated.


(Note: Use the axes provided in the exam booklet.)

- (b) Find the particular solution $y = f(x)$ to the differential equation with the initial condition $f(2) = 0$.

- (c) For the particular solution $y = f(x)$ described in part (b), find $\lim_{x \rightarrow \infty} f(x)$.



(a)



2 : $\begin{cases} 1 : \text{zero slopes} \\ 1 : \text{all other slopes} \end{cases}$

(b) $\frac{1}{y-1} dy = \frac{1}{x^2} dx$

$$\ln|y-1| = -\frac{1}{x} + C$$

$$|y-1| = e^{-\frac{1}{x} + C}$$

$$|y-1| = e^C e^{-\frac{1}{x}}$$

$$y-1 = ke^{-\frac{1}{x}}, \text{ where } k = \pm e^C$$

$$-1 = ke^{-\frac{1}{2}}$$

$$k = -e^{\frac{1}{2}}$$

$$f(x) = 1 - e^{\left(\frac{1}{2} - \frac{1}{x}\right)}, x > 0$$

6 : $\begin{cases} 1 : \text{separates variables} \\ 2 : \text{antidifferentiates} \\ 1 : \text{includes constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } y \end{cases}$

Note: max 3/6 [1-2-0-0-0] if no constant of integration

Note: 0/6 if no separation of variables


(c) $\lim_{x \rightarrow \infty} 1 - e^{\left(\frac{1}{2} - \frac{1}{x}\right)} = 1 - \sqrt{e}$

1 : limit


**AP[®] CALCULUS AB
2005 SCORING GUIDELINES (Form B)**

Question 6

Consider the differential equation $\frac{dy}{dx} = \frac{-xy^2}{2}$. Let $y = f(x)$ be the particular solution to this differential equation with the initial condition $f(-1) = 2$.



- (a) On the axes provided, sketch a slope field for the given differential equation at the twelve points indicated.
(Note: Use the axes provided in the test booklet.)
- (b) Write an equation for the line tangent to the graph of f at $x = -1$.
- (c) Find the solution $y = f(x)$ to the given differential equation with the initial condition $f(-1) = 2$.



2 : $\begin{cases} 1 : \text{zero slopes} \\ 1 : \text{nonzero slopes} \end{cases}$

(b) Slope = $\frac{-(-1)4}{2} = 2$
 $y - 2 = 2(x + 1)$

1 : equation

(c) $\frac{1}{y^2} dy = -\frac{x}{2} dx$
 $-\frac{1}{y} = -\frac{x^2}{4} + C$
 $-\frac{1}{2} = -\frac{1}{4} + C; C = -\frac{1}{4}$
 $y = \frac{1}{\frac{x^2}{4} + \frac{1}{4}} = \frac{4}{x^2 + 1}$

6 : $\begin{cases} 1 : \text{separates variables} \\ 2 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } y \end{cases}$



Note: max 3/6 [1-2-0-0-0] if no constant of integration
 Note: 0/6 if no separation of variables

**AP[®] CALCULUS AB
2004 SCORING GUIDELINES**

Question 6

Consider the differential equation $\frac{dy}{dx} = x^2(y - 1)$.

- (a) On the axes provided, sketch a slope field for the given differential equation at the twelve points indicated.
(Note: Use the axes provided in the pink test booklet.)
- (b) While the slope field in part (a) is drawn at only twelve points, it is defined at every point in the xy -plane. Describe all points in the xy -plane for which the slopes are positive.
- (c) Find the particular solution $y = f(x)$ to the given differential equation with the initial condition $f(0) = 3$.



- (b) Slopes are positive at points (x, y) where $x \neq 0$ and $y > 1$.

- 1 : zero slope at each point (x, y) where $x = 0$ or $y = 1$
- 2 : { positive slope at each point (x, y) where $x \neq 0$ and $y > 1$
- 1 : { negative slope at each point (x, y) where $x \neq 0$ and $y < 1$

1 : description

(c) $\frac{1}{y-1} dy = x^2 dx$
 $\ln|y-1| = \frac{1}{3}x^3 + C$
 $|y-1| = e^C e^{\frac{1}{3}x^3}$
 $y-1 = Ke^{\frac{1}{3}x^3}, K = \pm e^C$
 $2 = Ke^0 = K$
 $y = 1 + 2e^{\frac{1}{3}x^3}$

- 1 : separates variables
 2 : antiderivatives
 1 : constant of integration
 6 : { 1 : uses initial condition
 1 : solves for y
 0/1 if y is not exponential

Note: max 3/6 [1-2-0-0-0] if no constant of integration

Note: 0/6 if no separation of variables

AP[®] CALCULUS AB
2007 SCORING GUIDELINES (Form B)

Question 5

Consider the differential equation $\frac{dy}{dx} = \frac{1}{2}x + y - 1$.


- (a) On the axes provided, sketch a slope field for the given differential equation at the nine points indicated.

(Note: Use the axes provided in the exam booklet.)


- (b) Find $\frac{d^2y}{dx^2}$ in terms of x and y . Describe the region in the xy -plane in which all solution curves to the differential equation are concave up.

- (c) Let $y = f(x)$ be a particular solution to the differential equation with the initial condition $f(0) = 1$. Does f have a relative minimum, a relative maximum, or neither at $x = 0$? Justify your answer.

- (d) Find the values of the constants m and b , for which $y = mx + b$ is a solution to the differential equation.



(a)



2 : Sign of slope at each point and relative steepness of slope lines in rows and columns.

(b) $\frac{d^2y}{dx^2} = \frac{1}{2} + \frac{dy}{dx} = \frac{1}{2}x + y - \frac{1}{2}$

Solution curves will be concave up on the half-plane above the line

$$y = -\frac{1}{2}x + \frac{1}{2}.$$

3 : $\left\{ \begin{array}{l} 2 : \frac{d^2y}{dx^2} \\ 1 : \text{description} \end{array} \right.$

(c) $\left. \frac{dy}{dx} \right|_{(0,1)} = 0 + 1 - 1 = 0$ and $\left. \frac{d^2y}{dx^2} \right|_{(0,1)} = 0 + 1 - \frac{1}{2} > 0$

Thus, f has a relative minimum at $(0, 1)$.

2 : $\left\{ \begin{array}{l} 1 : \text{answer} \\ 1 : \text{justification} \end{array} \right.$

- (d) Substituting $y = mx + b$ into the differential equation:

$$m = \frac{1}{2}x + (mx + b) - 1 = \left(m + \frac{1}{2}\right)x + (b - 1)$$

Then $0 = m + \frac{1}{2}$ and $m = b - 1$: $m = -\frac{1}{2}$ and $b = \frac{1}{2}$.

2 : $\left\{ \begin{array}{l} 1 : \text{value for } m \\ 1 : \text{value for } b \end{array} \right.$

AP[®] CALCULUS AB
2002 SCORING GUIDELINES (Form B)

Question 5

Consider the differential equation $\frac{dy}{dx} = \frac{3-x}{y}$.

- (a) Let $y = f(x)$ be the particular solution to the given differential equation for $1 < x < 5$ such that the line $y = -2$ is tangent to the graph of f . Find the x -coordinate of the point of tangency, and determine whether f has a local maximum, local minimum, or neither at this point. Justify your answer.
- (b) Let $y = g(x)$ be the particular solution to the given differential equation for $-2 < x < 8$, with the initial condition $g(6) = -4$. Find $y = g(x)$.

(a) $\frac{dy}{dx} = 0$ when $x = 3$

$$\left. \frac{d^2y}{dx^2} \right|_{(3,-2)} = \left. \frac{-y - y'(3-x)}{y^2} \right|_{(3,-2)} = \frac{1}{2},$$

so f has a local minimum at this point.

or

Because f is continuous for $1 < x < 5$, there is an interval containing $x = 3$ on which

$y < 0$. On this interval, $\frac{dy}{dx}$ is negative to

the left of $x = 3$ and $\frac{dy}{dx}$ is positive to the

right of $x = 3$. Therefore f has a local

minimum at $x = 3$.

$$3 \left\{ \begin{array}{l} 1 : x = 3 \\ 1 : \text{local minimum} \\ 1 : \text{justification} \end{array} \right.$$

(b) $y \, dy = (3-x) \, dx$

$$\frac{1}{2}y^2 = 3x - \frac{1}{2}x^2 + C$$

$$8 = 18 - 18 + C; C = 8$$

$$y^2 = 6x - x^2 + 16$$

$$y = -\sqrt{6x - x^2 + 16}$$

$$6 \left\{ \begin{array}{l} 1 : \text{separates variables} \\ 1 : \text{antiderivative of } dy \text{ term} \\ 1 : \text{antiderivative of } dx \text{ term} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition } g(6) = -4 \\ 1 : \text{solves for } y \end{array} \right.$$

Note: max 3/6 [1-1-1-0-0-0] if no constant of integration

Note: 0/6 if no separation of variables

AP[®] CALCULUS AB
2011 SCORING GUIDELINES

Question 5

At the beginning of 2010, a landfill contained 1400 tons of solid waste. The increasing function W models the total amount of solid waste stored at the landfill. Planners estimate that W will satisfy the differential equation $\frac{dW}{dt} = \frac{1}{25}(W - 300)$ for the next 20 years. W is measured in tons, and t is measured in years from the start of 2010.

- (a) Use the line tangent to the graph of W at $t = 0$ to approximate the amount of solid waste that the landfill contains at the end of the first 3 months of 2010 (time $t = \frac{1}{4}$).
- (b) Find $\frac{d^2W}{dt^2}$ in terms of W . Use $\frac{d^2W}{dt^2}$ to determine whether your answer in part (a) is an underestimate or an overestimate of the amount of solid waste that the landfill contains at time $t = \frac{1}{4}$.
- (c) Find the particular solution $W = W(t)$ to the differential equation $\frac{dW}{dt} = \frac{1}{25}(W - 300)$ with initial condition $W(0) = 1400$.

(a) $\left. \frac{dW}{dt} \right|_{t=0} = \frac{1}{25}(W(0) - 300) = \frac{1}{25}(1400 - 300) = 44$

The tangent line is $y = 1400 + 44t$.

$$W\left(\frac{1}{4}\right) \approx 1400 + 44\left(\frac{1}{4}\right) = 1411 \text{ tons}$$

$$2 : \begin{cases} 1 : \frac{dW}{dt} \text{ at } t = 0 \\ 1 : \text{answer} \end{cases}$$

(b) $\frac{d^2W}{dt^2} = \frac{1}{25} \frac{dW}{dt} = \frac{1}{625}(W - 300)$ and $W \geq 1400$

Therefore $\frac{d^2W}{dt^2} > 0$ on the interval $0 \leq t \leq \frac{1}{4}$.

The answer in part (a) is an underestimate.

$$2 : \begin{cases} 1 : \frac{d^2W}{dt^2} \\ 1 : \text{answer with reason} \end{cases}$$

(c) $\frac{dW}{dt} = \frac{1}{25}(W - 300)$

$$\int \frac{1}{W - 300} dW = \int \frac{1}{25} dt$$

$$\ln|W - 300| = \frac{1}{25}t + C$$

$$\ln(1400 - 300) = \frac{1}{25}(0) + C \Rightarrow \ln(1100) = C$$

$$W - 300 = 1100e^{\frac{1}{25}t}$$

$$W(t) = 300 + 1100e^{\frac{1}{25}t}, \quad 0 \leq t \leq 20$$

$$5 : \begin{cases} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } W \end{cases}$$

Note: max 2/5 [1-1-0-0-0] if no constant of integration

Note: 0/5 if no separation of variables

AP[®] CALCULUS AB
2003 SCORING GUIDELINES (Form B)

Question 6

Let f be the function satisfying $f'(x) = x\sqrt{f(x)}$ for all real numbers x , where $f(3) = 25$.

- (a) Find $f''(3)$.
- (b) Write an expression for $y = f(x)$ by solving the differential equation $\frac{dy}{dx} = x\sqrt{y}$ with the initial condition $f(3) = 25$.

(a) $f''(x) = \sqrt{f(x)} + x \cdot \frac{f'(x)}{2\sqrt{f(x)}} = \sqrt{f(x)} + \frac{x^2}{2}$

$$f''(3) = \sqrt{25} + \frac{9}{2} = \frac{19}{2}$$

(b) $\frac{1}{\sqrt{y}} dy = x dx$

$$2\sqrt{y} = \frac{1}{2}x^2 + C$$

$$2\sqrt{25} = \frac{1}{2}(3)^2 + C; C = \frac{11}{2}$$

$$\sqrt{y} = \frac{1}{4}x^2 + \frac{11}{4}$$

$$y = \left(\frac{1}{4}x^2 + \frac{11}{4}\right)^2 = \frac{1}{16}(x^2 + 11)^2$$

3 : $\left\{ \begin{array}{l} 2 : f''(x) \\ < -2 > \text{ product or} \\ & \text{chain rule error} \\ 1 : \text{ value at } x = 3 \end{array} \right.$

6 : $\left\{ \begin{array}{l} 1 : \text{ separates variables} \\ 1 : \text{ antiderivative of } dy \text{ term} \\ 1 : \text{ antiderivative of } dx \text{ term} \\ 1 : \text{ constant of integration} \\ 1 : \text{ uses initial condition } f(3) = 25 \\ 1 : \text{ solves for } y \end{array} \right.$

Note: max 3/6 [1-1-1-0-0-0] if no constant of integration

Note: 0/6 if no separation of variables

AP[®] CALCULUS AB
2010 SCORING GUIDELINES

Question 6

Solutions to the differential equation $\frac{dy}{dx} = xy^3$ also satisfy $\frac{d^2y}{dx^2} = y^3(1 + 3x^2y^2)$. Let $y = f(x)$ be a particular solution to the differential equation $\frac{dy}{dx} = xy^3$ with $f(1) = 2$.

- (a) Write an equation for the line tangent to the graph of $y = f(x)$ at $x = 1$.
- (b) Use the tangent line equation from part (a) to approximate $f(1.1)$. Given that $f(x) > 0$ for $1 < x < 1.1$, is the approximation for $f(1.1)$ greater than or less than $f(1.1)$? Explain your reasoning.
- (c) Find the particular solution $y = f(x)$ with initial condition $f(1) = 2$.

(a) $f'(1) = \left. \frac{dy}{dx} \right|_{(1,2)} = 8$

An equation of the tangent line is $y = 2 + 8(x - 1)$.

(b) $f(1.1) \approx 2.8$

Since $y = f(x) > 0$ on the interval $1 \leq x < 1.1$,

$$\frac{d^2y}{dx^2} = y^3(1 + 3x^2y^2) > 0 \text{ on this interval.}$$

Therefore on the interval $1 < x < 1.1$, the line tangent to the graph of $y = f(x)$ at $x = 1$ lies below the curve and the approximation 2.8 is less than $f(1.1)$.

(c) $\frac{dy}{dx} = xy^3$

$$\int \frac{1}{y^3} dy = \int x dx$$

$$-\frac{1}{2y^2} = \frac{x^2}{2} + C$$

$$-\frac{1}{2 \cdot 2^2} = \frac{1^2}{2} + C \Rightarrow C = -\frac{5}{8}$$

$$y^2 = \frac{1}{\frac{5}{4} - x^2}$$

$$f(x) = \frac{2}{\sqrt{5 - 4x^2}}, \quad \frac{-\sqrt{5}}{2} < x < \frac{\sqrt{5}}{2}$$

2 : $\begin{cases} 1 : f'(1) \\ 1 : \text{answer} \end{cases}$

2 : $\begin{cases} 1 : \text{approximation} \\ 1 : \text{conclusion with explanation} \end{cases}$

5 : $\begin{cases} 1 : \text{separation of variables} \\ 1 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } y \end{cases}$

Note: max 2/5 [1-1-0-0-0] if no constant of integration

Note: 0/5 if no separation of variables